

Section 1.2

Row Reduction and Echelon Forms

Row Echelon Form

Let's come up with an *algorithm* for turning an arbitrary matrix into a “solved” matrix.

Row Echelon Form

Let's come up with an *algorithm* for turning an arbitrary matrix into a “solved” matrix. What do we mean by “solved”?

Row Echelon Form

Let's come up with an *algorithm* for turning an arbitrary matrix into a “solved” matrix. What do we mean by “solved”?

A matrix is in **row echelon form** if

Row Echelon Form

Let's come up with an *algorithm* for turning an arbitrary matrix into a “solved” matrix. What do we mean by “solved”?

A matrix is in **row echelon form** if

1. All zero rows are at the bottom.

Row Echelon Form

Let's come up with an *algorithm* for turning an arbitrary matrix into a “solved” matrix. What do we mean by “solved”?

A matrix is in **row echelon form** if

1. All zero rows are at the bottom.
2. Each leading nonzero entry of a row is to the *right* of the leading entry of the row above.

Row Echelon Form

Let's come up with an *algorithm* for turning an arbitrary matrix into a “solved” matrix. What do we mean by “solved”?

A matrix is in **row echelon form** if

1. All zero rows are at the bottom.
2. Each leading nonzero entry of a row is to the *right* of the leading entry of the row above.
3. Below a leading entry of a row, all entries are *zero*.

Row Echelon Form

Let's come up with an *algorithm* for turning an arbitrary matrix into a “solved” matrix. What do we mean by “solved”?

A matrix is in **row echelon form** if

1. All zero rows are at the bottom.
2. Each leading nonzero entry of a row is to the *right* of the leading entry of the row above.
3. Below a leading entry of a row, all entries are *zero*.

Picture:

$$\begin{pmatrix} \boxed{\star} & \star & \star & \star & \star \\ 0 & \boxed{\star} & \star & \star & \star \\ 0 & 0 & 0 & \boxed{\star} & \star \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

\star = any number

$\boxed{\star}$ = any nonzero number

Row Echelon Form

Let's come up with an *algorithm* for turning an arbitrary matrix into a “solved” matrix. What do we mean by “solved”?

A matrix is in **row echelon form** if

1. All zero rows are at the bottom.
2. Each leading nonzero entry of a row is to the *right* of the leading entry of the row above.
3. Below a leading entry of a row, all entries are *zero*.

Picture:

$$\begin{pmatrix} \boxed{\star} & \star & \star & \star & \star \\ 0 & \boxed{\star} & \star & \star & \star \\ 0 & 0 & 0 & \boxed{\star} & \star \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

\star = any number

$\boxed{\star}$ = any nonzero number

Definition

A **pivot** $\boxed{\star}$ is the first nonzero entry of a row of a matrix in row echelon form.

Reduced Row Echelon Form

A matrix is in **reduced row echelon form** if it is in row echelon form, and in addition,

Reduced Row Echelon Form

A matrix is in **reduced row echelon form** if it is in row echelon form, and in addition,

4. The pivot in each nonzero row is equal to 1.

Reduced Row Echelon Form

A matrix is in **reduced row echelon form** if it is in row echelon form, and in addition,

4. The pivot in each nonzero row is equal to 1.
5. Each pivot is the only nonzero entry in its column.

Reduced Row Echelon Form

A matrix is in **reduced row echelon form** if it is in row echelon form, and in addition,

4. The pivot in each nonzero row is equal to 1.
5. Each pivot is the only nonzero entry in its column.

Picture:

$$\begin{pmatrix} \color{red}{1} & 0 & \star & 0 & \star \\ 0 & \color{red}{1} & \star & 0 & \star \\ 0 & 0 & 0 & \color{red}{1} & \star \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

$\star =$ any number

$\color{red}{1} =$ pivot

Reduced Row Echelon Form

A matrix is in **reduced row echelon form** if it is in row echelon form, and in addition,

4. The pivot in each nonzero row is equal to 1.
5. Each pivot is the only nonzero entry in its column.

Picture:

$$\begin{pmatrix} \color{red}{1} & 0 & \star & 0 & \star \\ 0 & \color{red}{1} & \star & 0 & \star \\ 0 & 0 & 0 & \color{red}{1} & \star \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \quad \begin{array}{l} \star = \text{any number} \\ \color{red}{1} = \text{pivot} \end{array}$$

Note: Echelon forms do not care whether or not a column is augmented. Just ignore the vertical line.

Reduced Row Echelon Form

A matrix is in **reduced row echelon form** if it is in row echelon form, and in addition,

4. The pivot in each nonzero row is equal to 1.
5. Each pivot is the only nonzero entry in its column.

Picture:

$$\begin{pmatrix} \color{red}{1} & 0 & \star & 0 & \star \\ 0 & \color{red}{1} & \star & 0 & \star \\ 0 & 0 & 0 & \color{red}{1} & \star \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \quad \begin{array}{l} \star = \text{any number} \\ \color{red}{1} = \text{pivot} \end{array}$$

Note: Echelon forms do not care whether or not a column is augmented. Just ignore the vertical line.

Question

Can every matrix be put into reduced row echelon form only using row operations?

Reduced Row Echelon Form

A matrix is in **reduced row echelon form** if it is in row echelon form, and in addition,

4. The pivot in each nonzero row is equal to 1.
5. Each pivot is the only nonzero entry in its column.

Picture:

$$\begin{pmatrix} \color{red}{1} & 0 & \star & 0 & \star \\ 0 & \color{red}{1} & \star & 0 & \star \\ 0 & 0 & 0 & \color{red}{1} & \star \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \quad \begin{array}{l} \star = \text{any number} \\ \color{red}{1} = \text{pivot} \end{array}$$

Note: Echelon forms do not care whether or not a column is augmented. Just ignore the vertical line.

Question

Can every matrix be put into reduced row echelon form only using row operations?

Answer: Yes! Stay tuned.

Reduced Row Echelon Form

Continued

Why is this the “solved” version of the matrix?

Reduced Row Echelon Form

Continued

Why is this the “solved” version of the matrix?

$$\left(\begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & -2 \\ 0 & 0 & 1 & 3 \end{array} \right)$$

is in reduced row echelon form. It translates into

Reduced Row Echelon Form

Continued

Why is this the “solved” version of the matrix?

$$\left(\begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & -2 \\ 0 & 0 & 1 & 3 \end{array} \right)$$

is in reduced row echelon form. It translates into

which is clearly the solution.

Reduced Row Echelon Form

Continued

Why is this the “solved” version of the matrix?

$$\left(\begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & -2 \\ 0 & 0 & 1 & 3 \end{array} \right)$$

is in reduced row echelon form. It translates into

which is clearly the solution.

But what happens if there are fewer pivots than rows?

Reduced Row Echelon Form

Continued

Why is this the “solved” version of the matrix?

$$\left(\begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & -2 \\ 0 & 0 & 1 & 3 \end{array} \right)$$

is in reduced row echelon form. It translates into

which is clearly the solution.

But what happens if there are fewer pivots than rows? ... parametrized solution set (later).

Poll

Which of the following matrices are in reduced row echelon form?

A. $\begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}$ B. $\begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$

C. $\begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}$

D. $(0 \ 1 \ 0 \ 0)$

E. $(0 \ 1 \ 8 \ 0)$

F. $\left(\begin{array}{cc|c} 1 & 17 & 0 \\ 0 & 0 & 1 \end{array} \right)$

Reduced Row Echelon Form

Theorem

Every matrix is row equivalent to one and only one matrix in reduced row echelon form.

Reduced Row Echelon Form

Theorem

Every matrix is row equivalent to one and only one matrix in reduced row echelon form.

We'll give an algorithm, called **row reduction**, which demonstrates that every matrix is row equivalent to *at least one* matrix in reduced row echelon form.

Reduced Row Echelon Form

Theorem

Every matrix is row equivalent to one and only one matrix in reduced row echelon form.

We'll give an algorithm, called **row reduction**, which demonstrates that every matrix is row equivalent to *at least one* matrix in reduced row echelon form.

Note: Like echelon forms, the row reduction algorithm does not care if a column is augmented: ignore the vertical line when row reducing.

Reduced Row Echelon Form

Theorem

Every matrix is row equivalent to one and only one matrix in reduced row echelon form.

We'll give an algorithm, called **row reduction**, which demonstrates that every matrix is row equivalent to *at least one* matrix in reduced row echelon form.

Note: Like echelon forms, the row reduction algorithm does not care if a column is augmented: ignore the vertical line when row reducing.

The uniqueness statement is interesting—

Reduced Row Echelon Form

Theorem

Every matrix is row equivalent to one and only one matrix in reduced row echelon form.

We'll give an algorithm, called **row reduction**, which demonstrates that every matrix is row equivalent to *at least one* matrix in reduced row echelon form.

Note: Like echelon forms, the row reduction algorithm does not care if a column is augmented: ignore the vertical line when row reducing.

The uniqueness statement is interesting—it means that, no matter *how* you row reduce, you *always* get the same matrix in reduced row echelon form.

Reduced Row Echelon Form

Theorem

Every matrix is row equivalent to one and only one matrix in reduced row echelon form.

We'll give an algorithm, called **row reduction**, which demonstrates that every matrix is row equivalent to *at least one* matrix in reduced row echelon form.

Note: Like echelon forms, the row reduction algorithm does not care if a column is augmented: ignore the vertical line when row reducing.

The uniqueness statement is interesting—it means that, no matter *how* you row reduce, you *always* get the same matrix in reduced row echelon form. (Assuming you only do the three legal row operations.)

Reduced Row Echelon Form

Theorem

Every matrix is row equivalent to one and only one matrix in reduced row echelon form.

We'll give an algorithm, called **row reduction**, which demonstrates that every matrix is row equivalent to *at least one* matrix in reduced row echelon form.

Note: Like echelon forms, the row reduction algorithm does not care if a column is augmented: ignore the vertical line when row reducing.

The uniqueness statement is interesting—it means that, no matter *how* you row reduce, you *always* get the same matrix in reduced row echelon form. (Assuming you only do the three legal row operations.) (And you don't make any arithmetic errors.)

Reduced Row Echelon Form

Theorem

Every matrix is row equivalent to one and only one matrix in reduced row echelon form.

We'll give an algorithm, called **row reduction**, which demonstrates that every matrix is row equivalent to *at least one* matrix in reduced row echelon form.

Note: Like echelon forms, the row reduction algorithm does not care if a column is augmented: ignore the vertical line when row reducing.

The uniqueness statement is interesting—it means that, no matter *how* you row reduce, you *always* get the same matrix in reduced row echelon form. (Assuming you only do the three legal row operations.) (And you don't make any arithmetic errors.)

Maybe you can figure out why it's true!

Row Reduction Algorithm

Step 1a Swap the 1st row with a lower one so a leftmost nonzero entry is in 1st row (if necessary).

Example

$$\left(\begin{array}{ccc|c} 0 & -7 & -4 & 2 \\ 2 & 4 & 6 & 12 \\ 3 & 1 & -1 & -2 \end{array} \right)$$

Row Reduction Algorithm

Step 1a Swap the 1st row with a lower one so a leftmost nonzero entry is in 1st row (if necessary).

Step 1b Scale 1st row so that its leading entry is equal to 1.

Example

$$\left(\begin{array}{ccc|c} 0 & -7 & -4 & 2 \\ 2 & 4 & 6 & 12 \\ 3 & 1 & -1 & -2 \end{array} \right)$$

Row Reduction Algorithm

Step 1a Swap the 1st row with a lower one so a leftmost nonzero entry is in 1st row (if necessary).

Step 1b Scale 1st row so that its leading entry is equal to 1.

Step 1c Use row replacement so all entries above and below this 1 are 0.

Example

$$\left(\begin{array}{ccc|c} 0 & -7 & -4 & 2 \\ 2 & 4 & 6 & 12 \\ 3 & 1 & -1 & -2 \end{array} \right)$$

Row Reduction Algorithm

Step 1a Swap the 1st row with a lower one so a leftmost nonzero entry is in 1st row (if necessary).

Step 1b Scale 1st row so that its leading entry is equal to 1.

Step 1c Use row replacement so all entries above and below this 1 are 0.

Step 2a Cover the first row, swap the 2nd row with a lower one so that the leftmost nonzero (uncovered) entry is in 2nd row; uncover 1st row.

Example

$$\left(\begin{array}{ccc|c} 0 & -7 & -4 & 2 \\ 2 & 4 & 6 & 12 \\ 3 & 1 & -1 & -2 \end{array} \right)$$

Row Reduction Algorithm

Step 1a Swap the 1st row with a lower one so a leftmost nonzero entry is in 1st row (if necessary).

Step 1b Scale 1st row so that its leading entry is equal to 1.

Step 1c Use row replacement so all entries above and below this 1 are 0.

Step 2a Cover the first row, swap the 2nd row with a lower one so that the leftmost nonzero (uncovered) entry is in 2nd row; uncover 1st row.

Step 2b Scale 2nd row so that its leading entry is equal to 1.

Example

$$\left(\begin{array}{ccc|c} 0 & -7 & -4 & 2 \\ 2 & 4 & 6 & 12 \\ 3 & 1 & -1 & -2 \end{array} \right)$$

Row Reduction Algorithm

Step 1a Swap the 1st row with a lower one so a leftmost nonzero entry is in 1st row (if necessary).

Step 1b Scale 1st row so that its leading entry is equal to 1.

Step 1c Use row replacement so all entries above and below this 1 are 0.

Step 2a Cover the first row, swap the 2nd row with a lower one so that the leftmost nonzero (uncovered) entry is in 2nd row; uncover 1st row.

Step 2b Scale 2nd row so that its leading entry is equal to 1.

Step 2c Use row replacement so all entries above and below this 1 are 0.

Example

$$\left(\begin{array}{ccc|c} 0 & -7 & -4 & 2 \\ 2 & 4 & 6 & 12 \\ 3 & 1 & -1 & -2 \end{array} \right)$$

Row Reduction Algorithm

Step 1a Swap the 1st row with a lower one so a leftmost nonzero entry is in 1st row (if necessary).

Step 1b Scale 1st row so that its leading entry is equal to 1.

Step 1c Use row replacement so all entries above and below this 1 are 0.

Step 2a Cover the first row, swap the 2nd row with a lower one so that the leftmost nonzero (uncovered) entry is in 2nd row; uncover 1st row.

Step 2b Scale 2nd row so that its leading entry is equal to 1.

Step 2c Use row replacement so all entries above and below this 1 are 0.

Step 3a Cover the first two rows, swap the 3rd row with a lower one so that the leftmost nonzero (uncovered) entry is in 3rd row; uncover first two rows.

Example

$$\left(\begin{array}{ccc|c} 0 & -7 & -4 & 2 \\ 2 & 4 & 6 & 12 \\ 3 & 1 & -1 & -2 \end{array} \right)$$

Row Reduction Algorithm

Step 1a Swap the 1st row with a lower one so a leftmost nonzero entry is in 1st row (if necessary).

Step 1b Scale 1st row so that its leading entry is equal to 1.

Step 1c Use row replacement so all entries above and below this 1 are 0.

Step 2a Cover the first row, swap the 2nd row with a lower one so that the leftmost nonzero (uncovered) entry is in 2nd row; uncover 1st row.

Step 2b Scale 2nd row so that its leading entry is equal to 1.

Step 2c Use row replacement so all entries above and below this 1 are 0.

Step 3a Cover the first two rows, swap the 3rd row with a lower one so that the leftmost nonzero (uncovered) entry is in 3rd row; uncover first two rows.

etc.

Example

$$\left(\begin{array}{ccc|c} 0 & -7 & -4 & 2 \\ 2 & 4 & 6 & 12 \\ 3 & 1 & -1 & -2 \end{array} \right)$$

Row Reduction

Example

$$\left(\begin{array}{ccc|c} 0 & -7 & -4 & 2 \\ 2 & 4 & 6 & 12 \\ 3 & 1 & -1 & -2 \end{array} \right)$$

Row Reduction

Example, continued

$$\left(\begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 0 & -5 & -10 & -20 \\ 0 & -7 & -4 & 2 \end{array} \right)$$

Row Reduction

Example, continued

$$\left(\begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 0 & -5 & -10 & -20 \\ 0 & -7 & -4 & 2 \end{array} \right)$$

Note: Step 2 never messes up the first (nonzero) column of the matrix, because it looks like this:

“Active” row \rightarrow $\left(\begin{array}{ccc|c} 1 & * & * & * \\ 0 & * & * & * \\ 0 & * & * & * \end{array} \right)$

Row Reduction

Example, continued

$$\left(\begin{array}{ccc|c} 1 & 0 & -1 & -2 \\ 0 & 1 & 2 & 4 \\ 0 & 0 & 10 & 30 \end{array} \right)$$

Row Reduction

Example, continued

$$\left(\begin{array}{ccc|c} 1 & 0 & -1 & -2 \\ 0 & 1 & 2 & 4 \\ 0 & 0 & 10 & 30 \end{array} \right)$$

Note: Step 3 never messes up the columns to the left.

Row Reduction

Example, continued

$$\left(\begin{array}{ccc|c} 1 & 0 & -1 & -2 \\ 0 & 1 & 2 & 4 \\ 0 & 0 & 10 & 30 \end{array} \right)$$

Note: Step 3 never messes up the columns to the left.

Success! The reduced row echelon form is

$$\left(\begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & -2 \\ 0 & 0 & 1 & 3 \end{array} \right) \implies \begin{cases} x & = & 1 \\ y & = & -2 \\ z & = & 3 \end{cases}$$

Row Reduction

Example, continued

$$\left(\begin{array}{ccc|c} 1 & 0 & -1 & -2 \\ 0 & 1 & 2 & 4 \\ 0 & 0 & 10 & 30 \end{array} \right)$$

Note: Step 3 never messes up the columns to the left.

Success! The reduced row echelon form is

$$\left(\begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & -2 \\ 0 & 0 & 1 & 3 \end{array} \right) \implies \begin{cases} x & = & 1 \\ y & = & -2 \\ z & = & 3 \end{cases}$$

Step 4: profit?

Row Reduction

Another example

The linear system

$$2x + 10y = -1$$

$$3x + 15y = 2$$

gives rise to the matrix

Row Reduction

Another example

The linear system

$$2x + 10y = -1$$

$$3x + 15y = 2$$

gives rise to the matrix

Let's row reduce it:

$$\left(\begin{array}{cc|c} 2 & 10 & -1 \\ 3 & 15 & 2 \end{array} \right)$$

Row Reduction

Another example

The linear system

$$2x + 10y = -1$$

$$3x + 15y = 2$$

gives rise to the matrix

Let's row reduce it:

$$\left(\begin{array}{cc|c} 2 & 10 & -1 \\ 3 & 15 & 2 \end{array} \right)$$

The row reduced matrix

$$\left(\begin{array}{cc|c} 1 & 5 & 0 \\ 0 & 0 & 1 \end{array} \right)$$

corresponds to the
inconsistent system

$$\begin{aligned} x + 5y &= 0 \\ 0 &= 1. \end{aligned}$$

Inconsistent Matrices

Question

What does an augmented matrix in reduced row echelon form look like, if its system of linear equations is inconsistent?

Inconsistent Matrices

Question

What does an augmented matrix in reduced row echelon form look like, if its system of linear equations is inconsistent?

Answer:

$$\left(\begin{array}{cccc|c} 1 & 0 & \star & \star & 0 \\ 0 & 1 & \star & \star & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right)$$

Inconsistent Matrices

Question

What does an augmented matrix in reduced row echelon form look like, if its system of linear equations is inconsistent?

Answer:

$$\left(\begin{array}{cccc|c} 1 & 0 & \star & \star & 0 \\ 0 & 1 & \star & \star & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right)$$

An augmented matrix corresponds to an inconsistent system of equations if and only if *the last* (i.e., the augmented) *column is a pivot column*.

Another Example

The linear system

$$2x + y + 12z = 1$$

$$x + 2y + 9z = -1$$

gives rise to the matrix $\left(\begin{array}{ccc|c} 2 & 1 & 12 & 1 \\ 1 & 2 & 9 & -1 \end{array} \right)$.

Another Example

The linear system

$$2x + y + 12z = 1$$

$$x + 2y + 9z = -1$$

gives rise to the matrix $\left(\begin{array}{ccc|c} 2 & 1 & 12 & 1 \\ 1 & 2 & 9 & -1 \end{array} \right)$.

Let's row reduce it:

$$\left(\begin{array}{ccc|c} 2 & 1 & 12 & 1 \\ 1 & 2 & 9 & -1 \end{array} \right)$$

Another Example

The linear system

$$2x + y + 12z = 1$$

$$x + 2y + 9z = -1$$

gives rise to the matrix $\left(\begin{array}{ccc|c} 2 & 1 & 12 & 1 \\ 1 & 2 & 9 & -1 \end{array} \right)$.

Let's row reduce it:

$$\left(\begin{array}{ccc|c} 2 & 1 & 12 & 1 \\ 1 & 2 & 9 & -1 \end{array} \right)$$

The row reduced matrix

$$\left(\begin{array}{ccc|c} 1 & 0 & 5 & 1 \\ 0 & 1 & 2 & -1 \end{array} \right)$$

corresponds to the
linear system

$$\begin{cases} x + 5z = 1 \\ y + 2z = -1 \end{cases}$$

Another Example

Continued

The system

$$x + 5z = 1$$

$$y + 2z = -1$$

comes from a matrix in reduced row echelon form.

Another Example

Continued

The system

$$x + 5z = 1$$

$$y + 2z = -1$$

comes from a matrix in reduced row echelon form. Are we done? Is the system solved?

Another Example

Continued

The system

$$x + 5z = 1$$

$$y + 2z = -1$$

comes from a matrix in reduced row echelon form. Are we done? Is the system solved?

Yes! Rewrite:

$$x = 1 - 5z$$

$$y = -1 - 2z$$

For any value of z , there is exactly one value of x and y that makes the equations true. But z can be *anything we want*!

Another Example

Continued

The system

$$x + 5z = 1$$

$$y + 2z = -1$$

comes from a matrix in reduced row echelon form. Are we done? Is the system solved?

Yes! Rewrite:

$$x = 1 - 5z$$

$$y = -1 - 2z$$

For any value of z , there is exactly one value of x and y that makes the equations true. But z can be *anything we want*!

So we have found the solution set: it is all values x, y, z where

$$x = 1 - 5z$$

$$y = -1 - 2z \quad \text{for } z \text{ any real number.}$$

$$(z = z)$$

Another Example

Continued

The system

$$x + 5z = 1$$

$$y + 2z = -1$$

comes from a matrix in reduced row echelon form. Are we done? Is the system solved?

Yes! Rewrite:

$$x = 1 - 5z$$

$$y = -1 - 2z$$

For any value of z , there is exactly one value of x and y that makes the equations true. But z can be *anything we want*!

So we have found the solution set: it is all values x, y, z where

$$x = 1 - 5z$$

$$y = -1 - 2z \quad \text{for } z \text{ any real number.}$$

$$(z = z)$$

This is called the **parametric form** for the solution.

Free Variables

Definition

Consider a *consistent* linear system of equations in the variables x_1, \dots, x_n . Let A be a row echelon form of the matrix for this system.

Free Variables

Definition

Consider a *consistent* linear system of equations in the variables x_1, \dots, x_n . Let A be a row echelon form of the matrix for this system.

We say that x_i is a **free variable** if its corresponding column in A is *not* a pivot column.

Free Variables

Definition

Consider a *consistent* linear system of equations in the variables x_1, \dots, x_n . Let A be a row echelon form of the matrix for this system.

We say that x_i is a **free variable** if its corresponding column in A is *not* a pivot column.

Important

1. You can choose *any value* for the free variables in a (consistent) linear system.

Free Variables

Definition

Consider a *consistent* linear system of equations in the variables x_1, \dots, x_n . Let A be a row echelon form of the matrix for this system.

We say that x_i is a **free variable** if its corresponding column in A is *not* a pivot column.

Important

1. You can choose *any value* for the free variables in a (consistent) linear system.
2. Free variables come from *columns without pivots* in a matrix in row echelon form.

Free Variables

Definition

Consider a *consistent* linear system of equations in the variables x_1, \dots, x_n . Let A be a row echelon form of the matrix for this system.

We say that x_i is a **free variable** if its corresponding column in A is *not* a pivot column.

Important

1. You can choose *any value* for the free variables in a (consistent) linear system.
2. Free variables come from *columns without pivots* in a matrix in row echelon form.

In the previous example, z was free because the reduced row echelon form matrix was

$$\left(\begin{array}{ccc|c} 1 & 0 & 5 & 4 \\ 0 & 1 & 2 & -1 \end{array} \right).$$

Free Variables

Definition

Consider a *consistent* linear system of equations in the variables x_1, \dots, x_n . Let A be a row echelon form of the matrix for this system.

We say that x_i is a **free variable** if its corresponding column in A is *not* a pivot column.

Important

1. You can choose *any value* for the free variables in a (consistent) linear system.
2. Free variables come from *columns without pivots* in a matrix in row echelon form.

In the previous example, z was free because the reduced row echelon form matrix was

$$\left(\begin{array}{ccc|c} 1 & 0 & 5 & 4 \\ 0 & 1 & 2 & -1 \end{array} \right).$$

In this matrix:

$$\left(\begin{array}{cccc|c} 1 & \star & 0 & \star & \star \\ 0 & 0 & 1 & \star & \star \end{array} \right)$$

the free variables are _____.

Free Variables

Definition

Consider a *consistent* linear system of equations in the variables x_1, \dots, x_n . Let A be a row echelon form of the matrix for this system.

We say that x_i is a **free variable** if its corresponding column in A is *not* a pivot column.

Important

1. You can choose *any value* for the free variables in a (consistent) linear system.
2. Free variables come from *columns without pivots* in a matrix in row echelon form.

In the previous example, z was free because the reduced row echelon form matrix was

$$\left(\begin{array}{ccc|c} 1 & 0 & 5 & 4 \\ 0 & 1 & 2 & -1 \end{array} \right).$$

In this matrix:

$$\left(\begin{array}{cccc|c} 1 & \star & 0 & \star & \star \\ 0 & 0 & 1 & \star & \star \end{array} \right)$$

the free variables are x_2 and x_4 .

Free Variables

Definition

Consider a *consistent* linear system of equations in the variables x_1, \dots, x_n . Let A be a row echelon form of the matrix for this system.

We say that x_i is a **free variable** if its corresponding column in A is *not* a pivot column.

Important

1. You can choose *any value* for the free variables in a (consistent) linear system.
2. Free variables come from *columns without pivots* in a matrix in row echelon form.

In the previous example, z was free because the reduced row echelon form matrix was

$$\left(\begin{array}{ccc|c} 1 & 0 & 5 & 4 \\ 0 & 1 & 2 & -1 \end{array} \right).$$

In this matrix:

$$\left(\begin{array}{cccc|c} 1 & \star & 0 & \star & \star \\ 0 & 0 & 1 & \star & \star \end{array} \right)$$

the free variables are x_2 and x_4 . (What about the last column?)

One More Example

The reduced row echelon form of the matrix for a linear system in x_1, x_2, x_3, x_4 is

$$\left(\begin{array}{cccc|c} 1 & 0 & 0 & 3 & 2 \\ 0 & 0 & 1 & 4 & -1 \end{array} \right)$$

One More Example

The reduced row echelon form of the matrix for a linear system in x_1, x_2, x_3, x_4 is

$$\left(\begin{array}{cccc|c} 1 & 0 & 0 & 3 & 2 \\ 0 & 0 & 1 & 4 & -1 \end{array} \right)$$

The free variables are _____:

One More Example

The reduced row echelon form of the matrix for a linear system in x_1, x_2, x_3, x_4 is

$$\left(\begin{array}{cccc|c} 1 & 0 & 0 & 3 & 2 \\ 0 & 0 & 1 & 4 & -1 \end{array} \right)$$

The free variables are x_2 and x_4 : they are the ones whose columns are *not* pivot columns.

One More Example

The reduced row echelon form of the matrix for a linear system in x_1, x_2, x_3, x_4 is

$$\left(\begin{array}{cccc|c} 1 & 0 & 0 & 3 & 2 \\ 0 & 0 & 1 & 4 & -1 \end{array} \right)$$

The free variables are x_2 and x_4 : they are the ones whose columns are *not* pivot columns.

This translates into the system of equations

$$\begin{cases} x_1 & + 3x_4 = 2 \\ & x_3 + x_4 = -1 \end{cases}$$

One More Example

The reduced row echelon form of the matrix for a linear system in x_1, x_2, x_3, x_4 is

$$\left(\begin{array}{cccc|c} 1 & 0 & 0 & 3 & 2 \\ 0 & 0 & 1 & 4 & -1 \end{array} \right)$$

The free variables are x_2 and x_4 : they are the ones whose columns are *not* pivot columns.

This translates into the system of equations

$$\begin{cases} x_1 & + 3x_4 = 2 \\ x_3 + x_4 = -1 \end{cases} \implies \boxed{\begin{array}{l} x_1 = 2 - 3x_4 \\ x_3 = -1 - 4x_4 \end{array}}$$

One More Example

The reduced row echelon form of the matrix for a linear system in x_1, x_2, x_3, x_4 is

$$\left(\begin{array}{cccc|c} 1 & 0 & 0 & 3 & 2 \\ 0 & 0 & 1 & 4 & -1 \end{array} \right)$$

The free variables are x_2 and x_4 : they are the ones whose columns are *not* pivot columns.

This translates into the system of equations

$$\begin{cases} x_1 & + 3x_4 = 2 \\ & x_3 + x_4 = -1 \end{cases} \implies$$

$\begin{aligned} x_1 &= 2 - 3x_4 \\ x_3 &= -1 - 4x_4 \end{aligned}$

What happened to x_2 ? What is it allowed to be?

One More Example

The reduced row echelon form of the matrix for a linear system in x_1, x_2, x_3, x_4 is

$$\left(\begin{array}{cccc|c} 1 & 0 & 0 & 3 & 2 \\ 0 & 0 & 1 & 4 & -1 \end{array} \right)$$

The free variables are x_2 and x_4 : they are the ones whose columns are *not* pivot columns.

This translates into the system of equations

$$\begin{cases} x_1 & + 3x_4 = 2 \\ & x_3 + x_4 = -1 \end{cases} \implies \boxed{\begin{array}{l} x_1 = 2 - 3x_4 \\ x_3 = -1 - 4x_4 \end{array}}$$

What happened to x_2 ? What is it allowed to be? Anything! The general solution is

for any values of x_2 and x_4 .

One More Example

The reduced row echelon form of the matrix for a linear system in x_1, x_2, x_3, x_4 is

$$\left(\begin{array}{cccc|c} 1 & 0 & 0 & 3 & 2 \\ 0 & 0 & 1 & 4 & -1 \end{array} \right)$$

The free variables are x_2 and x_4 : they are the ones whose columns are *not* pivot columns.

This translates into the system of equations

$$\begin{cases} x_1 & + 3x_4 = 2 \\ & x_3 + x_4 = -1 \end{cases} \implies \boxed{\begin{array}{l} x_1 = 2 - 3x_4 \\ x_3 = -1 - 4x_4 \end{array}}$$

What happened to x_2 ? What is it allowed to be? Anything! The general solution is

for any values of x_2 and x_4 .

The boxed equation is called the **parametric form** of the general solution to the system of equations. It is obtained by moving all free variables to the right-hand side of the $=$.

Poll

Is it possible for a system of linear equations to have exactly two solutions?

Summary

There are *three possibilities* for the reduced row echelon form of the augmented matrix of a linear system.

Summary

There are *three possibilities* for the reduced row echelon form of the augmented matrix of a linear system.

1. The last column is a pivot column.

In this case, the system is *inconsistent*.

Summary

There are *three possibilities* for the reduced row echelon form of the augmented matrix of a linear system.

1. The last column is a pivot column.

In this case, the system is *inconsistent*. There are *zero* solutions, i.e. the solution set is *empty*. Picture:

Summary

There are *three possibilities* for the reduced row echelon form of the augmented matrix of a linear system.

1. The last column is a pivot column.

In this case, the system is *inconsistent*. There are *zero* solutions, i.e. the solution set is *empty*. Picture:

$$\left(\begin{array}{cc|c} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right)$$

Summary

There are *three possibilities* for the reduced row echelon form of the augmented matrix of a linear system.

1. The last column is a pivot column.

In this case, the system is *inconsistent*. There are *zero* solutions, i.e. the solution set is *empty*. Picture:

$$\left(\begin{array}{ccc|c} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{array} \right)$$

2. Every column except the last column is a pivot column.

In this case, the system has a *unique solution*. Picture:

$$\left(\begin{array}{ccc|c} 1 & 0 & 0 & \star \\ 0 & 1 & 0 & \star \\ 0 & 0 & 1 & \star \end{array} \right)$$

Summary

There are *three possibilities* for the reduced row echelon form of the augmented matrix of a linear system.

1. The last column is a pivot column.

In this case, the system is *inconsistent*. There are *zero* solutions, i.e. the solution set is *empty*. Picture:

$$\left(\begin{array}{cc|c} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right)$$

2. Every column except the last column is a pivot column.

In this case, the system has a *unique solution*. Picture:

$$\left(\begin{array}{ccc|c} 1 & 0 & 0 & \star \\ 0 & 1 & 0 & \star \\ 0 & 0 & 1 & \star \end{array} \right)$$

3. The last column is not a pivot column, and some other column isn't either.

In this case, the system has *infinitely many* solutions, corresponding to the infinitely many possible values of the free variable(s). Picture:

$$\left(\begin{array}{cccc|c} 1 & \star & 0 & \star & \star \\ 0 & 0 & 1 & \star & \star \end{array} \right)$$