

Section 1.3

Vector Equations


Motivation

We want to think about the *algebra* in linear algebra (systems of equations and their solution sets) in terms of *geometry* (points, lines, planes, etc).

Motivation

We want to think about the *algebra* in linear algebra (systems of equations and their solution sets) in terms of *geometry* (points, lines, planes, etc).


$$\begin{array}{rcl} x - 3y & = & -3 \\ 2x + y & = & 8 \end{array}$$



Motivation

We want to think about the *algebra* in linear algebra (systems of equations and their solution sets) in terms of *geometry* (points, lines, planes, etc).

$$\begin{array}{rcl} x - 3y & = & -3 \\ 2x + y & = & 8 \end{array}$$




This will give us better insight into the properties of systems of equations and their solution sets.

Motivation

We want to think about the *algebra* in linear algebra (systems of equations and their solution sets) in terms of *geometry* (points, lines, planes, etc).

$$\begin{array}{rcl} x - 3y & = & -3 \\ 2x + y & = & 8 \end{array}$$



This will give us better insight into the properties of systems of equations and their solution sets.

To do this, we need to introduce n -dimensional space \mathbf{R}^n , and **vectors** inside it.

Line, Plane, Space, ...

Recall that \mathbf{R} denotes the collection of all real numbers, i.e. the number line.

Recall that \mathbf{R} denotes the collection of all real numbers, i.e. the number line.

Definition

Let n be a positive whole number. We define

$$\mathbf{R}^n = \text{all ordered } n\text{-tuples of real numbers } (x_1, x_2, x_3, \dots, x_n).$$

Line, Plane, Space, ...

Recall that \mathbf{R} denotes the collection of all real numbers, i.e. the number line.


Definition

Let n be a positive whole number. We define

$$\mathbf{R}^n = \text{all ordered } n\text{-tuples of real numbers } (x_1, x_2, x_3, \dots, x_n).$$

Example

When $n = 1$, we just get \mathbf{R} back: $\mathbf{R}^1 = \mathbf{R}$. Geometrically, this is the *number line*.




Line, Plane, Space, ...

Continued

Example

When $n = 2$, we can think of \mathbf{R}^2 as the *plane*. This is because every point on the plane can be represented by an ordered pair of real numbers, namely, its x - and y -coordinates.




Line, Plane, Space, ...

Continued

Example

When $n = 2$, we can think of \mathbf{R}^2 as the *plane*. This is because every point on the plane can be represented by an ordered pair of real numbers, namely, its x - and y -coordinates.




We can use the elements of \mathbf{R}^2 to *label* points on the plane, but \mathbf{R}^2 is not defined to be the plane!

Line, Plane, Space, ...

Continued

Example

When $n = 3$, we can think of \mathbf{R}^3 as the *space* we (appear to) live in. This is because every point in space can be represented by an ordered triple of real numbers, namely, its x -, y -, and z -coordinates.




Line, Plane, Space, ...

Continued

Example

When $n = 3$, we can think of \mathbf{R}^3 as the *space* we (appear to) live in. This is because every point in space can be represented by an ordered triple of real numbers, namely, its x -, y -, and z -coordinates.



Again, we can use the elements of \mathbf{R}^3 to *label* points in space, but \mathbf{R}^3 is not defined to be space!

Line, Plane, Space, ...

Continued

So what is \mathbf{R}^4 ? or \mathbf{R}^5 ? or \mathbf{R}^n ?

Line, Plane, Space, ...

Continued

So what is \mathbf{R}^4 ? or \mathbf{R}^5 ? or \mathbf{R}^n ?

...go back to the *definition*: ordered n -tuples of real numbers

$$(x_1, x_2, x_3, \dots, x_n).$$

Line, Plane, Space, ...

Continued

So what is \mathbf{R}^4 ? or \mathbf{R}^5 ? or \mathbf{R}^n ?

...go back to the *definition*: ordered n -tuples of real numbers

$$(x_1, x_2, x_3, \dots, x_n).$$

They're still “geometric” spaces, in the sense that our intuition for \mathbf{R}^2 and \mathbf{R}^3 sometimes extends to \mathbf{R}^n , but they're harder to visualize.

Line, Plane, Space, ...

Continued

So what is \mathbf{R}^4 ? or \mathbf{R}^5 ? or \mathbf{R}^n ?

...go back to the *definition*: ordered n -tuples of real numbers

$$(x_1, x_2, x_3, \dots, x_n).$$

They're still “geometric” spaces, in the sense that our intuition for \mathbf{R}^2 and \mathbf{R}^3 sometimes extends to \mathbf{R}^n , but they're harder to visualize.

We'll make definitions and state theorems that apply to any \mathbf{R}^n , but we'll only draw pictures for \mathbf{R}^2 and \mathbf{R}^3 .


Vectors

In the previous slides, we were thinking of elements of \mathbf{R}^n as **points**: in the line, plane, space, etc.

Vectors

In the previous slides, we were thinking of elements of \mathbf{R}^n as **points**: in the line, plane, space, etc.


We can also think of them as **vectors**: arrows with a given length and direction.



Vectors

In the previous slides, we were thinking of elements of \mathbf{R}^n as **points**: in the line, plane, space, etc.

We can also think of them as **vectors**: arrows with a given length and direction.




So the vector points *horizontally* in the amount of its x-coordinate, and *vertically* in the amount of its y-coordinate.

Vectors

In the previous slides, we were thinking of elements of \mathbf{R}^n as **points**: in the line, plane, space, etc.

We can also think of them as **vectors**: arrows with a given length and direction.



So the vector points *horizontally* in the amount of its x -coordinate, and *vertically* in the amount of its y -coordinate.

When we think of an element of \mathbf{R}^n as a vector, we write it as a matrix with n rows and one column:

$$v = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}.$$

We'll see why this is useful later.


Points and Vectors

So what is the difference between a point and a vector?

Points and Vectors

So what is the difference between a point and a vector?

A vector need not start at the origin: *it can be located anywhere!* In other words, an arrow is determined by its length and its direction, not by its location.




These arrows all represent the vector $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$.

Points and Vectors

So what is the difference between a point and a vector?

A vector need not start at the origin: *it can be located anywhere!* In other words, an arrow is determined by its length and its direction, not by its location.



These arrows all represent the vector $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$.

However, unless otherwise specified, we'll assume a vector starts at the origin: we'll usually be sloppy and identify the vector $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$ with the point (1,2).

Points and Vectors

So what is the difference between a point and a vector?

A vector need not start at the origin: *it can be located anywhere!* In other words, an arrow is determined by its length and its direction, not by its location.



These arrows all represent the vector $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$.


However, unless otherwise specified, we'll assume a vector starts at the origin: we'll usually be sloppy and identify the vector $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$ with the point (1, 2).

This makes sense in the real world: many physical quantities, such as velocity, are represented as vectors. But it makes more sense to think of the velocity of a car as being located at the car.

Points and Vectors

So what is the difference between a point and a vector?

A vector need not start at the origin: *it can be located anywhere!* In other words, an arrow is determined by its length and its direction, not by its location.



These arrows all represent the vector $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$.

However, unless otherwise specified, we'll assume a vector starts at the origin: we'll usually be sloppy and identify the vector $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$ with the point (1, 2).

This makes sense in the real world: many physical quantities, such as velocity, are represented as vectors. But it makes more sense to think of the velocity of a car as being located at the car.

Another way to think about it: a vector is a *difference* between two points, or the arrow from one point to another.

Points and Vectors

So what is the difference between a point and a vector?

A vector need not start at the origin: *it can be located anywhere!* In other words, an arrow is determined by its length and its direction, not by its location.




These arrows all represent the vector $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$.

However, unless otherwise specified, we'll assume a vector starts at the origin: we'll usually be sloppy and identify the vector $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$ with the point $(1, 2)$.

This makes sense in the real world: many physical quantities, such as velocity, are represented as vectors. But it makes more sense to think of the velocity of a car as being located at the car.

Another way to think about it: a vector is a *difference* between two points, or the arrow from one point to another.

For instance, $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$ is the arrow from $(1, 1)$ to $(2, 3)$.



Definition

- ▶ We can add two vectors together:

$$\begin{pmatrix} a \\ b \\ c \end{pmatrix} + \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} a+x \\ b+y \\ c+z \end{pmatrix} .$$

Definition

- ▶ We can add two vectors together:

$$\begin{pmatrix} a \\ b \\ c \end{pmatrix} + \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} a + x \\ b + y \\ c + z \end{pmatrix} .$$

- ▶ We can multiply, or **scale**, a vector by a real number c :

$$c \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} c \cdot x \\ c \cdot y \\ c \cdot z \end{pmatrix} .$$

Definition

- ▶ We can add two vectors together:

$$\begin{pmatrix} a \\ b \\ c \end{pmatrix} + \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} a+x \\ b+y \\ c+z \end{pmatrix}.$$

- ▶ We can multiply, or **scale**, a vector by a real number c :

$$c \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} c \cdot x \\ c \cdot y \\ c \cdot z \end{pmatrix}.$$

We call c a **scalar** to distinguish it from a vector. If v is a vector and c is a scalar, cv is called a **scalar multiple** of v .

Definition

- ▶ We can add two vectors together:

$$\begin{pmatrix} a \\ b \\ c \end{pmatrix} + \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} a+x \\ b+y \\ c+z \end{pmatrix}.$$

- ▶ We can multiply, or **scale**, a vector by a real number c :

$$c \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} c \cdot x \\ c \cdot y \\ c \cdot z \end{pmatrix}.$$

We call c a **scalar** to distinguish it from a vector. If v is a vector and c is a scalar, cv is called a **scalar multiple** of v .

(And likewise for vectors of length n .)

Definition

- ▶ We can add two vectors together:

$$\begin{pmatrix} a \\ b \\ c \end{pmatrix} + \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} a+x \\ b+y \\ c+z \end{pmatrix}.$$

- ▶ We can multiply, or **scale**, a vector by a real number c :


$$c \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} c \cdot x \\ c \cdot y \\ c \cdot z \end{pmatrix}.$$

We call c a **scalar** to distinguish it from a vector. If v is a vector and c is a scalar, cv is called a **scalar multiple** of v .


(And likewise for vectors of length n .) For instance,

Vector Addition and Subtraction: Geometry

The parallelogram law for vector addition




Vector Addition and Subtraction: Geometry



The parallelogram law for vector addition

Geometrically, the sum of two vectors v, w is obtained as follows:


Vector Addition and Subtraction: Geometry



The parallelogram law for vector addition

Geometrically, the sum of two vectors v, w is obtained as follows: place the tail of w at the head of v .


Vector Addition and Subtraction: Geometry



The parallelogram law for vector addition

Geometrically, the sum of two vectors v, w is obtained as follows: place the tail of w at the head of v . Then $v + w$ is the vector whose tail is the tail of v and whose head is the head of w .


Vector Addition and Subtraction: Geometry



The parallelogram law for vector addition

Geometrically, the sum of two vectors v, w is obtained as follows: place the tail of w at the head of v . Then $v + w$ is the vector whose tail is the tail of v and whose head is the head of w . Doing this both ways creates a **parallelogram**.

Vector Addition and Subtraction: Geometry




The parallelogram law for vector addition

Geometrically, the sum of two vectors v, w is obtained as follows: place the tail of w at the head of v . Then $v + w$ is the vector whose tail is the tail of v and whose head is the head of w . Doing this both ways creates a **parallelogram**. For example,

$$\begin{pmatrix} 1 \\ 3 \end{pmatrix} + \begin{pmatrix} 4 \\ 2 \end{pmatrix} = \begin{pmatrix} 5 \\ 5 \end{pmatrix}.$$

Vector Addition and Subtraction: Geometry




The parallelogram law for vector addition

Geometrically, the sum of two vectors v, w is obtained as follows: place the tail of w at the head of v . Then $v + w$ is the vector whose tail is the tail of v and whose head is the head of w . Doing this both ways creates a **parallelogram**. For example,

$$\begin{pmatrix} 1 \\ 3 \end{pmatrix} + \begin{pmatrix} 4 \\ 2 \end{pmatrix} = \begin{pmatrix} 5 \\ 5 \end{pmatrix}.$$

Why? The width of $v + w$ is the sum of the widths, and likewise with the heights.

Vector Addition and Subtraction: Geometry




The parallelogram law for vector addition

Geometrically, the sum of two vectors v, w is obtained as follows: place the tail of w at the head of v . Then $v + w$ is the vector whose tail is the tail of v and whose head is the head of w . Doing this both ways creates a **parallelogram**. For example,


$$\begin{pmatrix} 1 \\ 3 \end{pmatrix} + \begin{pmatrix} 4 \\ 2 \end{pmatrix} = \begin{pmatrix} 5 \\ 5 \end{pmatrix}.$$

Why? The width of $v + w$ is the sum of the widths, and likewise with the heights.

Vector subtraction



Vector Addition and Subtraction: Geometry



The parallelogram law for vector addition


Geometrically, the sum of two vectors v, w is obtained as follows: place the tail of w at the head of v . Then $v + w$ is the vector whose tail is the tail of v and whose head is the head of w . Doing this both ways creates a **parallelogram**. For example,

$$\begin{pmatrix} 1 \\ 3 \end{pmatrix} + \begin{pmatrix} 4 \\ 2 \end{pmatrix} = \begin{pmatrix} 5 \\ 5 \end{pmatrix}.$$


Why? The width of $v + w$ is the sum of the widths, and likewise with the heights.

Vector subtraction

Geometrically, the difference of two vectors v, w is obtained as follows:



Vector Addition and Subtraction: Geometry



The parallelogram law for vector addition


Geometrically, the sum of two vectors v, w is obtained as follows: place the tail of w at the head of v . Then $v + w$ is the vector whose tail is the tail of v and whose head is the head of w . Doing this both ways creates a **parallelogram**. For example,

$$\begin{pmatrix} 1 \\ 3 \end{pmatrix} + \begin{pmatrix} 4 \\ 2 \end{pmatrix} = \begin{pmatrix} 5 \\ 5 \end{pmatrix}.$$


Why? The width of $v + w$ is the sum of the widths, and likewise with the heights.

Vector subtraction

Geometrically, the difference of two vectors v, w is obtained as follows: place the tail of v and w at the same point.



Vector Addition and Subtraction: Geometry



The parallelogram law for vector addition


Geometrically, the sum of two vectors v, w is obtained as follows: place the tail of w at the head of v . Then $v + w$ is the vector whose tail is the tail of v and whose head is the head of w . Doing this both ways creates a **parallelogram**. For example,

$$\begin{pmatrix} 1 \\ 3 \end{pmatrix} + \begin{pmatrix} 4 \\ 2 \end{pmatrix} = \begin{pmatrix} 5 \\ 5 \end{pmatrix}.$$


Why? The width of $v + w$ is the sum of the widths, and likewise with the heights.

Vector subtraction

Geometrically, the difference of two vectors v, w is obtained as follows: place the tail of v and w at the same point. Then $v - w$ is the vector from the head of v to the head of w .



Vector Addition and Subtraction: Geometry



The parallelogram law for vector addition

Geometrically, the sum of two vectors v, w is obtained as follows: place the tail of w at the head of v . Then $v + w$ is the vector whose tail is the tail of v and whose head is the head of w . Doing this both ways creates a **parallelogram**. For example,


$$\begin{pmatrix} 1 \\ 3 \end{pmatrix} + \begin{pmatrix} 4 \\ 2 \end{pmatrix} = \begin{pmatrix} 5 \\ 5 \end{pmatrix}.$$

Why? The width of $v + w$ is the sum of the widths, and likewise with the heights.


Vector subtraction

Geometrically, the difference of two vectors v, w is obtained as follows: place the tail of v and w at the same point. Then $v - w$ is the vector from the head of v to the head of w . For example,

$$\begin{pmatrix} 1 \\ 4 \end{pmatrix} - \begin{pmatrix} 4 \\ 2 \end{pmatrix} = \begin{pmatrix} -3 \\ 2 \end{pmatrix}.$$



Vector Addition and Subtraction: Geometry



The parallelogram law for vector addition

Geometrically, the sum of two vectors v, w is obtained as follows: place the tail of w at the head of v . Then $v + w$ is the vector whose tail is the tail of v and whose head is the head of w . Doing this both ways creates a **parallelogram**. For example,

$$\begin{pmatrix} 1 \\ 3 \end{pmatrix} + \begin{pmatrix} 4 \\ 2 \end{pmatrix} = \begin{pmatrix} 5 \\ 5 \end{pmatrix}.$$


Why? The width of $v + w$ is the sum of the widths, and likewise with the heights.

Vector subtraction


Geometrically, the difference of two vectors v, w is obtained as follows: place the tail of v and w at the same point. Then $v - w$ is the vector from the head of v to the head of w . For example,

$$\begin{pmatrix} 1 \\ 4 \end{pmatrix} - \begin{pmatrix} 4 \\ 2 \end{pmatrix} = \begin{pmatrix} -3 \\ 2 \end{pmatrix}.$$

Why? If you add $v - w$ to w , you get v .



Vector Addition and Subtraction: Geometry



The parallelogram law for vector addition

Geometrically, the sum of two vectors v, w is obtained as follows: place the tail of w at the head of v . Then $v + w$ is the vector whose tail is the tail of v and whose head is the head of w . Doing this both ways creates a **parallelogram**. For example,

$$\begin{pmatrix} 1 \\ 3 \end{pmatrix} + \begin{pmatrix} 4 \\ 2 \end{pmatrix} = \begin{pmatrix} 5 \\ 5 \end{pmatrix}.$$

Why? The width of $v + w$ is the sum of the widths, and likewise with the heights.


Vector subtraction

Geometrically, the difference of two vectors v, w is obtained as follows: place the tail of v and w at the same point. Then $v - w$ is the vector from the head of v to the head of w . For example,

$$\begin{pmatrix} 1 \\ 4 \end{pmatrix} - \begin{pmatrix} 4 \\ 2 \end{pmatrix} = \begin{pmatrix} -3 \\ 2 \end{pmatrix}.$$

Why? If you add $v - w$ to w , you get v .

This works in higher dimensions too!



Scalar Multiplication: Geometry

Scalar multiples of a vector

These have the same *direction* but a different *length*.

Some multiples of v .



$$v = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$$2v = \begin{pmatrix} 2 \\ 4 \end{pmatrix}$$

$$-\frac{1}{2}v = \begin{pmatrix} -\frac{1}{2} \\ -1 \end{pmatrix}$$

$$0v = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

Scalar Multiplication: Geometry

Scalar multiples of a vector

These have the same *direction* but a different *length*.

Some multiples of v .




$$v = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$$2v = \begin{pmatrix} 2 \\ 4 \end{pmatrix}$$

$$-\frac{1}{2}v = \begin{pmatrix} -\frac{1}{2} \\ -1 \end{pmatrix}$$

$$0v = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

All multiples of v .




Scalar Multiplication: Geometry

Scalar multiples of a vector


These have the same *direction* but a different *length*.

Some multiples of v .



$$\begin{aligned}v &= \begin{pmatrix} 1 \\ 2 \end{pmatrix} \\2v &= \begin{pmatrix} 2 \\ 4 \end{pmatrix} \\-\frac{1}{2}v &= \begin{pmatrix} -\frac{1}{2} \\ -1 \end{pmatrix} \\0v &= \begin{pmatrix} 0 \\ 0 \end{pmatrix}\end{aligned}$$

All multiples of v .




Scalar Multiplication: Geometry

Scalar multiples of a vector


These have the same *direction* but a different *length*.

Some multiples of v .



$$\begin{aligned}v &= \begin{pmatrix} 1 \\ 2 \end{pmatrix} \\2v &= \begin{pmatrix} 2 \\ 4 \end{pmatrix} \\-\frac{1}{2}v &= \begin{pmatrix} -\frac{1}{2} \\ -1 \end{pmatrix} \\0v &= \begin{pmatrix} 0 \\ 0 \end{pmatrix}\end{aligned}$$

All multiples of v .



So the scalar multiples of v form a *line*.

Linear Combinations

We can add and scalar multiply in the same equation:

Linear Combinations

We can add and scalar multiply in the same equation:

$$w = c_1 v_1 + c_2 v_2 + \cdots + c_p v_p$$

where c_1, c_2, \dots, c_p are _____,

Linear Combinations

We can add and scalar multiply in the same equation:

$$w = c_1 v_1 + c_2 v_2 + \cdots + c_p v_p$$

where c_1, c_2, \dots, c_p are scalars, v_1, v_2, \dots, v_p are _____

Linear Combinations

We can add and scalar multiply in the same equation:

$$w = c_1 v_1 + c_2 v_2 + \cdots + c_p v_p$$

where c_1, c_2, \dots, c_p are scalars, v_1, v_2, \dots, v_p are vectors in \mathbf{R}^n , and w is a

Linear Combinations

We can add and scalar multiply in the same equation:

$$w = c_1 v_1 + c_2 v_2 + \cdots + c_p v_p$$

where c_1, c_2, \dots, c_p are scalars, v_1, v_2, \dots, v_p are vectors in \mathbf{R}^n , and w is a vector in \mathbf{R}^n .

Linear Combinations

We can add and scalar multiply in the same equation:

$$w = c_1 v_1 + c_2 v_2 + \cdots + c_p v_p$$

where c_1, c_2, \dots, c_p are scalars, v_1, v_2, \dots, v_p are vectors in \mathbf{R}^n , and w is a vector in \mathbf{R}^n .

Definition

We call w a **linear combination** of the vectors v_1, v_2, \dots, v_p .

Linear Combinations

We can add and scalar multiply in the same equation:

$$w = c_1 v_1 + c_2 v_2 + \cdots + c_p v_p$$

where c_1, c_2, \dots, c_p are scalars, v_1, v_2, \dots, v_p are vectors in \mathbf{R}^n , and w is a vector in \mathbf{R}^n .

Definition

We call w a **linear combination** of the vectors v_1, v_2, \dots, v_p . The scalars c_1, c_2, \dots, c_p are called the **weights** or **coefficients**.

Linear Combinations

We can add and scalar multiply in the same equation:


$$w = c_1 v_1 + c_2 v_2 + \cdots + c_p v_p$$

where c_1, c_2, \dots, c_p are scalars, v_1, v_2, \dots, v_p are vectors in \mathbf{R}^n , and w is a vector in \mathbf{R}^n .

Definition

We call w a **linear combination** of the vectors v_1, v_2, \dots, v_p . The scalars c_1, c_2, \dots, c_p are called the **weights** or **coefficients**.

Example




Let $v = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$ and $w = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$.

What are some linear combinations of v and w ?

- ▶ $v + w$
- ▶ $v - w$
- ▶ $2v + 0w$
- ▶ $2w$
- ▶ $-v$

Poll


Is there any vector in \mathbf{R}^2 that is *not* a linear combination of v and w ?



Poll

Is there any vector in \mathbf{R}^2 that is *not* a linear combination of v and w ?


No: in fact, every vector in \mathbf{R}^2 is a combination of v and w .



Poll


Is there any vector in \mathbf{R}^2 that is *not* a linear combination of v and w ?

No: in fact, *every* vector in \mathbf{R}^2 is a combination of v and w .




(The purple lines are to help measure *how much* of v and w you need to get to a given point.)

More Examples




What are some linear combinations of $v = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$?

More Examples



What are some linear combinations of $v = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$?




Question


What are all linear combinations of

$$v = \begin{pmatrix} 2 \\ 2 \end{pmatrix} \quad \text{and} \quad w = \begin{pmatrix} -1 \\ -1 \end{pmatrix}?$$

More Examples



What are some linear combinations of $v = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$?




Question

What are all linear combinations of


$$v = \begin{pmatrix} 2 \\ 2 \end{pmatrix} \quad \text{and} \quad w = \begin{pmatrix} -1 \\ -1 \end{pmatrix}?$$

Answer: The line which contains both vectors.

More Examples



What are some linear combinations of $v = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$?



Question

What are all linear combinations of

$$v = \begin{pmatrix} 2 \\ 2 \end{pmatrix} \quad \text{and} \quad w = \begin{pmatrix} -1 \\ -1 \end{pmatrix}?$$

Answer: The line which contains both vectors.

What's different about this example and the one on the poll?

Systems of Linear Equations

Question

Is $\begin{pmatrix} 8 \\ 16 \\ 3 \end{pmatrix}$ a linear combination of $\begin{pmatrix} 1 \\ 2 \\ 6 \end{pmatrix}$ and $\begin{pmatrix} -1 \\ -2 \\ -1 \end{pmatrix}$?

Systems of Linear Equations

Continued

$$x - y = 8$$

$$2x - 2y = 16$$

$$6x - y = 3$$

Systems of Linear Equations

Continued

$$x - y = 8$$

$$2x - 2y = 16$$

$$6x - y = 3$$

matrix form
~~~~~>

$$\left( \begin{array}{cc|c} 1 & -1 & 8 \\ 2 & -2 & 16 \\ 6 & -1 & 3 \end{array} \right)$$

# Systems of Linear Equations

Continued

$$x - y = 8$$

$$2x - 2y = 16$$

$$6x - y = 3$$

matrix form  
~~~~~>

$$\left(\begin{array}{cc|c} 1 & -1 & 8 \\ 2 & -2 & 16 \\ 6 & -1 & 3 \end{array} \right)$$

row reduce
~~~~~>

$$\left( \begin{array}{cc|c} 1 & 0 & -1 \\ 0 & 1 & -9 \\ 0 & 0 & 0 \end{array} \right)$$

# Systems of Linear Equations

Continued

$$x - y = 8$$

$$2x - 2y = 16$$

$$6x - y = 3$$

matrix form

~~~~~>

$$\left(\begin{array}{cc|c} 1 & -1 & 8 \\ 2 & -2 & 16 \\ 6 & -1 & 3 \end{array} \right)$$

row reduce

~~~~~>

$$\left( \begin{array}{cc|c} 1 & 0 & -1 \\ 0 & 1 & -9 \\ 0 & 0 & 0 \end{array} \right)$$

solution

~~~~~>

$$x = -1$$

$$y = -9$$

Systems of Linear Equations

Continued

$$x - y = 8$$

$$2x - 2y = 16$$

$$6x - y = 3$$

matrix form
~~~~~>

$$\left( \begin{array}{cc|c} 1 & -1 & 8 \\ 2 & -2 & 16 \\ 6 & -1 & 3 \end{array} \right)$$

row reduce  
~~~~~>

$$\left(\begin{array}{cc|c} 1 & 0 & -1 \\ 0 & 1 & -9 \\ 0 & 0 & 0 \end{array} \right)$$

solution
~~~~~>

$$x = -1$$

$$y = -9$$

Conclusion:

$$-\begin{pmatrix} 1 \\ 2 \\ 6 \end{pmatrix} - 9 \begin{pmatrix} -1 \\ -2 \\ -1 \end{pmatrix} = \begin{pmatrix} 8 \\ 16 \\ 3 \end{pmatrix}$$

# Systems of Linear Equations

Continued

$$x - y = 8$$

$$2x - 2y = 16$$

$$6x - y = 3$$

matrix form  
~~~~~→

$$\left(\begin{array}{cc|c} 1 & -1 & 8 \\ 2 & -2 & 16 \\ 6 & -1 & 3 \end{array} \right)$$

row reduce
~~~~~→

$$\left( \begin{array}{cc|c} 1 & 0 & -1 \\ 0 & 1 & -9 \\ 0 & 0 & 0 \end{array} \right)$$

solution  
~~~~~→

$$x = -1$$

$$y = -9$$

Conclusion:

$$-\begin{pmatrix} 1 \\ 2 \\ 6 \end{pmatrix} - 9 \begin{pmatrix} -1 \\ -2 \\ -1 \end{pmatrix} = \begin{pmatrix} 8 \\ 16 \\ 3 \end{pmatrix}$$

What is the relationship between the original vectors and the matrix form of the linear equation?

Systems of Linear Equations

Continued

$$x - y = 8$$

$$2x - 2y = 16$$

$$6x - y = 3$$

matrix form
~~~~~>

$$\left( \begin{array}{cc|c} 1 & -1 & 8 \\ 2 & -2 & 16 \\ 6 & -1 & 3 \end{array} \right)$$

row reduce  
~~~~~>

$$\left(\begin{array}{cc|c} 1 & 0 & -1 \\ 0 & 1 & -9 \\ 0 & 0 & 0 \end{array} \right)$$

solution
~~~~~>

$$x = -1$$

$$y = -9$$

Conclusion:

$$-\begin{pmatrix} 1 \\ 2 \\ 6 \end{pmatrix} - 9\begin{pmatrix} -1 \\ -2 \\ -1 \end{pmatrix} = \begin{pmatrix} 8 \\ 16 \\ 3 \end{pmatrix}$$

What is the relationship between the original vectors and the matrix form of the linear equation? They have the same columns!



# Systems of Linear Equations

Continued

$$x - y = 8$$

$$2x - 2y = 16$$

$$6x - y = 3$$

matrix form  
~~~~~>

$$\left(\begin{array}{cc|c} 1 & -1 & 8 \\ 2 & -2 & 16 \\ 6 & -1 & 3 \end{array} \right)$$

row reduce
~~~~~>

$$\left( \begin{array}{cc|c} 1 & 0 & -1 \\ 0 & 1 & -9 \\ 0 & 0 & 0 \end{array} \right)$$

solution  
~~~~~>

$$x = -1$$

$$y = -9$$

Conclusion:

$$-\begin{pmatrix} 1 \\ 2 \\ 6 \end{pmatrix} - 9 \begin{pmatrix} -1 \\ -2 \\ -1 \end{pmatrix} = \begin{pmatrix} 8 \\ 16 \\ 3 \end{pmatrix}$$

What is the relationship between the original vectors and the matrix form of the linear equation? They have the same columns!

Shortcut: You can make the augmented matrix without writing down the system of linear equations first.

Vector Equations and Linear Equations

Summary

The vector equation

$$x_1 v_1 + x_2 v_2 + \cdots + x_p v_p = b,$$

where v_1, v_2, \dots, v_p, b are vectors in \mathbf{R}^n and x_1, x_2, \dots, x_p are scalars,

Vector Equations and Linear Equations

Summary

The vector equation

$$x_1 v_1 + x_2 v_2 + \cdots + x_p v_p = b,$$

where v_1, v_2, \dots, v_p, b are vectors in \mathbf{R}^n and x_1, x_2, \dots, x_p are scalars, has the same solution set as the linear system with augmented matrix

$$\left(\begin{array}{c|c|c|c|c} | & | & & | & | \\ v_1 & v_2 & \cdots & v_p & b \\ | & | & & | & | \end{array} \right),$$

where the v_i 's and b are the columns of the matrix.

Vector Equations and Linear Equations

Summary

The vector equation

$$x_1 v_1 + x_2 v_2 + \cdots + x_p v_p = b,$$

where v_1, v_2, \dots, v_p, b are vectors in \mathbf{R}^n and x_1, x_2, \dots, x_p are scalars, has the same solution set as the linear system with augmented matrix

$$\left(\begin{array}{c|c|c|c|c} | & | & & | & | \\ v_1 & v_2 & \cdots & v_p & b \\ | & | & & | & | \end{array} \right),$$

where the v_i 's and b are the columns of the matrix.

So we now have (at least) *two* equivalent ways of thinking about linear systems of equations:

Vector Equations and Linear Equations

Summary

The vector equation

$$x_1 v_1 + x_2 v_2 + \cdots + x_p v_p = b,$$

where v_1, v_2, \dots, v_p, b are vectors in \mathbf{R}^n and x_1, x_2, \dots, x_p are scalars, has the same solution set as the linear system with augmented matrix

$$\left(\begin{array}{c|c|c|c|c} | & | & & | & | \\ v_1 & v_2 & \cdots & v_p & b \\ | & | & & | & | \end{array} \right),$$

where the v_i 's and b are the columns of the matrix.

So we now have (at least) *two* equivalent ways of thinking about linear systems of equations:

1. Augmented matrices.

Vector Equations and Linear Equations

Summary

The vector equation

$$x_1 v_1 + x_2 v_2 + \cdots + x_p v_p = b,$$

where v_1, v_2, \dots, v_p, b are vectors in \mathbf{R}^n and x_1, x_2, \dots, x_p are scalars, has the same solution set as the linear system with augmented matrix

$$\left(\begin{array}{c|c|c|c|c} | & | & & | & | \\ v_1 & v_2 & \cdots & v_p & b \\ | & | & & | & | \end{array} \right),$$

where the v_i 's and b are the columns of the matrix.

So we now have (at least) *two* equivalent ways of thinking about linear systems of equations:

1. Augmented matrices.
2. Linear combinations of vectors (vector equations).

Vector Equations and Linear Equations

Summary

The vector equation

$$x_1 v_1 + x_2 v_2 + \cdots + x_p v_p = b,$$

where v_1, v_2, \dots, v_p, b are vectors in \mathbf{R}^n and x_1, x_2, \dots, x_p are scalars, has the same solution set as the linear system with augmented matrix

$$\left(\begin{array}{c|c|c|c|c} | & | & & | & | \\ v_1 & v_2 & \cdots & v_p & b \\ | & | & & | & | \end{array} \right),$$

where the v_i 's and b are the columns of the matrix.

So we now have (at least) *two* equivalent ways of thinking about linear systems of equations:

1. Augmented matrices.
2. Linear combinations of vectors (vector equations).

The last one is more geometric in nature.

Span

It is important to know what are *all* linear combinations of a set of vectors v_1, v_2, \dots, v_p in \mathbf{R}^n :

Span

It is important to know what are *all* linear combinations of a set of vectors v_1, v_2, \dots, v_p in \mathbf{R}^n : it's exactly the collection of all b in \mathbf{R}^n such that the vector equation (in the unknowns x_1, x_2, \dots, x_p)

$$x_1 v_1 + x_2 v_2 + \cdots + x_p v_p = b$$

has a solution (i.e., is consistent).

Span

It is important to know what are *all* linear combinations of a set of vectors v_1, v_2, \dots, v_p in \mathbf{R}^n : it's exactly the collection of all b in \mathbf{R}^n such that the vector equation (in the unknowns x_1, x_2, \dots, x_p)

$$x_1 v_1 + x_2 v_2 + \dots + x_p v_p = b$$

has a solution (i.e., is consistent).

Definition

Let v_1, v_2, \dots, v_p be vectors in \mathbf{R}^n . The **span** of v_1, v_2, \dots, v_p is the collection of all linear combinations of v_1, v_2, \dots, v_p , and is denoted $\text{Span}\{v_1, v_2, \dots, v_p\}$.

Span

It is important to know what are *all* linear combinations of a set of vectors v_1, v_2, \dots, v_p in \mathbf{R}^n : it's exactly the collection of all b in \mathbf{R}^n such that the vector equation (in the unknowns x_1, x_2, \dots, x_p)

$$x_1 v_1 + x_2 v_2 + \dots + x_p v_p = b$$

has a solution (i.e., is consistent).

Definition

Let v_1, v_2, \dots, v_p be vectors in \mathbf{R}^n . The **span** of v_1, v_2, \dots, v_p is the collection of all linear combinations of v_1, v_2, \dots, v_p , and is denoted $\text{Span}\{v_1, v_2, \dots, v_p\}$. In symbols:

$$\text{Span}\{v_1, v_2, \dots, v_p\} = \{ x_1 v_1 + x_2 v_2 + \dots + x_p v_p \mid x_1, x_2, \dots, x_p \text{ in } \mathbf{R} \}.$$

Span


It is important to know what are *all* linear combinations of a set of vectors v_1, v_2, \dots, v_p in \mathbf{R}^n : it's exactly the collection of all b in \mathbf{R}^n such that the vector equation (in the unknowns x_1, x_2, \dots, x_p)

$$x_1 v_1 + x_2 v_2 + \dots + x_p v_p = b$$

has a solution (i.e., is consistent).

Definition

Let v_1, v_2, \dots, v_p be vectors in \mathbf{R}^n . The **span** of v_1, v_2, \dots, v_p is the collection of all linear combinations of v_1, v_2, \dots, v_p , and is denoted $\text{Span}\{v_1, v_2, \dots, v_p\}$. In symbols:


$$\text{Span}\{v_1, v_2, \dots, v_p\} = \left\{ x_1 v_1 + x_2 v_2 + \dots + x_p v_p \mid x_1, x_2, \dots, x_p \text{ in } \mathbf{R} \right\}.$$

Span


It is important to know what are *all* linear combinations of a set of vectors v_1, v_2, \dots, v_p in \mathbf{R}^n : it's exactly the collection of all b in \mathbf{R}^n such that the vector equation (in the unknowns x_1, x_2, \dots, x_p)

$$x_1 v_1 + x_2 v_2 + \dots + x_p v_p = b$$

has a solution (i.e., is consistent).

Definition

Let v_1, v_2, \dots, v_p be vectors in \mathbf{R}^n . The **span** of v_1, v_2, \dots, v_p is the collection of all linear combinations of v_1, v_2, \dots, v_p , and is denoted $\text{Span}\{v_1, v_2, \dots, v_p\}$. In symbols:


$$\text{Span}\{v_1, v_2, \dots, v_p\} = \left\{ x_1 v_1 + x_2 v_2 + \dots + x_p v_p \mid x_1, x_2, \dots, x_p \text{ in } \mathbf{R} \right\}.$$

Synonyms: $\text{Span}\{v_1, v_2, \dots, v_p\}$ is the subset **spanned by** or **generated by** v_1, v_2, \dots, v_p .

Span


It is important to know what are *all* linear combinations of a set of vectors v_1, v_2, \dots, v_p in \mathbf{R}^n : it's exactly the collection of all b in \mathbf{R}^n such that the vector equation (in the unknowns x_1, x_2, \dots, x_p)

$$x_1 v_1 + x_2 v_2 + \dots + x_p v_p = b$$

has a solution (i.e., is consistent).

Definition

Let v_1, v_2, \dots, v_p be vectors in \mathbf{R}^n . The **span** of v_1, v_2, \dots, v_p is the collection of all linear combinations of v_1, v_2, \dots, v_p , and is denoted $\text{Span}\{v_1, v_2, \dots, v_p\}$. In symbols:


$$\text{Span}\{v_1, v_2, \dots, v_p\} = \{x_1 v_1 + x_2 v_2 + \dots + x_p v_p \mid x_1, x_2, \dots, x_p \text{ in } \mathbf{R}\}.$$

Synonyms: $\text{Span}\{v_1, v_2, \dots, v_p\}$ is the subset **spanned by** or **generated by** v_1, v_2, \dots, v_p .

This is the first of several definitions in this class that you simply **must learn**. I will give you other ways to think about Span, and ways to draw pictures, but *this is the definition*. Having a vague idea what Span means will not help you solve any exam problems!

Span

Continued

Now we have several equivalent ways of making the same statement:

Span

Continued

Now we have several equivalent ways of making the same statement:

1. A vector b is in the span of v_1, v_2, \dots, v_p .

Span

Continued

Now we have several equivalent ways of making the same statement:

1. A vector b is in the span of v_1, v_2, \dots, v_p .
2. The linear system with augmented matrix

$$\left(\begin{array}{c|c|c|c|c} | & | & & | & | \\ v_1 & v_2 & \cdots & v_p & b \\ | & | & & | & | \end{array} \right)$$

is consistent.

Span

Continued

Now we have several equivalent ways of making the same statement:

1. A vector b is in the span of v_1, v_2, \dots, v_p .
2. The linear system with augmented matrix

$$\left(\begin{array}{c|c|c|c|c} | & | & & | & \\ \hline v_1 & v_2 & \cdots & v_p & b \\ \hline | & | & & | & \end{array} \right)$$

is consistent.

3. The vector equation

$$x_1 v_1 + x_2 v_2 + \cdots + x_p v_p = b$$

has a solution.

Span

Continued

Now we have several equivalent ways of making the same statement:

1. A vector b is in the span of v_1, v_2, \dots, v_p .
2. The linear system with augmented matrix

$$\left(\begin{array}{c|c|c|c|c} | & | & & | & \\ \hline v_1 & v_2 & \cdots & v_p & b \\ \hline | & | & & | & \end{array} \right)$$

is consistent.

3. The vector equation

$$x_1 v_1 + x_2 v_2 + \cdots + x_p v_p = b$$

has a solution.


Note: **equivalent** means that, for any given list of vectors v_1, v_2, \dots, v_p, b , *either* all three statements are true, *or* all three statements are false.

Pictures of Span

Drawing a picture of $\text{Span}\{v_1, v_2, \dots, v_p\}$ is the same as drawing a picture of all linear combinations of v_1, v_2, \dots, v_p .


Pictures of Span

Drawing a picture of $\text{Span}\{v_1, v_2, \dots, v_p\}$ is the same as drawing a picture of all linear combinations of v_1, v_2, \dots, v_p .





Pictures of Span

Drawing a picture of $\text{Span}\{v_1, v_2, \dots, v_p\}$ is the same as drawing a picture of all linear combinations of v_1, v_2, \dots, v_p .





Pictures of Span

Drawing a picture of $\text{Span}\{v_1, v_2, \dots, v_p\}$ is the same as drawing a picture of all linear combinations of v_1, v_2, \dots, v_p .






Pictures of Span

Drawing a picture of $\text{Span}\{v_1, v_2, \dots, v_p\}$ is the same as drawing a picture of all linear combinations of v_1, v_2, \dots, v_p .






Pictures of Span

Drawing a picture of $\text{Span}\{v_1, v_2, \dots, v_p\}$ is the same as drawing a picture of all linear combinations of v_1, v_2, \dots, v_p .




Pictures of Span

Drawing a picture of $\text{Span}\{v_1, v_2, \dots, v_p\}$ is the same as drawing a picture of all linear combinations of v_1, v_2, \dots, v_p .




Pictures of Span

In \mathbb{R}^3





Pictures of Span

In \mathbb{R}^3





Pictures of Span

In \mathbb{R}^3






Pictures of Span

In \mathbb{R}^3






Pictures of Span

In \mathbb{R}^3







Pictures of Span

In \mathbb{R}^3







Pictures of Span

In \mathbb{R}^3



Pictures of Span

In \mathbb{R}^3



Poll

How many vectors are in $\text{Span} \left\{ \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \right\}$?

- A. Zero
- B. One
- C. Infinity

Poll

How many vectors are in $\text{Span} \left\{ \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \right\}$?

- A. Zero
- B. One
- C. Infinity

In general, it appears that $\text{Span}\{v_1, v_2, \dots, v_p\}$ is the smallest “linear space” (line, plane, etc.) containing the origin and all of the vectors v_1, v_2, \dots, v_p .

Poll

How many vectors are in $\text{Span} \left\{ \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \right\}$?

- A. Zero
- B. One
- C. Infinity

In general, it appears that $\text{Span}\{v_1, v_2, \dots, v_p\}$ is the smallest “linear space” (line, plane, etc.) containing the origin and all of the vectors v_1, v_2, \dots, v_p .

We will make this precise later.