

## Section 2.9

### Dimension and Rank

## Coefficients of Basis Vectors

**Recall:** a **basis** of a subspace  $V$  is a set of vectors that *spans*  $V$  and is *linearly independent*.

## Coefficients of Basis Vectors

**Recall:** a **basis** of a subspace  $V$  is a set of vectors that *spans*  $V$  and is *linearly independent*.

**Lemma** ← like a theorem, but less important

# Coefficients of Basis Vectors

**Recall:** a **basis** of a subspace  $V$  is a set of vectors that *spans*  $V$  and is *linearly independent*.

**Lemma**  like a theorem, but less important

If  $\mathcal{B} = \{v_1, v_2, \dots, v_m\}$  is a basis for a subspace  $V$ , then any vector  $x$  in  $V$  can be written as a linear combination

$$x = c_1 v_1 + c_2 v_2 + \cdots + c_m v_m$$

for *unique* coefficients  $c_1, c_2, \dots, c_m$ .

## Bases as Coordinate Systems

The unit coordinate vectors  $e_1, e_2, \dots, e_n$  form a basis for  $\mathbf{R}^n$ . Any vector is a unique linear combination of the  $e_i$ :

$$v = \begin{pmatrix} 3 \\ 5 \\ -2 \end{pmatrix} = 3 \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} + 5 \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} - 2 \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} = 3e_1 + 5e_2 - 2e_3.$$

## Bases as Coordinate Systems

The unit coordinate vectors  $e_1, e_2, \dots, e_n$  form a basis for  $\mathbf{R}^n$ . Any vector is a unique linear combination of the  $e_i$ :

$$v = \begin{pmatrix} 3 \\ 5 \\ -2 \end{pmatrix} = 3 \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} + 5 \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} - 2 \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} = 3e_1 + 5e_2 - 2e_3.$$

**Observe:** the *coordinates* of  $v$  are exactly the *coefficients* of  $e_1, e_2, e_3$ .

## Bases as Coordinate Systems

The unit coordinate vectors  $e_1, e_2, \dots, e_n$  form a basis for  $\mathbf{R}^n$ . Any vector is a unique linear combination of the  $e_i$ :

$$v = \begin{pmatrix} 3 \\ 5 \\ -2 \end{pmatrix} = 3 \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} + 5 \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} - 2 \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} = 3e_1 + 5e_2 - 2e_3.$$

**Observe:** the *coordinates* of  $v$  are exactly the *coefficients* of  $e_1, e_2, e_3$ .

We can go backwards: given any basis  $\mathcal{B}$ , we interpret the coefficients of a linear combination as “coordinates” with respect to  $\mathcal{B}$ .

## Bases as Coordinate Systems

The unit coordinate vectors  $e_1, e_2, \dots, e_n$  form a basis for  $\mathbf{R}^n$ . Any vector is a unique linear combination of the  $e_i$ :

$$v = \begin{pmatrix} 3 \\ 5 \\ -2 \end{pmatrix} = 3 \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} + 5 \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} - 2 \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} = 3e_1 + 5e_2 - 2e_3.$$

**Observe:** the *coordinates* of  $v$  are exactly the *coefficients* of  $e_1, e_2, e_3$ .

We can go backwards: given any basis  $\mathcal{B}$ , we interpret the coefficients of a linear combination as “coordinates” with respect to  $\mathcal{B}$ .

### Definition

Let  $\mathcal{B} = \{v_1, v_2, \dots, v_m\}$  be a basis of a subspace  $V$ . Any vector  $x$  in  $V$  can be written uniquely as a linear combination  $x = c_1 v_1 + c_2 v_2 + \dots + c_m v_m$ . The coefficients  $c_1, c_2, \dots, c_m$  are the **coordinates of  $x$  with respect to  $\mathcal{B}$** . The  **$\mathcal{B}$ -coordinate vector of  $x$**  is the vector

$$[x]_{\mathcal{B}} = \begin{pmatrix} c_1 \\ c_2 \\ \vdots \\ c_m \end{pmatrix} \quad \text{in } \mathbf{R}^m.$$



# Bases as Coordinate Systems

## Example 1

$$\text{Let } v_1 = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}, v_2 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \quad \mathcal{B} = \{v_1, v_2\}, \quad V = \text{Span}\{v_1, v_2\}.$$

# Bases as Coordinate Systems

## Example 1

Let  $v_1 = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$ ,  $v_2 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$ ,  $\mathcal{B} = \{v_1, v_2\}$ ,  $V = \text{Span}\{v_1, v_2\}$ .

Verify that  $\mathcal{B}$  is a basis:

# Bases as Coordinate Systems

## Example 1

Let  $v_1 = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$ ,  $v_2 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$ ,  $\mathcal{B} = \{v_1, v_2\}$ ,  $V = \text{Span}\{v_1, v_2\}$ .

Verify that  $\mathcal{B}$  is a basis:

Question: If  $[x]_{\mathcal{B}} = \begin{pmatrix} 5 \\ 2 \end{pmatrix}$ , then what is  $x$ ?

# Bases as Coordinate Systems

## Example 1

Let  $v_1 = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$ ,  $v_2 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$ ,  $\mathcal{B} = \{v_1, v_2\}$ ,  $V = \text{Span}\{v_1, v_2\}$ .

Verify that  $\mathcal{B}$  is a basis:

Question: If  $[x]_{\mathcal{B}} = \begin{pmatrix} 5 \\ 2 \end{pmatrix}$ , then what is  $x$ ?

Question: Find the  $\mathcal{B}$ -coordinates of  $x = \begin{pmatrix} 5 \\ 3 \\ 5 \end{pmatrix}$ .

# Bases as Coordinate Systems

## Example 2

$$\text{Let } v_1 = \begin{pmatrix} 2 \\ 3 \\ 2 \end{pmatrix}, v_2 = \begin{pmatrix} -1 \\ 1 \\ 1 \end{pmatrix}, v_3 = \begin{pmatrix} 2 \\ 8 \\ 6 \end{pmatrix}, \quad V = \text{Span}\{v_1, v_2, v_3\}.$$

# Bases as Coordinate Systems

## Example 2

$$\text{Let } v_1 = \begin{pmatrix} 2 \\ 3 \\ 2 \end{pmatrix}, v_2 = \begin{pmatrix} -1 \\ 1 \\ 1 \end{pmatrix}, v_3 = \begin{pmatrix} 2 \\ 8 \\ 6 \end{pmatrix}, \quad V = \text{Span}\{v_1, v_2, v_3\}.$$

**Question:** Find a basis for  $V$ .

# Bases as Coordinate Systems

## Example 2

Let  $v_1 = \begin{pmatrix} 2 \\ 3 \\ 2 \end{pmatrix}$ ,  $v_2 = \begin{pmatrix} -1 \\ 1 \\ 1 \end{pmatrix}$ ,  $v_3 = \begin{pmatrix} 2 \\ 8 \\ 6 \end{pmatrix}$ ,  $V = \text{Span}\{v_1, v_2, v_3\}$ .

**Question:** Find a basis for  $V$ .

**Question:** Find the  $\mathcal{B}$ -coordinates of  $x = \begin{pmatrix} 4 \\ 11 \\ 8 \end{pmatrix}$ .

# Bases as Coordinate Systems

## Summary

If  $\mathcal{B} = \{v_1, v_2, \dots, v_m\}$  is a basis for a subspace  $V$  and  $x$  is in  $V$ , then

$$[x]_{\mathcal{B}} = \begin{pmatrix} c_1 \\ c_2 \\ \vdots \\ c_m \end{pmatrix} \quad \text{means} \quad x = c_1 v_1 + c_2 v_2 + \cdots + c_m v_m.$$



# Bases as Coordinate Systems

## Summary

If  $\mathcal{B} = \{v_1, v_2, \dots, v_m\}$  is a basis for a subspace  $V$  and  $x$  is in  $V$ , then

$$[x]_{\mathcal{B}} = \begin{pmatrix} c_1 \\ c_2 \\ \vdots \\ c_m \end{pmatrix} \quad \text{means} \quad x = c_1 v_1 + c_2 v_2 + \cdots + c_m v_m.$$

Finding the  $\mathcal{B}$ -coordinates for  $x$  means solving the vector equation

$$x = c_1 v_1 + c_2 v_2 + \cdots + c_m v_m$$

in the unknowns  $c_1, c_2, \dots, c_m$ .

# Bases as Coordinate Systems

## Summary

If  $\mathcal{B} = \{v_1, v_2, \dots, v_m\}$  is a basis for a subspace  $V$  and  $x$  is in  $V$ , then

$$[x]_{\mathcal{B}} = \begin{pmatrix} c_1 \\ c_2 \\ \vdots \\ c_m \end{pmatrix} \quad \text{means} \quad x = c_1 v_1 + c_2 v_2 + \cdots + c_m v_m.$$

Finding the  $\mathcal{B}$ -coordinates for  $x$  means solving the vector equation

$$x = c_1 v_1 + c_2 v_2 + \cdots + c_m v_m$$

in the unknowns  $c_1, c_2, \dots, c_m$ . This (usually) means row reducing the augmented matrix

$$\left( \begin{array}{c|c|c|c|c} | & | & & | & | \\ v_1 & v_2 & \cdots & v_m & x \\ | & | & & | & | \end{array} \right).$$

# Bases as Coordinate Systems

## Summary

If  $\mathcal{B} = \{v_1, v_2, \dots, v_m\}$  is a basis for a subspace  $V$  and  $x$  is in  $V$ , then

$$[x]_{\mathcal{B}} = \begin{pmatrix} c_1 \\ c_2 \\ \vdots \\ c_m \end{pmatrix} \quad \text{means} \quad x = c_1 v_1 + c_2 v_2 + \cdots + c_m v_m.$$

Finding the  $\mathcal{B}$ -coordinates for  $x$  means solving the vector equation

$$x = c_1 v_1 + c_2 v_2 + \cdots + c_m v_m$$

in the unknowns  $c_1, c_2, \dots, c_m$ . This (usually) means row reducing the augmented matrix

$$\left( \begin{array}{c|c|c|c|c} | & | & & | & | \\ v_1 & v_2 & \cdots & v_m & x \\ | & | & & | & | \end{array} \right).$$

**Question:** What happens if you try to find the  $\mathcal{B}$ -coordinates of  $x$  *not* in  $V$ ?

# Bases as Coordinate Systems

Picture


Let

$$v_1 = \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \quad v_2 = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$$

These form a basis  $\mathcal{B}$  for the plane

$$V = \text{Span}\{v_1, v_2\}$$

in  $\mathbf{R}^3$ .



# Bases as Coordinate Systems

Picture


Let

$$v_1 = \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \quad v_2 = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$$

These form a basis  $\mathcal{B}$  for the plane

$$V = \text{Span}\{v_1, v_2\}$$

in  $\mathbf{R}^3$ .



**Question:** Estimate the  $\mathcal{B}$ -coordinates of these vectors:

$$[u_1]_{\mathcal{B}} =$$

$$[u_2]_{\mathcal{B}} =$$

$$[u_3]_{\mathcal{B}} =$$

$$[u_4]_{\mathcal{B}} =$$

# Bases as Coordinate Systems

Picture


Let

$$v_1 = \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} \quad v_2 = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$$

These form a basis  $\mathcal{B}$  for the plane

$$V = \text{Span}\{v_1, v_2\}$$

in  $\mathbf{R}^3$ .



**Question:** Estimate the  $\mathcal{B}$ -coordinates of these vectors:

$$[u_1]_{\mathcal{B}} =$$

$$[u_2]_{\mathcal{B}} =$$

$$[u_3]_{\mathcal{B}} =$$

$$[u_4]_{\mathcal{B}} =$$

## Remark

Many of you want to think of a plane in  $\mathbf{R}^3$  as “being”  $\mathbf{R}^2$ . Choosing a basis  $\mathcal{B}$  and using  $\mathcal{B}$ -coordinates is one way to make sense of that. But remember that the coordinates are the coefficients of a linear combination of the basis vectors.

# The Rank Theorem

Recall:

- ▶ The **dimension** of a subspace  $V$  is the number of vectors in a basis for  $V$ .

# The Rank Theorem

Recall:

- ▶ The **dimension** of a subspace  $V$  is the number of vectors in a basis for  $V$ .
- ▶ A basis for the column space of a matrix  $A$  is given by the



# The Rank Theorem

Recall:

- ▶ The **dimension** of a subspace  $V$  is the number of vectors in a basis for  $V$ .
- ▶ A basis for the column space of a matrix  $A$  is given by the pivot columns.

# The Rank Theorem

Recall:

- ▶ The **dimension** of a subspace  $V$  is the number of vectors in a basis for  $V$ .
- ▶ A basis for the column space of a matrix  $A$  is given by the pivot columns.
- ▶ A basis for the null space of  $A$  is given by the

# The Rank Theorem

## Recall:

- ▶ The **dimension** of a subspace  $V$  is the number of vectors in a basis for  $V$ .
- ▶ A basis for the column space of a matrix  $A$  is given by the pivot columns.
- ▶ A basis for the null space of  $A$  is given by the vectors attached to the free variables in the parametric vector form.

# The Rank Theorem

## Recall:

- ▶ The **dimension** of a subspace  $V$  is the number of vectors in a basis for  $V$ .
- ▶ A basis for the column space of a matrix  $A$  is given by the pivot columns.
- ▶ A basis for the null space of  $A$  is given by the vectors attached to the free variables in the parametric vector form.

## Definition

The **rank** of a matrix  $A$ , written  $\text{rank } A$ , is the dimension of the column space  $\text{Col } A$ .

# The Rank Theorem

## Recall:

- ▶ The **dimension** of a subspace  $V$  is the number of vectors in a basis for  $V$ .
- ▶ A basis for the column space of a matrix  $A$  is given by the pivot columns.
- ▶ A basis for the null space of  $A$  is given by the vectors attached to the free variables in the parametric vector form.

## Definition

The **rank** of a matrix  $A$ , written  $\text{rank } A$ , is the dimension of the column space  $\text{Col } A$ .

## Observe:

$$\text{rank } A = \dim \text{Col } A =$$

# The Rank Theorem

## Recall:

- ▶ The **dimension** of a subspace  $V$  is the number of vectors in a basis for  $V$ .
- ▶ A basis for the column space of a matrix  $A$  is given by the pivot columns.
- ▶ A basis for the null space of  $A$  is given by the vectors attached to the free variables in the parametric vector form.

## Definition

The **rank** of a matrix  $A$ , written  $\text{rank } A$ , is the dimension of the column space  $\text{Col } A$ .

## Observe:

$\text{rank } A = \dim \text{Col } A = \text{the number of columns with pivots}$

# The Rank Theorem

## Recall:

- ▶ The **dimension** of a subspace  $V$  is the number of vectors in a basis for  $V$ .
- ▶ A basis for the column space of a matrix  $A$  is given by the pivot columns.
- ▶ A basis for the null space of  $A$  is given by the vectors attached to the free variables in the parametric vector form.

## Definition

The **rank** of a matrix  $A$ , written  $\text{rank } A$ , is the dimension of the column space  $\text{Col } A$ .

## Observe:

$$\begin{aligned}\text{rank } A &= \dim \text{Col } A = \text{the number of columns with pivots} \\ \dim \text{Nul } A &= \end{aligned}$$

# The Rank Theorem

## Recall:

- ▶ The **dimension** of a subspace  $V$  is the number of vectors in a basis for  $V$ .
- ▶ A basis for the column space of a matrix  $A$  is given by the pivot columns.
- ▶ A basis for the null space of  $A$  is given by the vectors attached to the free variables in the parametric vector form.

## Definition

The **rank** of a matrix  $A$ , written  $\text{rank } A$ , is the dimension of the column space  $\text{Col } A$ .

## Observe:

$$\begin{aligned}\text{rank } A &= \dim \text{Col } A = \text{the number of columns with pivots} \\ \dim \text{Nul } A &= \text{the number of free variables}\end{aligned}$$



# The Rank Theorem

## Recall:

- ▶ The **dimension** of a subspace  $V$  is the number of vectors in a basis for  $V$ .
- ▶ A basis for the column space of a matrix  $A$  is given by the pivot columns.
- ▶ A basis for the null space of  $A$  is given by the vectors attached to the free variables in the parametric vector form.

## Definition

The **rank** of a matrix  $A$ , written  $\text{rank } A$ , is the dimension of the column space  $\text{Col } A$ .

## Observe:

$$\begin{aligned}\text{rank } A &= \dim \text{Col } A = \text{the number of columns with pivots} \\ \dim \text{Nul } A &= \text{the number of free variables} \\ &= \text{the number of columns without pivots.}\end{aligned}$$

# The Rank Theorem

## Recall:

- ▶ The **dimension** of a subspace  $V$  is the number of vectors in a basis for  $V$ .
- ▶ A basis for the column space of a matrix  $A$  is given by the pivot columns.
- ▶ A basis for the null space of  $A$  is given by the vectors attached to the free variables in the parametric vector form.

## Definition

The **rank** of a matrix  $A$ , written  $\text{rank } A$ , is the dimension of the column space  $\text{Col } A$ .

## Observe:

$$\begin{aligned}\text{rank } A &= \dim \text{Col } A = \text{the number of columns with pivots} \\ \dim \text{Nul } A &= \text{the number of free variables} \\ &= \text{the number of columns without pivots.}\end{aligned}$$

## Rank Theorem

If  $A$  is an  $m \times n$  matrix, then

$$\text{rank } A + \dim \text{Nul } A =$$

# The Rank Theorem

## Recall:

- ▶ The **dimension** of a subspace  $V$  is the number of vectors in a basis for  $V$ .
- ▶ A basis for the column space of a matrix  $A$  is given by the pivot columns.
- ▶ A basis for the null space of  $A$  is given by the vectors attached to the free variables in the parametric vector form.

## Definition

The **rank** of a matrix  $A$ , written  $\text{rank } A$ , is the dimension of the column space  $\text{Col } A$ .

## Observe:

$$\begin{aligned}\text{rank } A &= \dim \text{Col } A = \text{the number of columns with pivots} \\ \dim \text{Nul } A &= \text{the number of free variables} \\ &= \text{the number of columns without pivots.}\end{aligned}$$

## Rank Theorem

If  $A$  is an  $m \times n$  matrix, then

$$\text{rank } A + \dim \text{Nul } A = n = \text{the number of columns of } A.$$

# The Rank Theorem


## Example

$$A = \begin{pmatrix} 1 & 2 & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix} \xrightarrow{\text{rref}} \begin{pmatrix} 1 & 0 & -8 & -7 \\ 0 & 1 & 4 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

# The Rank Theorem

## Example


$$A = \begin{pmatrix} \boxed{1} & \boxed{2} & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix} \xrightarrow{\text{rref}} \begin{pmatrix} 1 & 0 & -8 & -7 \\ 0 & 1 & 4 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

  
basis of Col A

# The Rank Theorem

## Example

$$A = \begin{pmatrix} \boxed{1} & \boxed{2} & 0 & -1 \\ -2 & -3 & 4 & 5 \\ 2 & 4 & 0 & -2 \end{pmatrix} \xrightarrow{\text{rref}} \begin{pmatrix} 1 & 0 & \boxed{-8} & \boxed{-7} \\ 0 & 1 & 4 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

  
basis of Col A                      free variables

## Poll

Let  $A$  and  $B$  be  $3 \times 3$  matrices. Suppose that  $\text{rank}(A) = 2$  and  $\text{rank}(B) = 2$ . Is it possible that  $AB = 0$ ? Why or why not?

## Poll

Let  $A$  and  $B$  be  $3 \times 3$  matrices. Suppose that  $\text{rank}(A) = 2$  and  $\text{rank}(B) = 2$ . Is it possible that  $AB = 0$ ? Why or why not?

If  $AB = 0$ , then  $ABx = 0$  for every  $x$  in  $\mathbf{R}^3$ .



## Poll

Let  $A$  and  $B$  be  $3 \times 3$  matrices. Suppose that  $\text{rank}(A) = 2$  and  $\text{rank}(B) = 2$ . Is it possible that  $AB = 0$ ? Why or why not?

If  $AB = 0$ , then  $ABx = 0$  for every  $x$  in  $\mathbf{R}^3$ .

This means  $A(Bx) = 0$ , so  $Bx$  is in  $\text{Nul } A$ .

## Poll

Let  $A$  and  $B$  be  $3 \times 3$  matrices. Suppose that  $\text{rank}(A) = 2$  and  $\text{rank}(B) = 2$ . Is it possible that  $AB = 0$ ? Why or why not?

If  $AB = 0$ , then  $ABx = 0$  for every  $x$  in  $\mathbf{R}^3$ .

This means  $A(Bx) = 0$ , so  $Bx$  is in  $\text{Nul } A$ .

This is true for every  $x$ , so  $\text{Col } B$  is contained in  $\text{Nul } A$ .

## Poll

Let  $A$  and  $B$  be  $3 \times 3$  matrices. Suppose that  $\text{rank}(A) = 2$  and  $\text{rank}(B) = 2$ . Is it possible that  $AB = 0$ ? Why or why not?

If  $AB = 0$ , then  $ABx = 0$  for every  $x$  in  $\mathbf{R}^3$ .

This means  $A(Bx) = 0$ , so  $Bx$  is in  $\text{Nul } A$ .

This is true for every  $x$ , so  $\text{Col } B$  is contained in  $\text{Nul } A$ .

But  $\dim \text{Nul } A = 1$  and  $\dim \text{Col } B = 2$ , and a 1-dimensional space can't contain a 2-dimensional space.

## Poll

Let  $A$  and  $B$  be  $3 \times 3$  matrices. Suppose that  $\text{rank}(A) = 2$  and  $\text{rank}(B) = 2$ . Is it possible that  $AB = 0$ ? Why or why not?

If  $AB = 0$ , then  $ABx = 0$  for every  $x$  in  $\mathbf{R}^3$ .

This means  $A(Bx) = 0$ , so  $Bx$  is in  $\text{Nul } A$ .

This is true for every  $x$ , so  $\text{Col } B$  is contained in  $\text{Nul } A$ .

But  $\dim \text{Nul } A = 1$  and  $\dim \text{Col } B = 2$ , and a 1-dimensional space can't contain a 2-dimensional space.

Hence it can't happen.

## Poll

Let  $A$  and  $B$  be  $3 \times 3$  matrices. Suppose that  $\text{rank}(A) = 2$  and  $\text{rank}(B) = 2$ . Is it possible that  $AB = 0$ ? Why or why not?


If  $AB = 0$ , then  $ABx = 0$  for every  $x$  in  $\mathbf{R}^3$ .

This means  $A(Bx) = 0$ , so  $Bx$  is in  $\text{Nul } A$ .


This is true for every  $x$ , so  $\text{Col } B$  is contained in  $\text{Nul } A$ .

But  $\dim \text{Nul } A = 1$  and  $\dim \text{Col } B = 2$ , and a 1-dimensional space can't contain a 2-dimensional space.

Hence it can't happen.



does not  
contain



# The Basis Theorem

## Basis Theorem

Let  $V$  be a subspace of dimension  $m$ . Then:

# The Basis Theorem

## Basis Theorem

Let  $V$  be a subspace of dimension  $m$ . Then:

- ▶ Any  $m$  linearly independent vectors in  $V$  form a basis for  $V$ .

# The Basis Theorem

## Basis Theorem

Let  $V$  be a subspace of dimension  $m$ . Then:

- ▶ Any  $m$  linearly independent vectors in  $V$  form a basis for  $V$ .
- ▶ Any  $m$  vectors that span  $V$  form a basis for  $V$ .



# The Basis Theorem

## Basis Theorem

Let  $V$  be a subspace of dimension  $m$ . Then:

- ▶ Any  $m$  linearly independent vectors in  $V$  form a basis for  $V$ .
- ▶ Any  $m$  vectors that span  $V$  form a basis for  $V$ .

### Upshot

If you *already* know that  $\dim V = m$ , and you have  $m$  vectors  $\mathcal{B} = \{v_1, v_2, \dots, v_m\}$  in  $V$ , then you only have to check *one* of

# The Basis Theorem

## Basis Theorem

Let  $V$  be a subspace of dimension  $m$ . Then:

- ▶ Any  $m$  linearly independent vectors in  $V$  form a basis for  $V$ .
- ▶ Any  $m$  vectors that span  $V$  form a basis for  $V$ .

### Upshot

If you *already* know that  $\dim V = m$ , and you have  $m$  vectors  $\mathcal{B} = \{v_1, v_2, \dots, v_m\}$  in  $V$ , then you only have to check *one* of

1.  $\mathcal{B}$  is linearly independent, *or*

# The Basis Theorem

## Basis Theorem

Let  $V$  be a subspace of dimension  $m$ . Then:

- ▶ Any  $m$  linearly independent vectors in  $V$  form a basis for  $V$ .
- ▶ Any  $m$  vectors that span  $V$  form a basis for  $V$ .

### Upshot

If you *already* know that  $\dim V = m$ , and you have  $m$  vectors  $\mathcal{B} = \{v_1, v_2, \dots, v_m\}$  in  $V$ , then you only have to check *one* of

1.  $\mathcal{B}$  is linearly independent, *or*
2.  $\mathcal{B}$  spans  $V$

# The Basis Theorem

## Basis Theorem

Let  $V$  be a subspace of dimension  $m$ . Then:

- ▶ Any  $m$  linearly independent vectors in  $V$  form a basis for  $V$ .
- ▶ Any  $m$  vectors that span  $V$  form a basis for  $V$ .

### Upshot

If you *already* know that  $\dim V = m$ , and you have  $m$  vectors  $\mathcal{B} = \{v_1, v_2, \dots, v_m\}$  in  $V$ , then you only have to check *one* of

1.  $\mathcal{B}$  is linearly independent, *or*
2.  $\mathcal{B}$  spans  $V$

in order for  $\mathcal{B}$  to be a basis.

# The Invertible Matrix Theorem

## Addenda

### The Invertible Matrix Theorem

Let  $A$  be an  $n \times n$  matrix, and let  $T: \mathbf{R}^n \rightarrow \mathbf{R}^n$  be the linear transformation  $T(x) = Ax$ . The following statements are equivalent.

1.  $A$  is invertible.

# The Invertible Matrix Theorem

## Addenda

### The Invertible Matrix Theorem

Let  $A$  be an  $n \times n$  matrix, and let  $T: \mathbf{R}^n \rightarrow \mathbf{R}^n$  be the linear transformation  $T(x) = Ax$ . The following statements are equivalent.

1.  $A$  is invertible.
2.  $T$  is invertible.
3.  $A$  is row equivalent to  $I_n$ .
4.  $A$  has  $n$  pivots.
5.  $Ax = 0$  has only the trivial solution.
6. The columns of  $A$  are linearly independent.
7.  $T$  is one-to-one.
8.  $Ax = b$  is consistent for all  $b$  in  $\mathbf{R}^n$ .
9. The columns of  $A$  span  $\mathbf{R}^n$ .
10.  $T$  is onto.
11.  $A$  has a left inverse (there exists  $B$  such that  $BA = I_n$ ).
12.  $A$  has a right inverse (there exists  $B$  such that  $AB = I_n$ ).
13.  $A^T$  is invertible.

# The Invertible Matrix Theorem

## Addenda

### The Invertible Matrix Theorem

Let  $A$  be an  $n \times n$  matrix, and let  $T: \mathbf{R}^n \rightarrow \mathbf{R}^n$  be the linear transformation  $T(x) = Ax$ . The following statements are equivalent.

1.  $A$  is invertible.
2.  $T$  is invertible.
3.  $A$  is row equivalent to  $I_n$ .
4.  $A$  has  $n$  pivots.
5.  $Ax = 0$  has only the trivial solution.
6. The columns of  $A$  are linearly independent.
7.  $T$  is one-to-one.
8.  $Ax = b$  is consistent for all  $b$  in  $\mathbf{R}^n$ .
9. The columns of  $A$  span  $\mathbf{R}^n$ .
10.  $T$  is onto.
11.  $A$  has a left inverse (there exists  $B$  such that  $BA = I_n$ ).
12.  $A$  has a right inverse (there exists  $B$  such that  $AB = I_n$ ).
13.  $A^T$  is invertible.
14. The columns of  $A$  form a basis for  $\mathbf{R}^n$ .

# The Invertible Matrix Theorem

## Addenda

### The Invertible Matrix Theorem

Let  $A$  be an  $n \times n$  matrix, and let  $T: \mathbf{R}^n \rightarrow \mathbf{R}^n$  be the linear transformation  $T(x) = Ax$ . The following statements are equivalent.

1.  $A$  is invertible.
2.  $T$  is invertible.
3.  $A$  is row equivalent to  $I_n$ .
4.  $A$  has  $n$  pivots.
5.  $Ax = 0$  has only the trivial solution.
6. The columns of  $A$  are linearly independent.
7.  $T$  is one-to-one.
8.  $Ax = b$  is consistent for all  $b$  in  $\mathbf{R}^n$ .
9. The columns of  $A$  span  $\mathbf{R}^n$ .
10.  $T$  is onto.
11.  $A$  has a left inverse (there exists  $B$  such that  $BA = I_n$ ).
12.  $A$  has a right inverse (there exists  $B$  such that  $AB = I_n$ ).
13.  $A^T$  is invertible.
14. The columns of  $A$  form a basis for  $\mathbf{R}^n$ .
15.  $\text{Col } A = \mathbf{R}^n$ .



# The Invertible Matrix Theorem

## Addenda

### The Invertible Matrix Theorem

Let  $A$  be an  $n \times n$  matrix, and let  $T: \mathbf{R}^n \rightarrow \mathbf{R}^n$  be the linear transformation  $T(x) = Ax$ . The following statements are equivalent.

1.  $A$  is invertible.
2.  $T$  is invertible.
3.  $A$  is row equivalent to  $I_n$ .
4.  $A$  has  $n$  pivots.
5.  $Ax = 0$  has only the trivial solution.
6. The columns of  $A$  are linearly independent.
7.  $T$  is one-to-one.
8.  $Ax = b$  is consistent for all  $b$  in  $\mathbf{R}^n$ .
9. The columns of  $A$  span  $\mathbf{R}^n$ .
10.  $T$  is onto.
11.  $A$  has a left inverse (there exists  $B$  such that  $BA = I_n$ ).
12.  $A$  has a right inverse (there exists  $B$  such that  $AB = I_n$ ).
13.  $A^T$  is invertible.
14. The columns of  $A$  form a basis for  $\mathbf{R}^n$ .
15.  $\text{Col } A = \mathbf{R}^n$ .
16.  $\dim \text{Col } A = n$ .

# The Invertible Matrix Theorem

## Addenda

### The Invertible Matrix Theorem

Let  $A$  be an  $n \times n$  matrix, and let  $T: \mathbf{R}^n \rightarrow \mathbf{R}^n$  be the linear transformation  $T(x) = Ax$ . The following statements are equivalent.

1.  $A$  is invertible.
2.  $T$  is invertible.
3.  $A$  is row equivalent to  $I_n$ .
4.  $A$  has  $n$  pivots.
5.  $Ax = 0$  has only the trivial solution.
6. The columns of  $A$  are linearly independent.
7.  $T$  is one-to-one.
8.  $Ax = b$  is consistent for all  $b$  in  $\mathbf{R}^n$ .
9. The columns of  $A$  span  $\mathbf{R}^n$ .
10.  $T$  is onto.
11.  $A$  has a left inverse (there exists  $B$  such that  $BA = I_n$ ).
12.  $A$  has a right inverse (there exists  $B$  such that  $AB = I_n$ ).
13.  $A^T$  is invertible.
14. The columns of  $A$  form a basis for  $\mathbf{R}^n$ .
15.  $\text{Col } A = \mathbf{R}^n$ .
16.  $\dim \text{Col } A = n$ .
17.  $\text{rank } A = n$ .

# The Invertible Matrix Theorem

## Addenda

### The Invertible Matrix Theorem

Let  $A$  be an  $n \times n$  matrix, and let  $T: \mathbf{R}^n \rightarrow \mathbf{R}^n$  be the linear transformation  $T(x) = Ax$ . The following statements are equivalent.

1.  $A$  is invertible.
2.  $T$  is invertible.
3.  $A$  is row equivalent to  $I_n$ .
4.  $A$  has  $n$  pivots.
5.  $Ax = 0$  has only the trivial solution.
6. The columns of  $A$  are linearly independent.
7.  $T$  is one-to-one.
8.  $Ax = b$  is consistent for all  $b$  in  $\mathbf{R}^n$ .
9. The columns of  $A$  span  $\mathbf{R}^n$ .
10.  $T$  is onto.
11.  $A$  has a left inverse (there exists  $B$  such that  $BA = I_n$ ).
12.  $A$  has a right inverse (there exists  $B$  such that  $AB = I_n$ ).
13.  $A^T$  is invertible.
14. The columns of  $A$  form a basis for  $\mathbf{R}^n$ .
15.  $\text{Col } A = \mathbf{R}^n$ .
16.  $\dim \text{Col } A = n$ .
17.  $\text{rank } A = n$ .
18.  $\text{Nul } A = \{0\}$ .

# The Invertible Matrix Theorem

## Addenda

### The Invertible Matrix Theorem

Let  $A$  be an  $n \times n$  matrix, and let  $T: \mathbf{R}^n \rightarrow \mathbf{R}^n$  be the linear transformation  $T(x) = Ax$ . The following statements are equivalent.

1.  $A$  is invertible.
2.  $T$  is invertible.
3.  $A$  is row equivalent to  $I_n$ .
4.  $A$  has  $n$  pivots.
5.  $Ax = 0$  has only the trivial solution.
6. The columns of  $A$  are linearly independent.
7.  $T$  is one-to-one.
8.  $Ax = b$  is consistent for all  $b$  in  $\mathbf{R}^n$ .
9. The columns of  $A$  span  $\mathbf{R}^n$ .
10.  $T$  is onto.
11.  $A$  has a left inverse (there exists  $B$  such that  $BA = I_n$ ).
12.  $A$  has a right inverse (there exists  $B$  such that  $AB = I_n$ ).
13.  $A^T$  is invertible.
14. The columns of  $A$  form a basis for  $\mathbf{R}^n$ .
15.  $\text{Col } A = \mathbf{R}^n$ .
16.  $\dim \text{Col } A = n$ .
17.  $\text{rank } A = n$ .
18.  $\text{Nul } A = \{0\}$ .
19.  $\dim \text{Nul } A = 0$ .

# The Invertible Matrix Theorem

## Addenda

### The Invertible Matrix Theorem

Let  $A$  be an  $n \times n$  matrix, and let  $T: \mathbf{R}^n \rightarrow \mathbf{R}^n$  be the linear transformation  $T(x) = Ax$ . The following statements are equivalent.

1.  $A$  is invertible.
2.  $T$  is invertible.
3.  $A$  is row equivalent to  $I_n$ .
4.  $A$  has  $n$  pivots.
5.  $Ax = 0$  has only the trivial solution.
6. The columns of  $A$  are linearly independent.
7.  $T$  is one-to-one.
8.  $Ax = b$  is consistent for all  $b$  in  $\mathbf{R}^n$ .
9. The columns of  $A$  span  $\mathbf{R}^n$ .
10.  $T$  is onto.
11.  $A$  has a left inverse (there exists  $B$  such that  $BA = I_n$ ).
12.  $A$  has a right inverse (there exists  $B$  such that  $AB = I_n$ ).
13.  $A^T$  is invertible.
14. The columns of  $A$  form a basis for  $\mathbf{R}^n$ .
15.  $\text{Col } A = \mathbf{R}^n$ .
16.  $\dim \text{Col } A = n$ .
17.  $\text{rank } A = n$ .
18.  $\text{Nul } A = \{0\}$ .
19.  $\dim \text{Nul } A = 0$ .

These are equivalent to the previous conditions by the Rank Theorem and the Basis Theorem.