

# Section 1.3

## Vector Equations


# Motivation

We want to think about the *algebra* in linear algebra (systems of equations and their solution sets) in terms of *geometry* (points, lines, planes, etc).

# Motivation

We want to think about the *algebra* in linear algebra (systems of equations and their solution sets) in terms of *geometry* (points, lines, planes, etc).


$$\begin{array}{rcl} x - 3y & = & -3 \\ 2x + y & = & 8 \end{array}$$



# Motivation

We want to think about the *algebra* in linear algebra (systems of equations and their solution sets) in terms of *geometry* (points, lines, planes, etc).

$$\begin{array}{rcl} x - 3y & = & -3 \\ 2x + y & = & 8 \end{array}$$




This will give us better insight into the properties of systems of equations and their solution sets.

# Motivation

We want to think about the *algebra* in linear algebra (systems of equations and their solution sets) in terms of *geometry* (points, lines, planes, etc).

$$\begin{array}{rcl} x - 3y & = & -3 \\ 2x + y & = & 8 \end{array}$$



This will give us better insight into the properties of systems of equations and their solution sets.

To do this, we need to introduce  $n$ -dimensional space  $\mathbf{R}^n$ , and **vectors** inside it.

## Line, Plane, Space, ...

Recall that  $\mathbf{R}$  denotes the collection of all real numbers, i.e. the number line.

Recall that  $\mathbf{R}$  denotes the collection of all real numbers, i.e. the number line.

### Definition

Let  $n$  be a positive whole number. We define

$$\mathbf{R}^n = \text{all ordered } n\text{-tuples of real numbers } (x_1, x_2, x_3, \dots, x_n).$$

## Line, Plane, Space, ...

Recall that  $\mathbf{R}$  denotes the collection of all real numbers, i.e. the number line.


### Definition

Let  $n$  be a positive whole number. We define

$$\mathbf{R}^n = \text{all ordered } n\text{-tuples of real numbers } (x_1, x_2, x_3, \dots, x_n).$$

### Example

When  $n = 1$ , we just get  $\mathbf{R}$  back:  $\mathbf{R}^1 = \mathbf{R}$ . Geometrically, this is the *number line*.






# Line, Plane, Space, ...

Continued

## Example

When  $n = 2$ , we can think of  $\mathbf{R}^2$  as the *plane*. This is because every point on the plane can be represented by an ordered pair of real numbers, namely, its  $x$ - and  $y$ -coordinates.




# Line, Plane, Space, ...

Continued

## Example

When  $n = 2$ , we can think of  $\mathbf{R}^2$  as the *plane*. This is because every point on the plane can be represented by an ordered pair of real numbers, namely, its  $x$ - and  $y$ -coordinates.




We can use the elements of  $\mathbf{R}^2$  to *label* points on the plane, but  $\mathbf{R}^2$  is not defined to be the plane!

# Line, Plane, Space, ...

Continued

## Example

When  $n = 3$ , we can think of  $\mathbf{R}^3$  as the *space* we (appear to) live in. This is because every point in space can be represented by an ordered triple of real numbers, namely, its  $x$ -,  $y$ -, and  $z$ -coordinates.




# Line, Plane, Space, ...

Continued

## Example

When  $n = 3$ , we can think of  $\mathbf{R}^3$  as the *space* we (appear to) live in. This is because every point in space can be represented by an ordered triple of real numbers, namely, its  $x$ -,  $y$ -, and  $z$ -coordinates.



Again, we can use the elements of  $\mathbf{R}^3$  to *label* points in space, but  $\mathbf{R}^3$  is not defined to be space!

# Line, Plane, Space, ...

Continued

So what is  $\mathbf{R}^4$ ? or  $\mathbf{R}^5$ ? or  $\mathbf{R}^n$ ?

# Line, Plane, Space, ...

Continued

So what is  $\mathbf{R}^4$ ? or  $\mathbf{R}^5$ ? or  $\mathbf{R}^n$ ?

...go back to the *definition*: ordered  $n$ -tuples of real numbers

$$(x_1, x_2, x_3, \dots, x_n).$$

# Line, Plane, Space, ...

Continued

So what is  $\mathbf{R}^4$ ? or  $\mathbf{R}^5$ ? or  $\mathbf{R}^n$ ?

...go back to the *definition*: ordered  $n$ -tuples of real numbers

$$(x_1, x_2, x_3, \dots, x_n).$$

They're still “geometric” spaces, in the sense that our intuition for  $\mathbf{R}^2$  and  $\mathbf{R}^3$  sometimes extends to  $\mathbf{R}^n$ , but they're harder to visualize.

# Line, Plane, Space, ...

Continued

So what is  $\mathbf{R}^4$ ? or  $\mathbf{R}^5$ ? or  $\mathbf{R}^n$ ?

...go back to the *definition*: ordered  $n$ -tuples of real numbers

$$(x_1, x_2, x_3, \dots, x_n).$$

They're still “geometric” spaces, in the sense that our intuition for  $\mathbf{R}^2$  and  $\mathbf{R}^3$  sometimes extends to  $\mathbf{R}^n$ , but they're harder to visualize.

We'll make definitions and state theorems that apply to any  $\mathbf{R}^n$ , but we'll only draw pictures for  $\mathbf{R}^2$  and  $\mathbf{R}^3$ .




# Vectors

In the previous slides, we were thinking of elements of  $\mathbf{R}^n$  as **points**: in the line, plane, space, etc.

# Vectors

In the previous slides, we were thinking of elements of  $\mathbf{R}^n$  as **points**: in the line, plane, space, etc.


We can also think of them as **vectors**: arrows with a given length and direction.



# Vectors

In the previous slides, we were thinking of elements of  $\mathbf{R}^n$  as **points**: in the line, plane, space, etc.

We can also think of them as **vectors**: arrows with a given length and direction.




So the vector points *horizontally* in the amount of its x-coordinate, and *vertically* in the amount of its y-coordinate.

# Vectors

In the previous slides, we were thinking of elements of  $\mathbf{R}^n$  as **points**: in the line, plane, space, etc.

We can also think of them as **vectors**: arrows with a given length and direction.



So the vector points *horizontally* in the amount of its  $x$ -coordinate, and *vertically* in the amount of its  $y$ -coordinate.

When we think of an element of  $\mathbf{R}^n$  as a vector, we write it as a matrix with  $n$  rows and one column:

$$v = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}.$$

We'll see why this is useful later.


## Points and Vectors

So what is the difference between a point and a vector?

## Points and Vectors

So what is the difference between a point and a vector?

A vector need not start at the origin: *it can be located anywhere!* In other words, an arrow is determined by its length and its direction, not by its location.




These arrows all represent the vector  $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$ .

## Points and Vectors

So what is the difference between a point and a vector?

A vector need not start at the origin: *it can be located anywhere!* In other words, an arrow is determined by its length and its direction, not by its location.




These arrows all represent the vector  $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$ .

However, unless otherwise specified, we'll assume a vector starts at the origin: we'll usually be sloppy and identify the vector  $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$  with the point (1, 2).

## Points and Vectors

So what is the difference between a point and a vector?

A vector need not start at the origin: *it can be located anywhere!* In other words, an arrow is determined by its length and its direction, not by its location.



These arrows all represent the vector  $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$ .

However, unless otherwise specified, we'll assume a vector starts at the origin: we'll usually be sloppy and identify the vector  $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$  with the point (1, 2).


This makes sense in the real world: many physical quantities, such as velocity, are represented as vectors. But it makes more sense to think of the velocity of a car as being located at the car.



## Points and Vectors

So what is the difference between a point and a vector?

A vector need not start at the origin: *it can be located anywhere!* In other words, an arrow is determined by its length and its direction, not by its location.



These arrows all represent the vector  $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$ .

However, unless otherwise specified, we'll assume a vector starts at the origin: we'll usually be sloppy and identify the vector  $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$  with the point (1, 2).

This makes sense in the real world: many physical quantities, such as velocity, are represented as vectors. But it makes more sense to think of the velocity of a car as being located at the car.

Another way to think about it: a vector is a *difference* between two points, or the arrow from one point to another.

# Points and Vectors

So what is the difference between a point and a vector?

A vector need not start at the origin: *it can be located anywhere!* In other words, an arrow is determined by its length and its direction, not by its location.




These arrows all represent the vector  $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$ .

However, unless otherwise specified, we'll assume a vector starts at the origin: we'll usually be sloppy and identify the vector  $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$  with the point  $(1, 2)$ .

This makes sense in the real world: many physical quantities, such as velocity, are represented as vectors. But it makes more sense to think of the velocity of a car as being located at the car.

Another way to think about it: a vector is a *difference* between two points, or the arrow from one point to another.

For instance,  $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$  is the arrow from  $(1, 1)$  to  $(2, 3)$ .



## Definition

- ▶ We can add two vectors together:

$$\begin{pmatrix} a \\ b \\ c \end{pmatrix} + \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} a+x \\ b+y \\ c+z \end{pmatrix} .$$

## Definition

- ▶ We can add two vectors together:

$$\begin{pmatrix} a \\ b \\ c \end{pmatrix} + \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} a+x \\ b+y \\ c+z \end{pmatrix} .$$

- ▶ We can multiply, or **scale**, a vector by a real number  $c$ :

$$c \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} c \cdot x \\ c \cdot y \\ c \cdot z \end{pmatrix} .$$

## Definition

- ▶ We can add two vectors together:

$$\begin{pmatrix} a \\ b \\ c \end{pmatrix} + \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} a+x \\ b+y \\ c+z \end{pmatrix}.$$

- ▶ We can multiply, or **scale**, a vector by a real number  $c$ :

$$c \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} c \cdot x \\ c \cdot y \\ c \cdot z \end{pmatrix}.$$

We call  $c$  a **scalar** to distinguish it from a vector. If  $v$  is a vector and  $c$  is a scalar,  $cv$  is called a **scalar multiple** of  $v$ .

## Definition

- ▶ We can add two vectors together:

$$\begin{pmatrix} a \\ b \\ c \end{pmatrix} + \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} a + x \\ b + y \\ c + z \end{pmatrix}.$$

- ▶ We can multiply, or **scale**, a vector by a real number  $c$ :

$$c \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} c \cdot x \\ c \cdot y \\ c \cdot z \end{pmatrix}.$$

We call  $c$  a **scalar** to distinguish it from a vector. If  $v$  is a vector and  $c$  is a scalar,  $cv$  is called a **scalar multiple** of  $v$ .

(And likewise for vectors of length  $n$ .)

## Definition

- ▶ We can add two vectors together:

$$\begin{pmatrix} a \\ b \\ c \end{pmatrix} + \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} a+x \\ b+y \\ c+z \end{pmatrix}.$$

- ▶ We can multiply, or **scale**, a vector by a real number  $c$ :


$$c \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} c \cdot x \\ c \cdot y \\ c \cdot z \end{pmatrix}.$$

We call  $c$  a **scalar** to distinguish it from a vector. If  $v$  is a vector and  $c$  is a scalar,  $cv$  is called a **scalar multiple** of  $v$ .

(And likewise for vectors of length  $n$ .) For instance,


## Vector Addition and Subtraction: Geometry

The parallelogram law for vector addition






## Vector Addition and Subtraction: Geometry



### The parallelogram law for vector addition

Geometrically, the sum of two vectors  $v, w$  is obtained as follows:


## Vector Addition and Subtraction: Geometry



### The parallelogram law for vector addition

Geometrically, the sum of two vectors  $v, w$  is obtained as follows: place the tail of  $w$  at the head of  $v$ .


## Vector Addition and Subtraction: Geometry



### The parallelogram law for vector addition

Geometrically, the sum of two vectors  $v, w$  is obtained as follows: place the tail of  $w$  at the head of  $v$ . Then  $v + w$  is the vector whose tail is the tail of  $v$  and whose head is the head of  $w$ .


# Vector Addition and Subtraction: Geometry



## The parallelogram law for vector addition

Geometrically, the sum of two vectors  $v, w$  is obtained as follows: place the tail of  $w$  at the head of  $v$ . Then  $v + w$  is the vector whose tail is the tail of  $v$  and whose head is the head of  $w$ . Doing this both ways creates a **parallelogram**.

## Vector Addition and Subtraction: Geometry




### The parallelogram law for vector addition

Geometrically, the sum of two vectors  $v, w$  is obtained as follows: place the tail of  $w$  at the head of  $v$ . Then  $v + w$  is the vector whose tail is the tail of  $v$  and whose head is the head of  $w$ . Doing this both ways creates a **parallelogram**. For example,

$$\begin{pmatrix} 1 \\ 3 \end{pmatrix} + \begin{pmatrix} 4 \\ 2 \end{pmatrix} = \begin{pmatrix} 5 \\ 5 \end{pmatrix}.$$

# Vector Addition and Subtraction: Geometry




## The parallelogram law for vector addition

Geometrically, the sum of two vectors  $v, w$  is obtained as follows: place the tail of  $w$  at the head of  $v$ . Then  $v + w$  is the vector whose tail is the tail of  $v$  and whose head is the head of  $w$ . Doing this both ways creates a **parallelogram**. For example,

$$\begin{pmatrix} 1 \\ 3 \end{pmatrix} + \begin{pmatrix} 4 \\ 2 \end{pmatrix} = \begin{pmatrix} 5 \\ 5 \end{pmatrix}.$$

Why? The width of  $v + w$  is the sum of the widths, and likewise with the heights.

# Vector Addition and Subtraction: Geometry




## The parallelogram law for vector addition

Geometrically, the sum of two vectors  $v, w$  is obtained as follows: place the tail of  $w$  at the head of  $v$ . Then  $v + w$  is the vector whose tail is the tail of  $v$  and whose head is the head of  $w$ . Doing this both ways creates a **parallelogram**. For example,


$$\begin{pmatrix} 1 \\ 3 \end{pmatrix} + \begin{pmatrix} 4 \\ 2 \end{pmatrix} = \begin{pmatrix} 5 \\ 5 \end{pmatrix}.$$

Why? The width of  $v + w$  is the sum of the widths, and likewise with the heights.

## Vector subtraction



# Vector Addition and Subtraction: Geometry



## The parallelogram law for vector addition


Geometrically, the sum of two vectors  $v, w$  is obtained as follows: place the tail of  $w$  at the head of  $v$ . Then  $v + w$  is the vector whose tail is the tail of  $v$  and whose head is the head of  $w$ . Doing this both ways creates a **parallelogram**. For example,

$$\begin{pmatrix} 1 \\ 3 \end{pmatrix} + \begin{pmatrix} 4 \\ 2 \end{pmatrix} = \begin{pmatrix} 5 \\ 5 \end{pmatrix}.$$

Why? The width of  $v + w$  is the sum of the widths, and likewise with the heights.


## Vector subtraction

Geometrically, the difference of two vectors  $v, w$  is obtained as follows:





# Vector Addition and Subtraction: Geometry



## The parallelogram law for vector addition


Geometrically, the sum of two vectors  $v, w$  is obtained as follows: place the tail of  $w$  at the head of  $v$ . Then  $v + w$  is the vector whose tail is the tail of  $v$  and whose head is the head of  $w$ . Doing this both ways creates a **parallelogram**. For example,

$$\begin{pmatrix} 1 \\ 3 \end{pmatrix} + \begin{pmatrix} 4 \\ 2 \end{pmatrix} = \begin{pmatrix} 5 \\ 5 \end{pmatrix}.$$


Why? The width of  $v + w$  is the sum of the widths, and likewise with the heights.

## Vector subtraction

Geometrically, the difference of two vectors  $v, w$  is obtained as follows: place the tail of  $v$  and  $w$  at the same point.



# Vector Addition and Subtraction: Geometry



## The parallelogram law for vector addition


Geometrically, the sum of two vectors  $v, w$  is obtained as follows: place the tail of  $w$  at the head of  $v$ . Then  $v + w$  is the vector whose tail is the tail of  $v$  and whose head is the head of  $w$ . Doing this both ways creates a **parallelogram**. For example,

$$\begin{pmatrix} 1 \\ 3 \end{pmatrix} + \begin{pmatrix} 4 \\ 2 \end{pmatrix} = \begin{pmatrix} 5 \\ 5 \end{pmatrix}.$$


Why? The width of  $v + w$  is the sum of the widths, and likewise with the heights.

## Vector subtraction

Geometrically, the difference of two vectors  $v, w$  is obtained as follows: place the tail of  $v$  and  $w$  at the same point. Then  $v - w$  is the vector from the head of  $v$  to the head of  $w$ .



# Vector Addition and Subtraction: Geometry



## The parallelogram law for vector addition

Geometrically, the sum of two vectors  $v, w$  is obtained as follows: place the tail of  $w$  at the head of  $v$ . Then  $v + w$  is the vector whose tail is the tail of  $v$  and whose head is the head of  $w$ . Doing this both ways creates a **parallelogram**. For example,


$$\begin{pmatrix} 1 \\ 3 \end{pmatrix} + \begin{pmatrix} 4 \\ 2 \end{pmatrix} = \begin{pmatrix} 5 \\ 5 \end{pmatrix}.$$

Why? The width of  $v + w$  is the sum of the widths, and likewise with the heights.


## Vector subtraction

Geometrically, the difference of two vectors  $v, w$  is obtained as follows: place the tail of  $v$  and  $w$  at the same point. Then  $v - w$  is the vector from the head of  $v$  to the head of  $w$ . For example,

$$\begin{pmatrix} 1 \\ 4 \end{pmatrix} - \begin{pmatrix} 4 \\ 2 \end{pmatrix} = \begin{pmatrix} -3 \\ 2 \end{pmatrix}.$$



# Vector Addition and Subtraction: Geometry



## The parallelogram law for vector addition

Geometrically, the sum of two vectors  $v, w$  is obtained as follows: place the tail of  $w$  at the head of  $v$ . Then  $v + w$  is the vector whose tail is the tail of  $v$  and whose head is the head of  $w$ . Doing this both ways creates a **parallelogram**. For example,

$$\begin{pmatrix} 1 \\ 3 \end{pmatrix} + \begin{pmatrix} 4 \\ 2 \end{pmatrix} = \begin{pmatrix} 5 \\ 5 \end{pmatrix}.$$


Why? The width of  $v + w$  is the sum of the widths, and likewise with the heights.

## Vector subtraction


Geometrically, the difference of two vectors  $v, w$  is obtained as follows: place the tail of  $v$  and  $w$  at the same point. Then  $v - w$  is the vector from the head of  $v$  to the head of  $w$ . For example,

$$\begin{pmatrix} 1 \\ 4 \end{pmatrix} - \begin{pmatrix} 4 \\ 2 \end{pmatrix} = \begin{pmatrix} -3 \\ 2 \end{pmatrix}.$$

Why? If you add  $v - w$  to  $w$ , you get  $v$ .



# Vector Addition and Subtraction: Geometry



## The parallelogram law for vector addition

Geometrically, the sum of two vectors  $v, w$  is obtained as follows: place the tail of  $w$  at the head of  $v$ . Then  $v + w$  is the vector whose tail is the tail of  $v$  and whose head is the head of  $w$ . Doing this both ways creates a **parallelogram**. For example,

$$\begin{pmatrix} 1 \\ 3 \end{pmatrix} + \begin{pmatrix} 4 \\ 2 \end{pmatrix} = \begin{pmatrix} 5 \\ 5 \end{pmatrix}.$$

Why? The width of  $v + w$  is the sum of the widths, and likewise with the heights.


## Vector subtraction

Geometrically, the difference of two vectors  $v, w$  is obtained as follows: place the tail of  $v$  and  $w$  at the same point. Then  $v - w$  is the vector from the head of  $v$  to the head of  $w$ . For example,

$$\begin{pmatrix} 1 \\ 4 \end{pmatrix} - \begin{pmatrix} 4 \\ 2 \end{pmatrix} = \begin{pmatrix} -3 \\ 2 \end{pmatrix}.$$

Why? If you add  $v - w$  to  $w$ , you get  $v$ .

This works in higher dimensions too!




# Scalar Multiplication: Geometry

## Scalar multiples of a vector

These have the same *direction* but a different *length*.

Some multiples of  $v$ .



$$v = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$$2v = \begin{pmatrix} 2 \\ 4 \end{pmatrix}$$

$$-\frac{1}{2}v = \begin{pmatrix} -\frac{1}{2} \\ -1 \end{pmatrix}$$

$$0v = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

# Scalar Multiplication: Geometry

## Scalar multiples of a vector

These have the same *direction* but a different *length*.

Some multiples of  $v$ .




$$v = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$$2v = \begin{pmatrix} 2 \\ 4 \end{pmatrix}$$

$$-\frac{1}{2}v = \begin{pmatrix} -\frac{1}{2} \\ -1 \end{pmatrix}$$

$$0v = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

All multiples of  $v$ .




# Scalar Multiplication: Geometry

## Scalar multiples of a vector

These have the same *direction* but a different *length*.

Some multiples of  $v$ .




$$v = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$$2v = \begin{pmatrix} 2 \\ 4 \end{pmatrix}$$

$$-\frac{1}{2}v = \begin{pmatrix} -\frac{1}{2} \\ -1 \end{pmatrix}$$

$$0v = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

All multiples of  $v$ .






# Scalar Multiplication: Geometry

## Scalar multiples of a vector

These have the same *direction* but a different *length*.

Some multiples of  $v$ .




$$v = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$$2v = \begin{pmatrix} 2 \\ 4 \end{pmatrix}$$

$$-\frac{1}{2}v = \begin{pmatrix} -\frac{1}{2} \\ -1 \end{pmatrix}$$

$$0v = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

All multiples of  $v$ .



So the scalar multiples of  $v$  form a *line*.

## Linear Combinations

We can add and scalar multiply in the same equation:

# Linear Combinations

We can add and scalar multiply in the same equation:

$$w = c_1 v_1 + c_2 v_2 + \cdots + c_p v_p$$

where  $c_1, c_2, \dots, c_p$  are \_\_\_\_\_,

# Linear Combinations

We can add and scalar multiply in the same equation:

$$w = c_1 v_1 + c_2 v_2 + \cdots + c_p v_p$$

where  $c_1, c_2, \dots, c_p$  are scalars,  $v_1, v_2, \dots, v_p$  are \_\_\_\_\_

# Linear Combinations

We can add and scalar multiply in the same equation:

$$w = c_1 v_1 + c_2 v_2 + \cdots + c_p v_p$$

where  $c_1, c_2, \dots, c_p$  are scalars,  $v_1, v_2, \dots, v_p$  are vectors in  $\mathbf{R}^n$ , and  $w$  is a

---

# Linear Combinations

We can add and scalar multiply in the same equation:

$$w = c_1 v_1 + c_2 v_2 + \cdots + c_p v_p$$

where  $c_1, c_2, \dots, c_p$  are scalars,  $v_1, v_2, \dots, v_p$  are vectors in  $\mathbf{R}^n$ , and  $w$  is a vector in  $\mathbf{R}^n$ .

# Linear Combinations

We can add and scalar multiply in the same equation:

$$w = c_1 v_1 + c_2 v_2 + \cdots + c_p v_p$$

where  $c_1, c_2, \dots, c_p$  are scalars,  $v_1, v_2, \dots, v_p$  are vectors in  $\mathbf{R}^n$ , and  $w$  is a vector in  $\mathbf{R}^n$ .

## Definition

We call  $w$  a **linear combination** of the vectors  $v_1, v_2, \dots, v_p$ .

# Linear Combinations

We can add and scalar multiply in the same equation:

$$w = c_1 v_1 + c_2 v_2 + \cdots + c_p v_p$$

where  $c_1, c_2, \dots, c_p$  are scalars,  $v_1, v_2, \dots, v_p$  are vectors in  $\mathbf{R}^n$ , and  $w$  is a vector in  $\mathbf{R}^n$ .

## Definition

We call  $w$  a **linear combination** of the vectors  $v_1, v_2, \dots, v_p$ . The scalars  $c_1, c_2, \dots, c_p$  are called the **weights** or **coefficients**.



# Linear Combinations

We can add and scalar multiply in the same equation:


$$w = c_1 v_1 + c_2 v_2 + \cdots + c_p v_p$$

where  $c_1, c_2, \dots, c_p$  are scalars,  $v_1, v_2, \dots, v_p$  are vectors in  $\mathbf{R}^n$ , and  $w$  is a vector in  $\mathbf{R}^n$ .

## Definition

We call  $w$  a **linear combination** of the vectors  $v_1, v_2, \dots, v_p$ . The scalars  $c_1, c_2, \dots, c_p$  are called the **weights** or **coefficients**.

## Example




Let  $v = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$  and  $w = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ .

What are some linear combinations of  $v$  and  $w$ ?

- ▶  $v + w$
- ▶  $v - w$
- ▶  $2v + 0w$
- ▶  $2w$
- ▶  $-v$

Poll


Is there any vector in  $\mathbf{R}^2$  that is *not* a linear combination of  $v$  and  $w$ ?



Poll

Is there any vector in  $\mathbf{R}^2$  that is *not* a linear combination of  $v$  and  $w$ ?


No: in fact, every vector in  $\mathbf{R}^2$  is a combination of  $v$  and  $w$ .



Poll


Is there any vector in  $\mathbf{R}^2$  that is *not* a linear combination of  $v$  and  $w$ ?

No: in fact, *every* vector in  $\mathbf{R}^2$  is a combination of  $v$  and  $w$ .




(The purple lines are to help measure *how much* of  $v$  and  $w$  you need to get to a given point.)

## More Examples




What are some linear combinations of  $v = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$ ?

## More Examples



What are some linear combinations of  $v = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$ ?




### Question


What are all linear combinations of

$$v = \begin{pmatrix} 2 \\ 2 \end{pmatrix} \quad \text{and} \quad w = \begin{pmatrix} -1 \\ -1 \end{pmatrix}?$$

## More Examples



What are some linear combinations of  $v = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$ ?




### Question

What are all linear combinations of


$$v = \begin{pmatrix} 2 \\ 2 \end{pmatrix} \quad \text{and} \quad w = \begin{pmatrix} -1 \\ -1 \end{pmatrix}?$$

**Answer:** The line which contains both vectors.

## More Examples



What are some linear combinations of  $v = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$ ?



### Question

What are all linear combinations of

$$v = \begin{pmatrix} 2 \\ 2 \end{pmatrix} \quad \text{and} \quad w = \begin{pmatrix} -1 \\ -1 \end{pmatrix}?$$

**Answer:** The line which contains both vectors.

What's different about this example and the one on the poll?



## Systems of Linear Equations

### Question

Is  $\begin{pmatrix} 8 \\ 16 \\ 3 \end{pmatrix}$  a linear combination of  $\begin{pmatrix} 1 \\ 2 \\ 6 \end{pmatrix}$  and  $\begin{pmatrix} -1 \\ -2 \\ -1 \end{pmatrix}$ ?

# Systems of Linear Equations

Continued

$$x - y = 8$$

$$2x - 2y = 16$$

$$6x - y = 3$$

# Systems of Linear Equations

Continued

$$x - y = 8$$

$$2x - 2y = 16$$

$$6x - y = 3$$

matrix form  


$$\left( \begin{array}{cc|c} 1 & -1 & 8 \\ 2 & -2 & 16 \\ 6 & -1 & 3 \end{array} \right)$$

# Systems of Linear Equations

Continued

$$x - y = 8$$

$$2x - 2y = 16$$

$$6x - y = 3$$

matrix form  
~~~~~>

$$\left( \begin{array}{cc|c} 1 & -1 & 8 \\ 2 & -2 & 16 \\ 6 & -1 & 3 \end{array} \right)$$

row reduce  
~~~~~>

$$\left( \begin{array}{cc|c} 1 & 0 & -1 \\ 0 & 1 & -9 \\ 0 & 0 & 0 \end{array} \right)$$

# Systems of Linear Equations


Continued

$$x - y = 8$$

$$2x - 2y = 16$$


$$6x - y = 3$$

matrix form



$$\left( \begin{array}{cc|c} 1 & -1 & 8 \\ 2 & -2 & 16 \\ 6 & -1 & 3 \end{array} \right)$$

row reduce



$$\left( \begin{array}{cc|c} 1 & 0 & -1 \\ 0 & 1 & -9 \\ 0 & 0 & 0 \end{array} \right)$$

solution



$$x = -1$$

$$y = -9$$

# Systems of Linear Equations

Continued

$$x - y = 8$$

$$2x - 2y = 16$$

$$6x - y = 3$$

matrix form  
~~~~~>

$$\left( \begin{array}{cc|c} 1 & -1 & 8 \\ 2 & -2 & 16 \\ 6 & -1 & 3 \end{array} \right)$$

row reduce  
~~~~~>

$$\left( \begin{array}{cc|c} 1 & 0 & -1 \\ 0 & 1 & -9 \\ 0 & 0 & 0 \end{array} \right)$$

solution  
~~~~~>

$$x = -1$$

$$y = -9$$

Conclusion:

$$-\begin{pmatrix} 1 \\ 2 \\ 6 \end{pmatrix} - 9 \begin{pmatrix} -1 \\ -2 \\ -1 \end{pmatrix} = \begin{pmatrix} 8 \\ 16 \\ 3 \end{pmatrix}$$

# Systems of Linear Equations

Continued

$$x - y = 8$$

$$2x - 2y = 16$$

$$6x - y = 3$$

matrix form  
~~~~~>

$$\left( \begin{array}{cc|c} 1 & -1 & 8 \\ 2 & -2 & 16 \\ 6 & -1 & 3 \end{array} \right)$$

row reduce  
~~~~~>

$$\left( \begin{array}{cc|c} 1 & 0 & -1 \\ 0 & 1 & -9 \\ 0 & 0 & 0 \end{array} \right)$$

solution  
~~~~~>

$$x = -1$$

$$y = -9$$

Conclusion:

$$-\begin{pmatrix} 1 \\ 2 \\ 6 \end{pmatrix} - 9 \begin{pmatrix} -1 \\ -2 \\ -1 \end{pmatrix} = \begin{pmatrix} 8 \\ 16 \\ 3 \end{pmatrix}$$

What is the relationship between the original vectors and the matrix form of the linear equation?

# Systems of Linear Equations

Continued

$$x - y = 8$$

$$2x - 2y = 16$$

$$6x - y = 3$$

matrix form  
~~~~~→

$$\left( \begin{array}{cc|c} 1 & -1 & 8 \\ 2 & -2 & 16 \\ 6 & -1 & 3 \end{array} \right)$$

row reduce  
~~~~~→

$$\left( \begin{array}{cc|c} 1 & 0 & -1 \\ 0 & 1 & -9 \\ 0 & 0 & 0 \end{array} \right)$$

solution  
~~~~~→

$$x = -1$$

$$y = -9$$

Conclusion:

$$-\begin{pmatrix} 1 \\ 2 \\ 6 \end{pmatrix} - 9 \begin{pmatrix} -1 \\ -2 \\ -1 \end{pmatrix} = \begin{pmatrix} 8 \\ 16 \\ 3 \end{pmatrix}$$

What is the relationship between the original vectors and the matrix form of the linear equation? They have the same columns!



# Systems of Linear Equations

Continued

$$x - y = 8$$

$$2x - 2y = 16$$

$$6x - y = 3$$

matrix form  
~~~~~>

$$\left( \begin{array}{cc|c} 1 & -1 & 8 \\ 2 & -2 & 16 \\ 6 & -1 & 3 \end{array} \right)$$

row reduce  
~~~~~>

$$\left( \begin{array}{cc|c} 1 & 0 & -1 \\ 0 & 1 & -9 \\ 0 & 0 & 0 \end{array} \right)$$

solution  
~~~~~>

$$x = -1$$

$$y = -9$$

Conclusion:

$$-\begin{pmatrix} 1 \\ 2 \\ 6 \end{pmatrix} - 9 \begin{pmatrix} -1 \\ -2 \\ -1 \end{pmatrix} = \begin{pmatrix} 8 \\ 16 \\ 3 \end{pmatrix}$$

What is the relationship between the original vectors and the matrix form of the linear equation? They have the same columns!

**Shortcut:** You can make the augmented matrix without writing down the system of linear equations first.

# Vector Equations and Linear Equations

## Summary

The vector equation

$$x_1 v_1 + x_2 v_2 + \cdots + x_p v_p = b,$$

where  $v_1, v_2, \dots, v_p, b$  are vectors in  $\mathbf{R}^n$  and  $x_1, x_2, \dots, x_p$  are scalars,

# Vector Equations and Linear Equations

## Summary

The vector equation

$$x_1 v_1 + x_2 v_2 + \cdots + x_p v_p = b,$$

where  $v_1, v_2, \dots, v_p, b$  are vectors in  $\mathbf{R}^n$  and  $x_1, x_2, \dots, x_p$  are scalars, has the same solution set as the linear system with augmented matrix

$$\left( \begin{array}{ccc|c} | & | & & | \\ v_1 & v_2 & \cdots & v_p \\ | & | & & | \end{array} \middle| \begin{array}{c} | \\ b \\ | \end{array} \right),$$

where the  $v_i$ 's and  $b$  are the columns of the matrix.

# Vector Equations and Linear Equations

## Summary

The vector equation

$$x_1 v_1 + x_2 v_2 + \cdots + x_p v_p = b,$$

where  $v_1, v_2, \dots, v_p, b$  are vectors in  $\mathbf{R}^n$  and  $x_1, x_2, \dots, x_p$  are scalars, has the same solution set as the linear system with augmented matrix

$$\left( \begin{array}{c|c|c|c|c} | & | & & | & | \\ v_1 & v_2 & \cdots & v_p & b \\ | & | & & | & | \end{array} \right),$$

where the  $v_i$ 's and  $b$  are the columns of the matrix.

So we now have (at least) *two* equivalent ways of thinking about linear systems of equations:

# Vector Equations and Linear Equations

## Summary

The vector equation

$$x_1 v_1 + x_2 v_2 + \cdots + x_p v_p = b,$$

where  $v_1, v_2, \dots, v_p, b$  are vectors in  $\mathbf{R}^n$  and  $x_1, x_2, \dots, x_p$  are scalars, has the same solution set as the linear system with augmented matrix

$$\left( \begin{array}{c|c|c|c|c} | & | & & | & | \\ v_1 & v_2 & \cdots & v_p & b \\ | & | & & | & | \end{array} \right),$$

where the  $v_i$ 's and  $b$  are the columns of the matrix.

So we now have (at least) *two* equivalent ways of thinking about linear systems of equations:

1. Augmented matrices.

# Vector Equations and Linear Equations

## Summary

The vector equation

$$x_1 v_1 + x_2 v_2 + \cdots + x_p v_p = b,$$

where  $v_1, v_2, \dots, v_p, b$  are vectors in  $\mathbf{R}^n$  and  $x_1, x_2, \dots, x_p$  are scalars, has the same solution set as the linear system with augmented matrix

$$\left( \begin{array}{c|c|c|c|c} | & | & & | & | \\ v_1 & v_2 & \cdots & v_p & b \\ | & | & & | & | \end{array} \right),$$

where the  $v_i$ 's and  $b$  are the columns of the matrix.

So we now have (at least) *two* equivalent ways of thinking about linear systems of equations:

1. Augmented matrices.
2. Linear combinations of vectors (vector equations).

# Vector Equations and Linear Equations

## Summary

The vector equation

$$x_1 v_1 + x_2 v_2 + \cdots + x_p v_p = b,$$

where  $v_1, v_2, \dots, v_p, b$  are vectors in  $\mathbf{R}^n$  and  $x_1, x_2, \dots, x_p$  are scalars, has the same solution set as the linear system with augmented matrix

$$\left( \begin{array}{c|c|c|c|c} | & | & & | & | \\ v_1 & v_2 & \cdots & v_p & b \\ | & | & & | & | \end{array} \right),$$

where the  $v_i$ 's and  $b$  are the columns of the matrix.

So we now have (at least) *two* equivalent ways of thinking about linear systems of equations:

1. Augmented matrices.
2. Linear combinations of vectors (vector equations).

The last one is more geometric in nature.

# Span

It is important to know what are *all* linear combinations of a set of vectors  $v_1, v_2, \dots, v_p$  in  $\mathbf{R}^n$ :



# Span

It is important to know what are *all* linear combinations of a set of vectors  $v_1, v_2, \dots, v_p$  in  $\mathbf{R}^n$ : it's exactly the collection of all  $b$  in  $\mathbf{R}^n$  such that the vector equation (in the unknowns  $x_1, x_2, \dots, x_p$ )

$$x_1 v_1 + x_2 v_2 + \cdots + x_p v_p = b$$

has a solution (i.e., is consistent).

# Span

It is important to know what are *all* linear combinations of a set of vectors  $v_1, v_2, \dots, v_p$  in  $\mathbf{R}^n$ : it's exactly the collection of all  $b$  in  $\mathbf{R}^n$  such that the vector equation (in the unknowns  $x_1, x_2, \dots, x_p$ )

$$x_1 v_1 + x_2 v_2 + \dots + x_p v_p = b$$

has a solution (i.e., is consistent).

## Definition

Let  $v_1, v_2, \dots, v_p$  be vectors in  $\mathbf{R}^n$ . The **span** of  $v_1, v_2, \dots, v_p$  is the collection of all linear combinations of  $v_1, v_2, \dots, v_p$ , and is denoted  $\text{Span}\{v_1, v_2, \dots, v_p\}$ .

# Span

It is important to know what are *all* linear combinations of a set of vectors  $v_1, v_2, \dots, v_p$  in  $\mathbf{R}^n$ : it's exactly the collection of all  $b$  in  $\mathbf{R}^n$  such that the vector equation (in the unknowns  $x_1, x_2, \dots, x_p$ )

$$x_1 v_1 + x_2 v_2 + \dots + x_p v_p = b$$

has a solution (i.e., is consistent).

## Definition

Let  $v_1, v_2, \dots, v_p$  be vectors in  $\mathbf{R}^n$ . The **span** of  $v_1, v_2, \dots, v_p$  is the collection of all linear combinations of  $v_1, v_2, \dots, v_p$ , and is denoted  $\text{Span}\{v_1, v_2, \dots, v_p\}$ . In symbols:

$$\text{Span}\{v_1, v_2, \dots, v_p\} = \{ x_1 v_1 + x_2 v_2 + \dots + x_p v_p \mid x_1, x_2, \dots, x_p \text{ in } \mathbf{R} \}.$$

# Span


It is important to know what are *all* linear combinations of a set of vectors  $v_1, v_2, \dots, v_p$  in  $\mathbf{R}^n$ : it's exactly the collection of all  $b$  in  $\mathbf{R}^n$  such that the vector equation (in the unknowns  $x_1, x_2, \dots, x_p$ )

$$x_1 v_1 + x_2 v_2 + \dots + x_p v_p = b$$

has a solution (i.e., is consistent).

## Definition

Let  $v_1, v_2, \dots, v_p$  be vectors in  $\mathbf{R}^n$ . The **span** of  $v_1, v_2, \dots, v_p$  is the collection of all linear combinations of  $v_1, v_2, \dots, v_p$ , and is denoted  $\text{Span}\{v_1, v_2, \dots, v_p\}$ . In symbols:


$$\text{Span}\{v_1, v_2, \dots, v_p\} = \left\{ x_1 v_1 + x_2 v_2 + \dots + x_p v_p \mid x_1, x_2, \dots, x_p \text{ in } \mathbf{R} \right\}.$$

# Span


It is important to know what are *all* linear combinations of a set of vectors  $v_1, v_2, \dots, v_p$  in  $\mathbf{R}^n$ : it's exactly the collection of all  $b$  in  $\mathbf{R}^n$  such that the vector equation (in the unknowns  $x_1, x_2, \dots, x_p$ )

$$x_1 v_1 + x_2 v_2 + \dots + x_p v_p = b$$

has a solution (i.e., is consistent).

## Definition

Let  $v_1, v_2, \dots, v_p$  be vectors in  $\mathbf{R}^n$ . The **span** of  $v_1, v_2, \dots, v_p$  is the collection of all linear combinations of  $v_1, v_2, \dots, v_p$ , and is denoted  $\text{Span}\{v_1, v_2, \dots, v_p\}$ . In symbols:


$$\text{Span}\{v_1, v_2, \dots, v_p\} = \left\{ x_1 v_1 + x_2 v_2 + \dots + x_p v_p \mid x_1, x_2, \dots, x_p \text{ in } \mathbf{R} \right\}.$$

**Synonyms:**  $\text{Span}\{v_1, v_2, \dots, v_p\}$  is the subset **spanned by** or **generated by**  $v_1, v_2, \dots, v_p$ .

# Span


It is important to know what are *all* linear combinations of a set of vectors  $v_1, v_2, \dots, v_p$  in  $\mathbf{R}^n$ : it's exactly the collection of all  $b$  in  $\mathbf{R}^n$  such that the vector equation (in the unknowns  $x_1, x_2, \dots, x_p$ )

$$x_1 v_1 + x_2 v_2 + \dots + x_p v_p = b$$

has a solution (i.e., is consistent).

## Definition

Let  $v_1, v_2, \dots, v_p$  be vectors in  $\mathbf{R}^n$ . The **span** of  $v_1, v_2, \dots, v_p$  is the collection of all linear combinations of  $v_1, v_2, \dots, v_p$ , and is denoted  $\text{Span}\{v_1, v_2, \dots, v_p\}$ . In symbols:


$$\text{Span}\{v_1, v_2, \dots, v_p\} = \{ x_1 v_1 + x_2 v_2 + \dots + x_p v_p \mid x_1, x_2, \dots, x_p \text{ in } \mathbf{R} \}.$$

**Synonyms:**  $\text{Span}\{v_1, v_2, \dots, v_p\}$  is the subset **spanned by** or **generated by**  $v_1, v_2, \dots, v_p$ .

This is the first of several definitions in this class that you simply **must learn**. I will give you other ways to think about Span, and ways to draw pictures, but *this is the definition*. Having a vague idea what Span means will not help you solve any exam problems!

# Span

Continued

Now we have several equivalent ways of making the same statement:

# Span

## Continued

Now we have several equivalent ways of making the same statement:

1. A vector  $b$  is in the span of  $v_1, v_2, \dots, v_p$ .



# Span

## Continued

Now we have several equivalent ways of making the same statement:

1. A vector  $b$  is in the span of  $v_1, v_2, \dots, v_p$ .
2. The linear system with augmented matrix

$$\left( \begin{array}{c|c|c|c|c} | & | & & | & | \\ v_1 & v_2 & \cdots & v_p & b \\ | & | & & | & | \end{array} \right)$$

is consistent.

# Span

## Continued

Now we have several equivalent ways of making the same statement:

1. A vector  $b$  is in the span of  $v_1, v_2, \dots, v_p$ .
2. The linear system with augmented matrix

$$\left( \begin{array}{c|c|c|c|c} | & | & & | & \\ \hline v_1 & v_2 & \cdots & v_p & b \\ \hline | & | & & | & \end{array} \right)$$

is consistent.

3. The vector equation

$$x_1 v_1 + x_2 v_2 + \cdots + x_p v_p = b$$

has a solution.

# Span

## Continued

Now we have several equivalent ways of making the same statement:

1. A vector  $b$  is in the span of  $v_1, v_2, \dots, v_p$ .
2. The linear system with augmented matrix

$$\left( \begin{array}{c|c|c|c|c} | & | & & | & \\ \hline v_1 & v_2 & \cdots & v_p & b \\ \hline | & | & & | & \end{array} \right)$$

is consistent.

3. The vector equation

$$x_1 v_1 + x_2 v_2 + \cdots + x_p v_p = b$$

has a solution.


**Note:** **equivalent** means that, for any given list of vectors  $v_1, v_2, \dots, v_p, b$ , *either* all three statements are true, *or* all three statements are false.

## Pictures of Span

Drawing a picture of  $\text{Span}\{v_1, v_2, \dots, v_p\}$  is the same as drawing a picture of all linear combinations of  $v_1, v_2, \dots, v_p$ .


## Pictures of Span

Drawing a picture of  $\text{Span}\{v_1, v_2, \dots, v_p\}$  is the same as drawing a picture of all linear combinations of  $v_1, v_2, \dots, v_p$ .





## Pictures of Span

Drawing a picture of  $\text{Span}\{v_1, v_2, \dots, v_p\}$  is the same as drawing a picture of all linear combinations of  $v_1, v_2, \dots, v_p$ .





# Pictures of Span

Drawing a picture of  $\text{Span}\{v_1, v_2, \dots, v_p\}$  is the same as drawing a picture of all linear combinations of  $v_1, v_2, \dots, v_p$ .



# Pictures of Span




Drawing a picture of  $\text{Span}\{v_1, v_2, \dots, v_p\}$  is the same as drawing a picture of all linear combinations of  $v_1, v_2, \dots, v_p$ .








# Pictures of Span

Drawing a picture of  $\text{Span}\{v_1, v_2, \dots, v_p\}$  is the same as drawing a picture of all linear combinations of  $v_1, v_2, \dots, v_p$ .




# Pictures of Span

Drawing a picture of  $\text{Span}\{v_1, v_2, \dots, v_p\}$  is the same as drawing a picture of all linear combinations of  $v_1, v_2, \dots, v_p$ .




# Pictures of Span

In  $\mathbb{R}^3$





# Pictures of Span

In  $\mathbb{R}^3$





# Pictures of Span

In  $\mathbb{R}^3$






# Pictures of Span

In  $\mathbb{R}^3$






# Pictures of Span

In  $\mathbb{R}^3$



# Pictures of Span





In  $\mathbb{R}^3$









# Pictures of Span

In  $\mathbb{R}^3$



# Pictures of Span

In  $\mathbb{R}^3$



## Poll

How many vectors are in  $\text{Span} \left\{ \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \right\}$ ?

- A. Zero
- B. One
- C. Infinity

Poll

How many vectors are in  $\text{Span} \left\{ \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \right\}$ ?

- A. Zero
- B. One
- C. Infinity

In general, it appears that  $\text{Span}\{v_1, v_2, \dots, v_p\}$  is the smallest “linear space” (line, plane, etc.) containing the origin and all of the vectors  $v_1, v_2, \dots, v_p$ .

Poll

How many vectors are in  $\text{Span} \left\{ \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \right\}$ ?

- A. Zero
- B. One
- C. Infinity

In general, it appears that  $\text{Span}\{v_1, v_2, \dots, v_p\}$  is the smallest “linear space” (line, plane, etc.) containing the origin and all of the vectors  $v_1, v_2, \dots, v_p$ .

We will make this precise later.