

# Chapter 1

## Linear Equations

# Section 1.1

## Systems of Linear Equations

# One Linear Equation

What does the solution set of a linear equation look like?

# One Linear Equation

What does the solution set of a linear equation look like?


- ▶  $x + y = 1$

# One Linear Equation

What does the solution set of a linear equation look like?

►  $x + y = 1$

~~~~~→ a line in the plane:  $y = 1 - x$




# One Linear Equation

What does the solution set of a linear equation look like?

►  $x + y = 1$

~~~~~→ a line in the plane:  $y = 1 - x$




►  $x + y + z = 1$

# One Linear Equation

What does the solution set of a linear equation look like?


►  $x + y = 1$

~~~~~> a line in the plane:  $y = 1 - x$



►  $x + y + z = 1$

~~~~~> a plane in space:  $z = 1 - x - y$




# One Linear Equation

What does the solution set of a linear equation look like?


►  $x + y = 1$

~~~~~→ a line in the plane:  $y = 1 - x$



►  $x + y + z = 1$

~~~~~→ a plane in space:  $z = 1 - x - y$



►  $x + y + z + w = 1$




# One Linear Equation

What does the solution set of a linear equation look like?


►  $x + y = 1$

~~~~~> a line in the plane:  $y = 1 - x$



►  $x + y + z = 1$

~~~~~> a plane in space:  $z = 1 - x - y$



►  $x + y + z + w = 1$

~~~~~> a "3-plane" in "4-space"...

[not pictured here]

# Systems of Linear Equations

What does the solution set of a *system* of more than one linear equation look like?

# Systems of Linear Equations

What does the solution set of a *system* of more than one linear equation look like?

$$x - 3y = -3$$


$$2x + y = 8$$

# Systems of Linear Equations

What does the solution set of a *system* of more than one linear equation look like?

$$x - 3y = -3$$

$$2x + y = 8$$




# Systems of Linear Equations

What does the solution set of a *system* of more than one linear equation look like?

$$x - 3y = -3$$

$$2x + y = 8$$



In general it's an intersection of lines, planes, etc.

## Kinds of Solution Sets

In what other ways can two lines intersect?

## Kinds of Solution Sets

In what other ways can two lines intersect?

$$x - 3y = -3$$


$$x - 3y = 3$$

## Kinds of Solution Sets

In what other ways can two lines intersect?

$$x - 3y = -3$$

$$x - 3y = 3$$






## Kinds of Solution Sets

In what other ways can two lines intersect?

$$x - 3y = -3$$

$$x - 3y = 3$$



A system of equations with no solutions is called **inconsistent**.

## Kinds of Solution Sets

In what other ways can two lines intersect?

## Kinds of Solution Sets

In what other ways can two lines intersect?

$$x - 3y = -3$$


$$2x - 6y = -6$$

## Kinds of Solution Sets

In what other ways can two lines intersect?

$$x - 3y = -3$$


$$2x - 6y = -6$$



## Kinds of Solution Sets

In what other ways can two lines intersect?

$$\begin{aligned}x - 3y &= -3 \\ 2x - 6y &= -6\end{aligned}$$




Note that multiplying an equation by a nonzero number gives the *same solution set*.

## Kinds of Solution Sets

In what other ways can two lines intersect?

$$\begin{aligned}x - 3y &= -3 \\ 2x - 6y &= -6\end{aligned}$$



Note that multiplying an equation by a nonzero number gives the *same solution set*. In other words, they are *equivalent* (systems of) equations.

## Poll

What about in three variables?

What about in three variables?

Poll

In how many different ways can three planes intersect in space?

- A. One
- B. Two
- C. Three
- D. Four
- E. Five
- F. Six
- G. Seven



# Solving Systems of Equations

## Example

Solve the system of equations

$$x + 2y + 3z = 6$$

$$2x - 3y + 2z = 14$$

$$3x + y - z = -2$$

This is the kind of problem we'll talk about for the first half of the course.

# Solving Systems of Equations

## Example

Solve the system of equations

$$x + 2y + 3z = 6$$

$$2x - 3y + 2z = 14$$

$$3x + y - z = -2$$

This is the kind of problem we'll talk about for the first half of the course.

- ▶ A **solution** is a list of numbers  $x, y, z, \dots$  that make *all* of the equations true.

# Solving Systems of Equations

## Example

Solve the system of equations

$$\begin{aligned}x + 2y + 3z &= 6 \\ 2x - 3y + 2z &= 14 \\ 3x + y - z &= -2\end{aligned}$$

This is the kind of problem we'll talk about for the first half of the course.

- ▶ A **solution** is a list of numbers  $x, y, z, \dots$  that make *all* of the equations true.
- ▶ The **solution set** is the collection of all solutions.

# Solving Systems of Equations

## Example

Solve the system of equations

$$\begin{aligned}x + 2y + 3z &= 6 \\ 2x - 3y + 2z &= 14 \\ 3x + y - z &= -2\end{aligned}$$

This is the kind of problem we'll talk about for the first half of the course.

- ▶ A **solution** is a list of numbers  $x, y, z, \dots$  that make *all* of the equations true.
- ▶ The **solution set** is the collection of all solutions.
- ▶ **Solving** the system means finding the solution set.

# Solving Systems of Equations

## Example

Solve the system of equations

$$x + 2y + 3z = 6$$

$$2x - 3y + 2z = 14$$

$$3x + y - z = -2$$

This is the kind of problem we'll talk about for the first half of the course.

- ▶ A **solution** is a list of numbers  $x, y, z, \dots$  that make *all* of the equations true.
- ▶ The **solution set** is the collection of all solutions.
- ▶ **Solving** the system means finding the solution set.

What is a *systematic* way to solve a system of equations?

# Solving Systems of Equations

## Example

Solve the system of equations

$$x + 2y + 3z = 6$$

$$2x - 3y + 2z = 14$$

$$3x + y - z = -2$$

# Solving Systems of Equations

## Example

Solve the system of equations

$$x + 2y + 3z = 6$$

$$2x - 3y + 2z = 14$$

$$3x + y - z = -2$$

What strategies do you know?

# Solving Systems of Equations

## Example

Solve the system of equations

$$x + 2y + 3z = 6$$

$$2x - 3y + 2z = 14$$

$$3x + y - z = -2$$

**Elimination method:** in what ways can you manipulate the equations?



# Solving Systems of Equations

## Example

Solve the system of equations

$$x + 2y + 3z = 6$$

$$2x - 3y + 2z = 14$$

$$3x + y - z = -2$$

# Solving Systems of Equations

## Example

Solve the system of equations

$$x + 2y + 3z = 6$$

$$2x - 3y + 2z = 14$$

$$3x + y - z = -2$$

Now I've eliminated  $x$  from the last equation!

# Solving Systems of Equations

## Example

Solve the system of equations

$$x + 2y + 3z = 6$$

$$2x - 3y + 2z = 14$$

$$3x + y - z = -2$$

Now I've eliminated  $x$  from the last equation!

...but there's a long way to go still. Can we make our lives easier?

# Solving Systems of Equations

Better notation

It sure is a pain to have to write  $x, y, z$ , and  $=$  over and over again.

# Solving Systems of Equations

Better notation

It sure is a pain to have to write  $x, y, z$ , and  $=$  over and over again.

**Matrix notation:** write just the numbers, in a box, instead!

$$\begin{array}{rcl} x + 2y + 3z & = & 6 \\ 2x - 3y + 2z & = & 14 \\ 3x + y - z & = & -2 \end{array} \quad \begin{array}{c} \text{becomes} \\ \text{~~~~~} \rightarrow \end{array} \quad \left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 2 & -3 & 2 & 14 \\ 3 & 1 & -1 & -2 \end{array} \right)$$

# Solving Systems of Equations

Better notation

It sure is a pain to have to write  $x, y, z$ , and  $=$  over and over again.

**Matrix notation:** write just the numbers, in a box, instead!

$$\begin{array}{rcl} x + 2y + 3z & = & 6 \\ 2x - 3y + 2z & = & 14 \\ 3x + y - z & = & -2 \end{array} \quad \begin{array}{c} \text{becomes} \\ \text{~~~~~} \end{array} \quad \left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 2 & -3 & 2 & 14 \\ 3 & 1 & -1 & -2 \end{array} \right)$$

This is called an **(augmented) matrix**. Our equation manipulations become **elementary row operations**:

# Solving Systems of Equations

Better notation

It sure is a pain to have to write  $x, y, z$ , and  $=$  over and over again.

**Matrix notation:** write just the numbers, in a box, instead!

$$\begin{array}{rcl} x + 2y + 3z & = & 6 \\ 2x - 3y + 2z & = & 14 \\ 3x + y - z & = & -2 \end{array} \quad \begin{array}{c} \text{becomes} \\ \text{~~~~~} \end{array} \quad \left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 2 & -3 & 2 & 14 \\ 3 & 1 & -1 & -2 \end{array} \right)$$

This is called an **(augmented) matrix**. Our equation manipulations become **elementary row operations**:

- ▶ Multiply all entries in a row by a nonzero number.

(scale)

# Solving Systems of Equations

Better notation

It sure is a pain to have to write  $x, y, z$ , and  $=$  over and over again.

**Matrix notation:** write just the numbers, in a box, instead!

$$\begin{array}{rcl} x + 2y + 3z & = & 6 \\ 2x - 3y + 2z & = & 14 \\ 3x + y - z & = & -2 \end{array} \quad \begin{array}{c} \text{becomes} \\ \text{~~~~~} \end{array} \quad \left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 2 & -3 & 2 & 14 \\ 3 & 1 & -1 & -2 \end{array} \right)$$

This is called an **(augmented) matrix**. Our equation manipulations become **elementary row operations**:

- ▶ Multiply all entries in a row by a nonzero number. (scale)
- ▶ Add a multiple of each entry of one row to the corresponding entry in another. (row replacement)



# Solving Systems of Equations

Better notation

It sure is a pain to have to write  $x, y, z$ , and  $=$  over and over again.

**Matrix notation:** write just the numbers, in a box, instead!

$$\begin{array}{rcl} x + 2y + 3z & = & 6 \\ 2x - 3y + 2z & = & 14 \\ 3x + y - z & = & -2 \end{array} \quad \begin{array}{c} \text{becomes} \\ \text{~~~~~} \end{array} \quad \left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 2 & -3 & 2 & 14 \\ 3 & 1 & -1 & -2 \end{array} \right)$$

This is called an **(augmented) matrix**. Our equation manipulations become **elementary row operations**:

- ▶ Multiply all entries in a row by a nonzero number. (scale)
- ▶ Add a multiple of each entry of one row to the corresponding entry in another. (row replacement)
- ▶ Swap two rows. (swap)

## Row Operations

### Example

Solve the system of equations

$$x + 2y + 3z = 6$$

$$2x - 3y + 2z = 14$$

$$3x + y - z = -2$$

## Row Operations

### Example

Solve the system of equations

$$x + 2y + 3z = 6$$

$$2x - 3y + 2z = 14$$

$$3x + y - z = -2$$

Start:

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 2 & -3 & 2 & 14 \\ 3 & 1 & -1 & -2 \end{array} \right)$$

# Row Operations

## Example

Solve the system of equations

$$x + 2y + 3z = 6$$

$$2x - 3y + 2z = 14$$

$$3x + y - z = -2$$

Start:

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 2 & -3 & 2 & 14 \\ 3 & 1 & -1 & -2 \end{array} \right)$$

**Goal:** we want our elimination method to eventually produce a system of equations like

$$x = A$$

$$y = B$$

$$z = C$$

# Row Operations

## Example

Solve the system of equations

$$x + 2y + 3z = 6$$

$$2x - 3y + 2z = 14$$

$$3x + y - z = -2$$

Start:

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 2 & -3 & 2 & 14 \\ 3 & 1 & -1 & -2 \end{array} \right)$$

**Goal:** we want our elimination method to eventually produce a system of equations like

$$x = A$$

$$y = B$$

$$z = C$$

or in matrix form,

# Row Operations

## Example

Solve the system of equations

$$x + 2y + 3z = 6$$

$$2x - 3y + 2z = 14$$

$$3x + y - z = -2$$

Start:

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 2 & -3 & 2 & 14 \\ 3 & 1 & -1 & -2 \end{array} \right)$$

**Goal:** we want our elimination method to eventually produce a system of equations like

$$x = A$$

$$y = B \quad \text{or in matrix form,}$$

$$z = C$$

So we need to do row operations that make the start matrix look like the end one.

# Row Operations

## Example

Solve the system of equations

$$x + 2y + 3z = 6$$

$$2x - 3y + 2z = 14$$

$$3x + y - z = -2$$

Start:

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 2 & -3 & 2 & 14 \\ 3 & 1 & -1 & -2 \end{array} \right)$$

**Goal:** we want our elimination method to eventually produce a system of equations like

$$x = A$$

$$y = B$$

$$z = C$$

or in matrix form,

So we need to do row operations that make the start matrix look like the end one.

**Strategy:** fiddle with it so we only have ones and zeros.

# Row Operations

Continued

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 2 & -3 & 2 & 14 \\ 3 & 1 & -1 & -2 \end{array} \right)$$



# Row Operations


Continued

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 2 & -3 & 2 & 14 \\ 3 & 1 & -1 & -2 \end{array} \right)$$

We want these to be zero.

# Row Operations

Continued


$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 2 & -3 & 2 & 14 \\ 3 & 1 & -1 & -2 \end{array} \right)$$


We want these to be zero.

So we subtract multiples of the first row.

# Row Operations

Continued

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 2 & -3 & 2 & 14 \\ 3 & 1 & -1 & -2 \end{array} \right)$$



We want these to be zero.

So we subtract multiples of the first row.

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 0 & -7 & -4 & 2 \\ 0 & -5 & -10 & -20 \end{array} \right)$$


# Row Operations

Continued

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 2 & -3 & 2 & 14 \\ 3 & 1 & -1 & -2 \end{array} \right)$$


We want these to be zero.

So we subtract multiples of the first row.

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 0 & -7 & -4 & 2 \\ 0 & -5 & -10 & -20 \end{array} \right)$$


We want these to be zero.

# Row Operations

Continued

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 2 & -3 & 2 & 14 \\ 3 & 1 & -1 & -2 \end{array} \right)$$

We want these to be zero.  
So we subtract multiples of the first row.

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 0 & -7 & -4 & 2 \\ 0 & -5 & -10 & -20 \end{array} \right)$$

We want these to be zero.

It would be nice if this were a 1.

# Row Operations

Continued

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 2 & -3 & 2 & 14 \\ 3 & 1 & -1 & -2 \end{array} \right)$$

We want these to be zero.  
So we subtract multiples of the first row.


$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 0 & -7 & -4 & 2 \\ 0 & -5 & -10 & -20 \end{array} \right)$$

We want these to be zero.


It would be nice if this were a 1.  
We could divide by  $-7$ , but that  
would produce ugly fractions.

# Row Operations

Continued

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 2 & -3 & 2 & 14 \\ 3 & 1 & -1 & -2 \end{array} \right)$$


We want these to be zero.  
So we subtract multiples of the first row.

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & 6 \\ 0 & -7 & -4 & 2 \\ 0 & -5 & -10 & -20 \end{array} \right)$$


We want these to be zero.

It would be nice if this were a 1.  
We could divide by  $-7$ , but that  
would produce ugly fractions.

Let's swap the last two rows first.

# Row Operations


Continued

$$\left( \begin{array}{ccc|c} 1 & 0 & -1 & -2 \\ 0 & 1 & 2 & 4 \\ 0 & 0 & 10 & 30 \end{array} \right)$$



# Row Operations

Continued

$$\left( \begin{array}{ccc|c} 1 & 0 & -1 & -2 \\ 0 & 1 & 2 & 4 \\ 0 & 0 & 10 & 30 \end{array} \right)$$


We want these to be zero.

# Row Operations

Continued

$$\left( \begin{array}{cc|c} 1 & 0 & -1 & -2 \\ 0 & 1 & 2 & 4 \\ 0 & 0 & 10 & 30 \end{array} \right)$$

We want these to be zero.

Let's make this a 1 first.

# Row Operations

Continued

$$\left( \begin{array}{cc|c} 1 & 0 & -1 & -2 \\ 0 & 1 & 2 & 4 \\ 0 & 0 & 10 & 30 \end{array} \right)$$

We want these to be zero.

Let's make this a 1 first.

Success!

# Row Operations

Continued

$$\left( \begin{array}{ccc|c} 1 & 0 & -1 & -2 \\ 0 & 1 & 2 & 4 \\ 0 & 0 & 10 & 30 \end{array} \right)$$

We want these to be zero.

Let's make this a 1 first.

Success!

Check:

$$\begin{array}{rcl} x + 2y + 3z & = & 6 \\ 2x - 3y + 2z & = & 14 \\ 3x + y - z & = & -2 \end{array}$$

substitute solution  
~~~~~>

## Row Equivalence

### Important

The process of doing row operations to a matrix does not change the solution set of the corresponding linear equations!

# Row Equivalence

## Important

The process of doing row operations to a matrix does not change the solution set of the corresponding linear equations!

## Definition

Two matrices are called **row equivalent** if one can be obtained from the other by doing some number of elementary row operations.

# Row Equivalence

## Important

The process of doing row operations to a matrix does not change the solution set of the corresponding linear equations!

## Definition

Two matrices are called **row equivalent** if one can be obtained from the other by doing some number of elementary row operations.

So the linear equations of row-equivalent matrices have the *same solution set*.

## A Bad Example

### Example

Solve the system of equations

$$x + y = 2$$

$$3x + 4y = 5$$

$$4x + 5y = 9$$



# A Bad Example

## Example

Solve the system of equations

$$x + y = 2$$

$$3x + 4y = 5$$

$$4x + 5y = 9$$

Let's try doing row operations:

## A Bad Example

### Example

Solve the system of equations

$$x + y = 2$$

$$3x + 4y = 5$$

$$4x + 5y = 9$$

Let's try doing row operations:

$$\left( \begin{array}{cc|c} 1 & 1 & 2 \\ 3 & 4 & 5 \\ 4 & 5 & 9 \end{array} \right)$$

# A Bad Example

## Example


Solve the system of equations

$$x + y = 2$$

$$3x + 4y = 5$$

$$4x + 5y = 9$$

Let's try doing row operations:

First clear these by  
subtracting multiples  
of the first row. 

$$\left( \begin{array}{cc|c} 1 & 1 & 2 \\ 3 & 4 & 5 \\ 4 & 5 & 9 \end{array} \right)$$

# A Bad Example

## Example

Solve the system of equations

$$x + y = 2$$

$$3x + 4y = 5$$

$$4x + 5y = 9$$

Let's try doing row operations:

First clear these by  
subtracting multiples  
of the first row.

$$\left( \begin{array}{cc|c} 1 & 1 & 2 \\ 3 & 4 & 5 \\ 4 & 5 & 9 \end{array} \right)$$

$$\left( \begin{array}{cc|c} 1 & 1 & 2 \\ 0 & 1 & -1 \\ 0 & 1 & 1 \end{array} \right)$$

# A Bad Example

## Example

Solve the system of equations

$$x + y = 2$$

$$3x + 4y = 5$$

$$4x + 5y = 9$$

Let's try doing row operations:

First clear these by  
subtracting multiples  
of the first row.

$$\left( \begin{array}{cc|c} 1 & 1 & 2 \\ 3 & 4 & 5 \\ 4 & 5 & 9 \end{array} \right)$$

Now clear this by  
subtracting  
the second row.

$$\left( \begin{array}{cc|c} 1 & 1 & 2 \\ 0 & 1 & -1 \\ 0 & 1 & 1 \end{array} \right)$$

# A Bad Example

Continued

$$\left( \begin{array}{cc|c} 1 & 1 & 2 \\ 0 & 1 & -1 \\ 0 & 0 & 2 \end{array} \right) \begin{array}{l} \text{translates into} \\ \text{~~~~~}\rightarrow \end{array}$$

# A Bad Example

Continued

$$\left( \begin{array}{cc|c} 1 & 1 & 2 \\ 0 & 1 & -1 \\ 0 & 0 & 2 \end{array} \right) \begin{array}{l} \text{translates into} \\ \text{~~~~~}\rightarrow \end{array}$$

In other words, the original equations

$$\begin{array}{lcl} x + y = 2 \\ 3x + 4y = 5 \\ 4x + 5y = 9 \end{array} \quad \text{have the same solutions as}$$

$$\begin{array}{lcl} x + y = 2 \\ y = -1 \\ 0 = 2 \end{array}$$

## Continued

translates into

$$0 = 2$$

But the latter system obviously has no solutions (there is no way to make them all true), so our original system has no solutions either.



## Continued

translates into

$$\begin{array}{rcl} x + y = 2 & & x + y = 2 \\ 3x + 4y = 5 & \text{have the same solutions as} & y = -1 \\ 4x + 5y = 9 & & 0 = 2 \end{array}$$

## Definition

A system of equations is called **inconsistent** if it has no solution. It is **consistent** otherwise.