

Section 2.8

Subspaces of \mathbf{R}^n

Motivation

Today we will discuss **subspaces** of \mathbf{R}^n .

Motivation

Today we will discuss **subspaces** of \mathbf{R}^n .


A subspace turns out to be the same as a span, except we don't know *which* vectors it's the span of.

Motivation

Today we will discuss **subspaces** of \mathbf{R}^n .

A subspace turns out to be the same as a span, except we don't know *which* vectors it's the span of.

This arises naturally when you have, say, a plane through the origin in \mathbf{R}^3 which is *not* defined (a priori) as a span, but you still want to say something about it.



Definition of Subspace

Definition

A **subspace** of \mathbf{R}^n is a subset V of \mathbf{R}^n satisfying:

1. The zero vector is in V .

Definition of Subspace

Definition

A **subspace** of \mathbf{R}^n is a subset V of \mathbf{R}^n satisfying:

1. The zero vector is in V .

“not empty”

Definition of Subspace

Definition

A **subspace** of \mathbf{R}^n is a subset V of \mathbf{R}^n satisfying:

1. The zero vector is in V .
2. If u and v are in V , then $u + v$ is also in V .

“not empty”

Definition of Subspace

Definition

A **subspace** of \mathbf{R}^n is a subset V of \mathbf{R}^n satisfying:

1. The zero vector is in V .
2. If u and v are in V , then $u + v$ is also in V .

“not empty”

“closed under addition”

Definition of Subspace

Definition

A **subspace** of \mathbf{R}^n is a subset V of \mathbf{R}^n satisfying:

1. The zero vector is in V .
2. If u and v are in V , then $u + v$ is also in V .
3. If u is in V and c is in \mathbf{R} , then cu is in V .

“not empty”

“closed under addition”

Definition of Subspace

Definition

A **subspace** of \mathbf{R}^n is a subset V of \mathbf{R}^n satisfying:

1. The zero vector is in V .
2. If u and v are in V , then $u + v$ is also in V .
3. If u is in V and c is in \mathbf{R} , then cu is in V .

“not empty”

“closed under addition”

“closed under \times scalars”

Definition of Subspace

Definition

A **subspace** of \mathbf{R}^n is a subset V of \mathbf{R}^n satisfying:

1. The zero vector is in V . "not empty"
2. If u and v are in V , then $u + v$ is also in V . "closed under addition"
3. If u is in V and c is in \mathbf{R} , then cu is in V . "closed under \times scalars"

What does this mean?

- If v is in V , then all scalar multiples of v are in V by (3).

Definition of Subspace

Definition

A **subspace** of \mathbf{R}^n is a subset V of \mathbf{R}^n satisfying:

1. The zero vector is in V . "not empty"
2. If u and v are in V , then $u + v$ is also in V . "closed under addition"
3. If u is in V and c is in \mathbf{R} , then cu is in V . "closed under \times scalars"

What does this mean?

- If v is in V , then all scalar multiples of v are in V by (3). That is, the line through v is in V .

Definition of Subspace

Definition

A **subspace** of \mathbf{R}^n is a subset V of \mathbf{R}^n satisfying:

1. The zero vector is in V . "not empty"
2. If u and v are in V , then $u + v$ is also in V . "closed under addition"
3. If u is in V and c is in \mathbf{R} , then cu is in V . "closed under \times scalars"

What does this mean?

- ▶ If v is in V , then all scalar multiples of v are in V by (3). That is, the line through v is in V .
- ▶ If u, v are in V , then xu and yv are in V for scalars x, y by (3).

Definition of Subspace

Definition

A **subspace** of \mathbf{R}^n is a subset V of \mathbf{R}^n satisfying:

1. The zero vector is in V . "not empty"
2. If u and v are in V , then $u + v$ is also in V . "closed under addition"
3. If u is in V and c is in \mathbf{R} , then cu is in V . "closed under \times scalars"

What does this mean?

- ▶ If v is in V , then all scalar multiples of v are in V by (3). That is, the line through v is in V .
- ▶ If u, v are in V , then xu and yv are in V for scalars x, y by (3). So $xu + yv$ is in V by (2).

Definition of Subspace

Definition

A **subspace** of \mathbf{R}^n is a subset V of \mathbf{R}^n satisfying:

1. The zero vector is in V . "not empty"
2. If u and v are in V , then $u + v$ is also in V . "closed under addition"
3. If u is in V and c is in \mathbf{R} , then cu is in V . "closed under \times scalars"

What does this mean?

- ▶ If v is in V , then all scalar multiples of v are in V by (3). That is, the line through v is in V .
- ▶ If u, v are in V , then xu and yv are in V for scalars x, y by (3). So $xu + yv$ is in V by (2). So $\text{Span}\{u, v\}$ is contained in V .

Definition of Subspace

Definition

A **subspace** of \mathbf{R}^n is a subset V of \mathbf{R}^n satisfying:

1. The zero vector is in V . "not empty"
2. If u and v are in V , then $u + v$ is also in V . "closed under addition"
3. If u is in V and c is in \mathbf{R} , then cu is in V . "closed under \times scalars"

What does this mean?

- ▶ If v is in V , then all scalar multiples of v are in V by (3). That is, the line through v is in V .
- ▶ If u, v are in V , then xu and yv are in V for scalars x, y by (3). So $xu + yv$ is in V by (2). So $\text{Span}\{u, v\}$ is contained in V .
- ▶ Likewise, if v_1, v_2, \dots, v_n are all in V , then $\text{Span}\{v_1, v_2, \dots, v_n\}$ is contained in V .

Definition of Subspace

Definition

A **subspace** of \mathbf{R}^n is a subset V of \mathbf{R}^n satisfying:

1. The zero vector is in V . "not empty"
2. If u and v are in V , then $u + v$ is also in V . "closed under addition"
3. If u is in V and c is in \mathbf{R} , then cu is in V . "closed under \times scalars"

What does this mean?


- ▶ If v is in V , then all scalar multiples of v are in V by (3). That is, the line through v is in V .
- ▶ If u, v are in V , then xu and yv are in V for scalars x, y by (3). So $xu + yv$ is in V by (2). So $\text{Span}\{u, v\}$ is contained in V .
- ▶ Likewise, if v_1, v_2, \dots, v_n are all in V , then $\text{Span}\{v_1, v_2, \dots, v_n\}$ is contained in V .

A subspace V contains the span of any set of vectors in V .

Examples

Example


A line L through the origin: this contains the span of any vector in L .



Examples


Example

A line L through the origin: this contains the span of any vector in L .



Example


A plane P through the origin: this contains the span of any vectors in P .



Examples


Example

A line L through the origin: this contains the span of any vector in L .



Example

A plane P through the origin: this contains the span of any vectors in P .




Example

All of \mathbf{R}^n : this contains 0 , and is closed under addition and scalar multiplication.

Examples


Example

A line L through the origin: this contains the span of any vector in L .



Example

A plane P through the origin: this contains the span of any vectors in P .



Example

All of \mathbf{R}^n : this contains 0 , and is closed under addition and scalar multiplication.


Example

The subset $\{0\}$: this subspace contains only one vector.

Examples


Example

A line L through the origin: this contains the span of any vector in L .



Example

A plane P through the origin: this contains the span of any vectors in P .



Example

All of \mathbf{R}^n : this contains 0 , and is closed under addition and scalar multiplication.

Example

The subset $\{0\}$: this subspace contains only one vector.


Note these are all pictures of spans! (Line, plane, space, etc.)

Non-Examples

Non-Example

A line L (or any other set) that doesn't contain the origin is not a subspace.

Fails:




Non-Examples

Non-Example

A line L (or any other set) that doesn't contain the origin is not a subspace.

Fails: 1.




Non-Examples

Non-Example


A line L (or any other set) that doesn't contain the origin is not a subspace.

Fails: 1.



Non-Example

A circle C is not a subspace. Fails:



Non-Examples

Non-Example


A line L (or any other set) that doesn't contain the origin is not a subspace.

Fails: 1.

Non-Example

A circle C is not a subspace. Fails:

1,2,3.



Non-Examples


Non-Example

A line L (or any other set) that doesn't contain the origin is not a subspace.

Fails: 1.

Non-Example


A circle C is not a subspace. Fails: 1,2,3. Think: a circle isn't a "linear space."



Non-Examples


Non-Example

A line L (or any other set) that doesn't contain the origin is not a subspace. Fails: 1.




Non-Example

A circle C is not a subspace. Fails: 1,2,3. Think: a circle isn't a "linear space."



Non-Example


The first quadrant in \mathbf{R}^2 is not a subspace. Fails:



Non-Examples


Non-Example

A line L (or any other set) that doesn't contain the origin is not a subspace. Fails: 1.




Non-Example

A circle C is not a subspace. Fails: 1,2,3. Think: a circle isn't a "linear space."



Non-Example


The first quadrant in \mathbf{R}^2 is not a subspace. Fails: 3 only.



Non-Examples


Non-Example

A line L (or any other set) that doesn't contain the origin is not a subspace. Fails: 1.




Non-Example

A circle C is not a subspace. Fails: 1,2,3. Think: a circle isn't a "linear space."




Non-Example

The first quadrant in \mathbf{R}^2 is not a subspace. Fails: 3 only.



Non-Example

A line union a plane in \mathbf{R}^3 is not a subspace. Fails:




Non-Examples

Non-Example


A line L (or any other set) that doesn't contain the origin is not a subspace.

Fails: 1.




Non-Example

A circle C is not a subspace. Fails: 1,2,3. Think: a circle isn't a "linear space."




Non-Example

The first quadrant in \mathbf{R}^2 is not a subspace. Fails: 3 only.



Non-Example

A line union a plane in \mathbf{R}^3 is not a subspace. Fails: 2 only.



Subsets and Subspaces

They aren't the same thing

A **subset** of \mathbf{R}^n is any collection of vectors whatsoever.

Subsets and Subspaces

They aren't the same thing

A **subset** of \mathbf{R}^n is any collection of vectors whatsoever.

All of the non-examples are still subsets.

Subsets and Subspaces

They aren't the same thing

A **subset** of \mathbf{R}^n is any collection of vectors whatsoever.

All of the non-examples are still subsets.

A **subspace** is a special kind of subset, which satisfies the three defining properties.


Subsets and Subspaces

They aren't the same thing

A **subset** of \mathbf{R}^n is any collection of vectors whatsoever.

All of the non-examples are still subsets.

A **subspace** is a special kind of subset, which satisfies the three defining properties.



Subset: *yes*

Subspace: *no*

Spans are Subspaces

Theorem

Any $\text{Span}\{v_1, v_2, \dots, v_n\}$ is a subspace.

Spans are Subspaces

Theorem

Any $\text{Span}\{v_1, v_2, \dots, v_n\}$ is a subspace.

!!!

Every subspace is a span, and every span is a subspace.

Spans are Subspaces

Theorem

Any $\text{Span}\{v_1, v_2, \dots, v_n\}$ is a subspace.

!!!

Every subspace is a span, and every span is a subspace.

Definition

If $V = \text{Span}\{v_1, v_2, \dots, v_n\}$, we say that V is the subspace **generated by** or **spanned by** the vectors v_1, v_2, \dots, v_n .

Poll

Is the empty set $\{\}$ a subspace? If not, which property(ies) does it fail?

Poll

Is the empty set $\{\}$ a subspace? If not, which property(ies) does it fail?

The zero vector is not contained in the empty set, so it is *not* a subspace.

Poll

Is the empty set $\{\}$ a subspace? If not, which property(ies) does it fail?

The zero vector is not contained in the empty set, so it is *not* a subspace.

Question: What is the difference between $\{\}$ and $\{0\}$?

Subspaces

Verification

Let $V = \left\{ \begin{pmatrix} a \\ b \end{pmatrix} \text{ in } \mathbf{R}^2 \mid ab = 0 \right\}$. Let's check if V is a subspace or not.

Subspaces

Verification


Let $V = \left\{ \begin{pmatrix} a \\ b \end{pmatrix} \text{ in } \mathbf{R}^2 \mid ab = 0 \right\}$. Let's check if V is a subspace or not.

We conclude that V is *not* a subspace.

Subspaces

Verification

Let $V = \left\{ \begin{pmatrix} a \\ b \end{pmatrix} \text{ in } \mathbf{R}^2 \mid ab = 0 \right\}$. Let's check if V is a subspace or not.



We conclude that V is *not* a subspace. A picture is above. (It doesn't look like a span.)

Column Space and Null Space

An $m \times n$ matrix A naturally gives rise to *two* subspaces.

Column Space and Null Space

An $m \times n$ matrix A naturally gives rise to *two* subspaces.

Definition

- ▶ The **column space** of A is the subspace of \mathbf{R}^m —spanned by the columns of A . It is written $\text{Col } A$.

Column Space and Null Space

An $m \times n$ matrix A naturally gives rise to *two* subspaces.

Definition

- ▶ The **column space** of A is the subspace of \mathbf{R}^m spanned by the columns of A . It is written $\text{Col } A$.

Column Space and Null Space

An $m \times n$ matrix A naturally gives rise to *two* subspaces.

Definition

- ▶ The **column space** of A is the subspace of \mathbf{R}^m spanned by the columns of A . It is written $\text{Col } A$.
- ▶ The **null space** of A is the set of all solutions of the homogeneous equation $Ax = 0$:

$$\text{Nul } A = \{x \text{ in } \mathbf{R}^n \mid Ax = 0\}.$$

This is a subspace of \mathbf{R}^n .

Column Space and Null Space

An $m \times n$ matrix A naturally gives rise to *two* subspaces.

Definition

- ▶ The **column space** of A is the subspace of \mathbf{R}^m spanned by the columns of A . It is written $\text{Col } A$.
- ▶ The **null space** of A is the set of all solutions of the homogeneous equation $Ax = 0$:

$$\text{Nul } A = \{x \text{ in } \mathbf{R}^n \mid Ax = 0\}.$$

This is a subspace of \mathbf{R}^n .

Column Space and Null Space

An $m \times n$ matrix A naturally gives rise to *two* subspaces.

Definition

- ▶ The **column space** of A is the subspace of \mathbf{R}^m spanned by the columns of A . It is written $\text{Col } A$.
- ▶ The **null space** of A is the set of all solutions of the homogeneous equation $Ax = 0$:

$$\text{Nul } A = \{x \text{ in } \mathbf{R}^n \mid Ax = 0\}.$$

This is a subspace of \mathbf{R}^n .

The column space is defined as a span, so we know it is a subspace.

Column Space and Null Space

An $m \times n$ matrix A naturally gives rise to *two* subspaces.

Definition

- ▶ The **column space** of A is the subspace of \mathbf{R}^m spanned by the columns of A . It is written $\text{Col } A$.
- ▶ The **null space** of A is the set of all solutions of the homogeneous equation $Ax = 0$:

$$\text{Nul } A = \{x \text{ in } \mathbf{R}^n \mid Ax = 0\}.$$

This is a subspace of \mathbf{R}^n .

The column space is defined as a span, so we know it is a subspace. It is the range (as opposed to the codomain) of the transformation $T(x) = Ax$.

Column Space and Null Space

An $m \times n$ matrix A naturally gives rise to *two* subspaces.

Definition

- ▶ The **column space** of A is the subspace of \mathbf{R}^m spanned by the columns of A . It is written $\text{Col } A$.
- ▶ The **null space** of A is the set of all solutions of the homogeneous equation $Ax = 0$:

$$\text{Nul } A = \{x \text{ in } \mathbf{R}^n \mid Ax = 0\}.$$

This is a subspace of \mathbf{R}^n .

The column space is defined as a span, so we know it is a subspace. It is the range (as opposed to the codomain) of the transformation $T(x) = Ax$.

Check that the null space is a subspace:

Column Space and Null Space

Example

$$\text{Let } A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \\ 1 & 1 \end{pmatrix}.$$


Let's compute the column space:

Column Space and Null Space

Example

$$\text{Let } A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \\ 1 & 1 \end{pmatrix}.$$

Let's compute the column space:




Column Space and Null Space

Example

$$\text{Let } A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \\ 1 & 1 \end{pmatrix}.$$

Let's compute the column space:




Let's compute the null space:

Column Space and Null Space


Example

$$\text{Let } A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \\ 1 & 1 \end{pmatrix}.$$

Let's compute the column space:



Let's compute the null space:



The Null Space is a Span

The column space of a matrix A is defined to be a span (of the columns).

The Null Space is a Span

The column space of a matrix A is defined to be a span (of the columns).

The null space is defined to be the solution set to $Ax = 0$.

The Null Space is a Span

The column space of a matrix A is defined to be a span (of the columns).

The null space is defined to be the solution set to $Ax = 0$. It is a subspace, so it is a span.

The Null Space is a Span

The column space of a matrix A is defined to be a span (of the columns).

The null space is defined to be the solution set to $Ax = 0$. It is a subspace, so it is a span.

Question

How to find vectors which span the null space?

The Null Space is a Span

The column space of a matrix A is defined to be a span (of the columns).

The null space is defined to be the solution set to $Ax = 0$. It is a subspace, so it is a span.

Question

How to find vectors which span the null space?

Answer: Parametric vector form!

The Null Space is a Span

The column space of a matrix A is defined to be a span (of the columns).

The null space is defined to be the solution set to $Ax = 0$. It is a subspace, so it is a span.

Question

How to find vectors which span the null space?

Answer: Parametric vector form! We know that the solution set to $Ax = 0$ has a parametric form that looks like

$$x_3 \begin{pmatrix} 1 \\ 2 \\ 1 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} -2 \\ 3 \\ 0 \\ 1 \end{pmatrix} \quad \begin{array}{l} \text{if, say, } x_3 \text{ and } x_4 \\ \text{are the free} \\ \text{variables.} \end{array}$$

The Null Space is a Span

The column space of a matrix A is defined to be a span (of the columns).

The null space is defined to be the solution set to $Ax = 0$. It is a subspace, so it is a span.

Question

How to find vectors which span the null space?

Answer: Parametric vector form! We know that the solution set to $Ax = 0$ has a parametric form that looks like

$$x_3 \begin{pmatrix} 1 \\ 2 \\ 1 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} -2 \\ 3 \\ 0 \\ 1 \end{pmatrix} \quad \text{if, say, } x_3 \text{ and } x_4 \text{ are the free variables. So} \quad \text{Nul } A = \text{Span} \left\{ \begin{pmatrix} 1 \\ 2 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -2 \\ 3 \\ 0 \\ 1 \end{pmatrix} \right\}.$$

The Null Space is a Span

The column space of a matrix A is defined to be a span (of the columns).

The null space is defined to be the solution set to $Ax = 0$. It is a subspace, so it is a span.

Question

How to find vectors which span the null space?

Answer: Parametric vector form! We know that the solution set to $Ax = 0$ has a parametric form that looks like

$$x_3 \begin{pmatrix} 1 \\ 2 \\ 1 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} -2 \\ 3 \\ 0 \\ 1 \end{pmatrix} \quad \begin{array}{l} \text{if, say, } x_3 \text{ and } x_4 \\ \text{are the free} \\ \text{variables. So} \end{array} \quad \text{Nul } A = \text{Span} \left\{ \begin{pmatrix} 1 \\ 2 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -2 \\ 3 \\ 0 \\ 1 \end{pmatrix} \right\}.$$

Refer back to the slides for §1.5 (Solution Sets).

The Null Space is a Span

The column space of a matrix A is defined to be a span (of the columns).

The null space is defined to be the solution set to $Ax = 0$. It is a subspace, so it is a span.

Question

How to find vectors which span the null space?

Answer: Parametric vector form! We know that the solution set to $Ax = 0$ has a parametric form that looks like

$$x_3 \begin{pmatrix} 1 \\ 2 \\ 1 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} -2 \\ 3 \\ 0 \\ 1 \end{pmatrix} \quad \begin{array}{l} \text{if, say, } x_3 \text{ and } x_4 \\ \text{are the free} \\ \text{variables. So} \end{array} \quad \text{Nul } A = \text{Span} \left\{ \begin{pmatrix} 1 \\ 2 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -2 \\ 3 \\ 0 \\ 1 \end{pmatrix} \right\}.$$

Refer back to the slides for §1.5 (Solution Sets).

Note: It is much easier to define the null space first as a subspace, then find spanning vectors *later*, if we need them. This is one reason subspaces are so useful.

The Null Space is a Span


Example, revisited

Find vector(s) that span the null space of $A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \\ 1 & 1 \end{pmatrix}$.

The Null Space is a Span

Example, revisited

Find vector(s) that span the null space of $A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \\ 1 & 1 \end{pmatrix}$.



Subspaces

Summary

How do you check if a subset is a subspace?

- ▶ Is it a span? Can it be written as a span?

Subspaces

Summary

How do you check if a subset is a subspace?

- ▶ Is it a span? Can it be written as a span?
- ▶ Can it be written as the column space of a matrix?

Subspaces

Summary

How do you check if a subset is a subspace?

- ▶ Is it a span? Can it be written as a span?
- ▶ Can it be written as the column space of a matrix?
- ▶ Can it be written as the null space of a matrix?

Subspaces

Summary

How do you check if a subset is a subspace?

- ▶ Is it a span? Can it be written as a span?
- ▶ Can it be written as the column space of a matrix?
- ▶ Can it be written as the null space of a matrix?
- ▶ Is it all of \mathbf{R}^n or the zero subspace $\{0\}$?

Subspaces

Summary

How do you check if a subset is a subspace?

- ▶ Is it a span? Can it be written as a span?
- ▶ Can it be written as the column space of a matrix?
- ▶ Can it be written as the null space of a matrix?
- ▶ Is it all of \mathbf{R}^n or the zero subspace $\{0\}$?
- ▶ Can it be written as a type of subspace that we'll learn about later (eigenspaces, ...)?

Subspaces

Summary

How do you check if a subset is a subspace?

- ▶ Is it a span? Can it be written as a span?
- ▶ Can it be written as the column space of a matrix?
- ▶ Can it be written as the null space of a matrix?
- ▶ Is it all of \mathbf{R}^n or the zero subspace $\{0\}$?
- ▶ Can it be written as a type of subspace that we'll learn about later (eigenspaces, ...)?

If so, then it's automatically a subspace.

Subspaces

Summary

How do you check if a subset is a subspace?

- ▶ Is it a span? Can it be written as a span?
- ▶ Can it be written as the column space of a matrix?
- ▶ Can it be written as the null space of a matrix?
- ▶ Is it all of \mathbf{R}^n or the zero subspace $\{0\}$?
- ▶ Can it be written as a type of subspace that we'll learn about later (eigenspaces, ...)?

If so, then it's automatically a subspace.

If all else fails:

Subspaces

Summary

How do you check if a subset is a subspace?

- ▶ Is it a span? Can it be written as a span?
- ▶ Can it be written as the column space of a matrix?
- ▶ Can it be written as the null space of a matrix?
- ▶ Is it all of \mathbf{R}^n or the zero subspace $\{0\}$?
- ▶ Can it be written as a type of subspace that we'll learn about later (eigenspaces, ...)?

If so, then it's automatically a subspace.

If all else fails:

- ▶ Can you verify directly that it satisfies the three defining properties?

Basis of a Subspace

What is the *smallest number* of vectors that are needed to span a subspace?

Basis of a Subspace

What is the *smallest number* of vectors that are needed to span a subspace?

Definition

Let V be a subspace of \mathbf{R}^n . A **basis** of V is a set of vectors $\{v_1, v_2, \dots, v_m\}$ in V such that:

Basis of a Subspace

What is the *smallest number* of vectors that are needed to span a subspace?

Definition

Let V be a subspace of \mathbf{R}^n . A **basis** of V is a set of vectors $\{v_1, v_2, \dots, v_m\}$ in V such that:

1. $V = \text{Span}\{v_1, v_2, \dots, v_m\}$, and

Basis of a Subspace

What is the *smallest number* of vectors that are needed to span a subspace?

Definition

Let V be a subspace of \mathbf{R}^n . A **basis** of V is a set of vectors $\{v_1, v_2, \dots, v_m\}$ in V such that:

1. $V = \text{Span}\{v_1, v_2, \dots, v_m\}$, and
2. $\{v_1, v_2, \dots, v_m\}$ is linearly independent.

Basis of a Subspace

What is the *smallest number* of vectors that are needed to span a subspace?

Definition

Let V be a subspace of \mathbf{R}^n . A **basis** of V is a set of vectors $\{v_1, v_2, \dots, v_m\}$ in V such that:

1. $V = \text{Span}\{v_1, v_2, \dots, v_m\}$, and
2. $\{v_1, v_2, \dots, v_m\}$ is linearly independent.

The number of vectors in a basis is the **dimension** of V , and is written $\dim V$.

Basis of a Subspace

What is the *smallest number* of vectors that are needed to span a subspace?

Definition

Let V be a subspace of \mathbf{R}^n . A **basis** of V is a set of vectors $\{v_1, v_2, \dots, v_m\}$ in V such that:

1. $V = \text{Span}\{v_1, v_2, \dots, v_m\}$, and
2. $\{v_1, v_2, \dots, v_m\}$ is linearly independent.

The number of vectors in a basis is the **dimension** of V , and is written $\dim V$.

Note the big
red border here

Basis of a Subspace

What is the *smallest number* of vectors that are needed to span a subspace?

Definition

Let V be a subspace of \mathbf{R}^n . A **basis** of V is a set of vectors $\{v_1, v_2, \dots, v_m\}$ in V such that:

1. $V = \text{Span}\{v_1, v_2, \dots, v_m\}$, and
2. $\{v_1, v_2, \dots, v_m\}$ is linearly independent.

The number of vectors in a basis is the **dimension** of V , and is written $\dim V$.

Note the big
red border here

Why is a basis the smallest number of vectors needed to span?

Basis of a Subspace

What is the *smallest number* of vectors that are needed to span a subspace?

Definition

Let V be a subspace of \mathbf{R}^n . A **basis** of V is a set of vectors $\{v_1, v_2, \dots, v_m\}$ in V such that:

1. $V = \text{Span}\{v_1, v_2, \dots, v_m\}$, and
2. $\{v_1, v_2, \dots, v_m\}$ is linearly independent.

The number of vectors in a basis is the **dimension** of V , and is written $\dim V$.

Note the big
red border here

Why is a basis the smallest number of vectors needed to span?

Recall: *linearly independent* means that every time you add another vector, the span gets bigger.

Basis of a Subspace

What is the *smallest number* of vectors that are needed to span a subspace?

Definition

Let V be a subspace of \mathbf{R}^n . A **basis** of V is a set of vectors $\{v_1, v_2, \dots, v_m\}$ in V such that:

1. $V = \text{Span}\{v_1, v_2, \dots, v_m\}$, and
2. $\{v_1, v_2, \dots, v_m\}$ is linearly independent.

The number of vectors in a basis is the **dimension** of V , and is written $\dim V$.

Note the big
red border here

Why is a basis the smallest number of vectors needed to span?

Recall: *linearly independent* means that every time you add another vector, the span gets bigger.

Hence, if we remove any vector, the span gets *smaller*: so any smaller set can't span V .

Basis of a Subspace

What is the *smallest number* of vectors that are needed to span a subspace?

Definition

Let V be a subspace of \mathbf{R}^n . A **basis** of V is a set of vectors $\{v_1, v_2, \dots, v_m\}$ in V such that:

1. $V = \text{Span}\{v_1, v_2, \dots, v_m\}$, and
2. $\{v_1, v_2, \dots, v_m\}$ is linearly independent.

The number of vectors in a basis is the **dimension** of V , and is written $\dim V$.

Note the big
red border here

Why is a basis the smallest number of vectors needed to span?

Recall: *linearly independent* means that every time you add another vector, the span gets bigger.

Hence, if we remove any vector, the span gets *smaller*: so any smaller set can't span V .

Important

A subspace has *many different* bases, but they all have the same number of vectors (see the exercises in §2.9).

Bases of \mathbf{R}^2

Question

What is a basis for \mathbf{R}^2 ?

Bases of \mathbf{R}^2

Question

What is a basis for \mathbf{R}^2 ?


We need two vectors that *span* \mathbf{R}^2 and are *linearly independent*.

Bases of \mathbf{R}^2

Question

What is a basis for \mathbf{R}^2 ?

We need two vectors that *span* \mathbf{R}^2 and are *linearly independent*. $\{e_1, e_2\}$ is one basis.




Bases of \mathbf{R}^2

Question

What is a basis for \mathbf{R}^2 ?

We need two vectors that *span* \mathbf{R}^2 and are *linearly independent*. $\{e_1, e_2\}$ is one basis.

1. They span: $\begin{pmatrix} a \\ b \end{pmatrix} =$




Bases of \mathbf{R}^2

Question

What is a basis for \mathbf{R}^2 ?

We need two vectors that *span* \mathbf{R}^2 and are *linearly independent*. $\{e_1, e_2\}$ is one basis.

1. They span: $\begin{pmatrix} a \\ b \end{pmatrix} = ae_1 + be_2$.




Bases of \mathbf{R}^2

Question

What is a basis for \mathbf{R}^2 ?

We need two vectors that *span* \mathbf{R}^2 and are *linearly independent*. $\{e_1, e_2\}$ is one basis.

1. They span: $\begin{pmatrix} a \\ b \end{pmatrix} = ae_1 + be_2$.
2. They are linearly independent because they are not collinear.




Bases of \mathbf{R}^2

Question

What is a basis for \mathbf{R}^2 ?

We need two vectors that *span* \mathbf{R}^2 and are *linearly independent*. $\{e_1, e_2\}$ is one basis.

1. They span: $\begin{pmatrix} a \\ b \end{pmatrix} = ae_1 + be_2$.
2. They are linearly independent because they are not collinear.



Question

What is another basis for \mathbf{R}^2 ?


Bases of \mathbf{R}^2

Question

What is a basis for \mathbf{R}^2 ?

We need two vectors that *span* \mathbf{R}^2 and are *linearly independent*. $\{e_1, e_2\}$ is one basis.

1. They span: $\begin{pmatrix} a \\ b \end{pmatrix} = ae_1 + be_2$.
2. They are linearly independent because they are not collinear.



Question

What is another basis for \mathbf{R}^2 ?

Any two nonzero vectors that are not collinear.


Bases of \mathbf{R}^2

Question

What is a basis for \mathbf{R}^2 ?

We need two vectors that *span* \mathbf{R}^2 and are *linearly independent*. $\{e_1, e_2\}$ is one basis.


1. They span: $\begin{pmatrix} a \\ b \end{pmatrix} = ae_1 + be_2$.
2. They are linearly independent because they are not collinear.



Question

What is another basis for \mathbf{R}^2 ?

Any two nonzero vectors that are not collinear. $\left\{ \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right\}$ is also a basis.




Bases of \mathbf{R}^2

Question

What is a basis for \mathbf{R}^2 ?

We need two vectors that *span* \mathbf{R}^2 and are *linearly independent*. $\{e_1, e_2\}$ is one basis.

1. They span: $\begin{pmatrix} a \\ b \end{pmatrix} = ae_1 + be_2$.
2. They are linearly independent because they are not collinear.




Question

What is another basis for \mathbf{R}^2 ?

Any two nonzero vectors that are not collinear. $\left\{ \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right\}$ is also a basis.

1. They span: $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ has a pivot in every row.




Bases of \mathbf{R}^2

Question

What is a basis for \mathbf{R}^2 ?

We need two vectors that *span* \mathbf{R}^2 and are *linearly independent*. $\{e_1, e_2\}$ is one basis.

1. They span: $\begin{pmatrix} a \\ b \end{pmatrix} = ae_1 + be_2$.
2. They are linearly independent because they are not collinear.




Question

What is another basis for \mathbf{R}^2 ?

Any two nonzero vectors that are not collinear. $\left\{ \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right\}$ is also a basis.

1. They span: $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ has a pivot in every row.
2. They are linearly independent: $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ has a pivot in every column.



Bases of \mathbf{R}^n

The unit coordinate vectors

$$e_1 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad \dots, \quad e_{n-1} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ 0 \end{pmatrix}, \quad e_n = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}$$

are a basis for \mathbf{R}^n .

Bases of \mathbf{R}^n

The unit coordinate vectors

$$e_1 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad \dots, \quad e_{n-1} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ 0 \end{pmatrix}, \quad e_n = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}$$

are a basis for \mathbf{R}^n .

1. They span: I_n has a pivot in every row.

Bases of \mathbf{R}^n

The unit coordinate vectors

$$e_1 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad \dots, \quad e_{n-1} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ 0 \end{pmatrix}, \quad e_n = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}$$

are a basis for \mathbf{R}^n .

The identity matrix has columns e_1, e_2, \dots, e_n .

1. They span: I_n has a pivot in every row.

Bases of \mathbf{R}^n

The unit coordinate vectors

$$e_1 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad \dots, \quad e_{n-1} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ 0 \end{pmatrix}, \quad e_n = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}$$

are a basis for \mathbf{R}^n .  The identity matrix has columns e_1, e_2, \dots, e_n .

1. They span: I_n has a pivot in every row.
2. They are linearly independent: I_n has a pivot in every column.

Bases of \mathbf{R}^n

The unit coordinate vectors

$$e_1 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad \dots, \quad e_{n-1} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ 0 \end{pmatrix}, \quad e_n = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}$$

are a basis for \mathbf{R}^n .  The identity matrix has columns e_1, e_2, \dots, e_n .

1. They span: I_n has a pivot in every row.
2. They are linearly independent: I_n has a pivot in every column.

In general: $\{v_1, v_2, \dots, v_n\}$ is a basis for \mathbf{R}^n if and only if the matrix

$$A = \begin{pmatrix} | & | & & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & & | \end{pmatrix}$$

has a pivot in every row and every column, i.e. if A is _____.

Bases of \mathbf{R}^n

The unit coordinate vectors

$$e_1 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad \dots, \quad e_{n-1} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ 0 \end{pmatrix}, \quad e_n = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}$$

are a basis for \mathbf{R}^n .  The identity matrix has columns e_1, e_2, \dots, e_n .

1. They span: I_n has a pivot in every row.
2. They are linearly independent: I_n has a pivot in every column.

In general: $\{v_1, v_2, \dots, v_n\}$ is a basis for \mathbf{R}^n if and only if the matrix

$$A = \begin{pmatrix} | & | & & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & & | \end{pmatrix}$$

has a pivot in every row and every column, i.e. if A is *invertible*.

Basis of a Subspace

Example

Example

Let

$$V = \left\{ \begin{pmatrix} x \\ y \\ z \end{pmatrix} \text{ in } \mathbf{R}^3 \mid x + 3y + z = 0 \right\} \quad \mathcal{B} = \left\{ \begin{pmatrix} -3 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ -3 \end{pmatrix} \right\}.$$

Verify that \mathcal{B} is a basis for V .

Basis for $\text{Nul } A$

Fact

The vectors in the parametric vector form of the general solution to $Ax = 0$ always form a basis for $\text{Nul } A$.

Basis for $\text{Nul } A$

Fact

The vectors in the parametric vector form of the general solution to $Ax = 0$ always form a basis for $\text{Nul } A$.

Example

Basis for Col A

Fact

The *pivot columns* of A always form a basis for Col A .

Basis for Col A

Fact

The *pivot columns* of A always form a basis for Col A .

Warning: I mean the pivot columns of the *original* matrix A , not the row-reduced form.

Basis for Col A

Fact

The *pivot columns* of A always form a basis for Col A .

Warning: I mean the pivot columns of the *original* matrix A , not the row-reduced form. (Row reduction changes the column space.)

Basis for Col A

Fact

The *pivot columns* of A always form a basis for Col A .

Warning: I mean the pivot columns of the *original* matrix A , not the row-reduced form. (Row reduction changes the column space.)

Example

Basis for Col A

Fact

The *pivot columns* of A always form a basis for Col A .

Warning: I mean the pivot columns of the *original* matrix A , not the row-reduced form. (Row reduction changes the column space.)

Example

Why? End of §2.8, or ask in office hours.