

Chapter 2

Matrix Algebra

Section 2.1

Matrix Operations

Motivation

Recall: we can turn any system of linear equations into a matrix equation

$$Ax = b.$$

Motivation

Recall: we can turn any system of linear equations into a matrix equation

$$Ax = b.$$

This notation is suggestive. Can we solve the equation by “dividing by A”?

$$x \stackrel{??}{=} \frac{b}{A}$$

Motivation

Recall: we can turn any system of linear equations into a matrix equation

$$Ax = b.$$

This notation is suggestive. Can we solve the equation by “dividing by A”?

$$x \stackrel{??}{=} \frac{b}{A}$$

Answer: Sometimes, but you have to know what you're doing.

Motivation

Recall: we can turn any system of linear equations into a matrix equation

$$Ax = b.$$

This notation is suggestive. Can we solve the equation by “dividing by A”?

$$x \stackrel{??}{=} \frac{b}{A}$$

Answer: Sometimes, but you have to know what you're doing.

Today we'll study *matrix algebra*: adding and multiplying matrices.

More Notation for Matrices

Let A be an $m \times n$ matrix.

More Notation for Matrices

Let A be an $m \times n$ matrix.

We write a_{ij} for the entry in the i th row and the j th column. It is called the **ij th entry** of the matrix.

$$\begin{pmatrix} a_{11} & \cdots & a_{1j} & \cdots & a_{1n} \\ \vdots & & \vdots & & \vdots \\ a_{i1} & \cdots & a_{ij} & \cdots & a_{in} \\ \vdots & & \vdots & & \vdots \\ a_{m1} & \cdots & a_{mj} & \cdots & a_{mn} \end{pmatrix}$$

j th column

i th row

More Notation for Matrices

Let A be an $m \times n$ matrix.

We write a_{ij} for the entry in the i th row and the j th column. It is called the **ij th entry** of the matrix.

$$\begin{pmatrix} a_{11} & \cdots & a_{1j} & \cdots & a_{1n} \\ \vdots & & \vdots & & \vdots \\ a_{i1} & \cdots & a_{ij} & \cdots & a_{in} \\ \vdots & & \vdots & & \vdots \\ a_{m1} & \cdots & a_{mj} & \cdots & a_{mn} \end{pmatrix}$$

j th column

i th row

The entries $a_{11}, a_{22}, a_{33}, \dots$ are the **diagonal entries**; they form the **main diagonal** of the matrix.

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{pmatrix} \quad \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \\ a_{31} & a_{32} \end{pmatrix}$$

More Notation for Matrices

Let A be an $m \times n$ matrix.

We write a_{ij} for the entry in the i th row and the j th column. It is called the **ij th entry** of the matrix.

$$\begin{pmatrix} a_{11} & \cdots & a_{1j} & \cdots & a_{1n} \\ \vdots & & \vdots & & \vdots \\ a_{i1} & \cdots & a_{ij} & \cdots & a_{in} \\ \vdots & & \vdots & & \vdots \\ a_{m1} & \cdots & a_{mj} & \cdots & a_{mn} \end{pmatrix}$$

j th column

i th row

The entries $a_{11}, a_{22}, a_{33}, \dots$ are the **diagonal entries**; they form the **main diagonal** of the matrix.

A **diagonal matrix** is a *square* matrix whose only nonzero entries are on the main diagonal.

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{pmatrix} \quad \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \\ a_{31} & a_{32} \end{pmatrix}$$

$$\begin{pmatrix} a_{11} & 0 & 0 \\ 0 & a_{22} & 0 \\ 0 & 0 & a_{33} \end{pmatrix}$$

More Notation for Matrices

Let A be an $m \times n$ matrix.

We write a_{ij} for the entry in the i th row and the j th column. It is called the **ij th entry** of the matrix.

$$\begin{pmatrix} a_{11} & \cdots & a_{1j} & \cdots & a_{1n} \\ \vdots & & \vdots & & \vdots \\ a_{i1} & \cdots & a_{ij} & \cdots & a_{in} \\ \vdots & & \vdots & & \vdots \\ a_{m1} & \cdots & a_{mj} & \cdots & a_{mn} \end{pmatrix}$$

The entries $a_{11}, a_{22}, a_{33}, \dots$ are the **diagonal entries**; they form the **main diagonal** of the matrix.

A **diagonal matrix** is a *square* matrix whose only nonzero entries are on the main diagonal.

The $n \times n$ **identity matrix** I_n is the diagonal matrix with all diagonal entries equal to 1. It is special because $I_n \mathbf{v} = \mathbf{v}$ for all \mathbf{v} in \mathbf{R}^n .

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \quad \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \\ a_{31} & a_{32} \end{pmatrix}$$

$$\begin{pmatrix} a_{11} & 0 & 0 \\ 0 & a_{22} & 0 \\ 0 & 0 & a_{33} \end{pmatrix}$$

$$I_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

More Notation for Matrices

Let A be an $m \times n$ matrix.

We write a_{ij} for the entry in the i th row and the j th column. It is called the **ij th entry** of the matrix.

$$\begin{pmatrix} a_{11} & \cdots & a_{1j} & \cdots & a_{1n} \\ \vdots & & \vdots & & \vdots \\ a_{i1} & \cdots & a_{ij} & \cdots & a_{in} \\ \vdots & & \vdots & & \vdots \\ a_{m1} & \cdots & a_{mj} & \cdots & a_{mn} \end{pmatrix}$$

The entries $a_{11}, a_{22}, a_{33}, \dots$ are the **diagonal entries**; they form the **main diagonal** of the matrix.

A **diagonal matrix** is a *square* matrix whose only nonzero entries are on the main diagonal.

The $n \times n$ **identity matrix** I_n is the diagonal matrix with all diagonal entries equal to 1. It is special because $I_n \mathbf{v} = \mathbf{v}$ for all \mathbf{v} in \mathbf{R}^n .

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{pmatrix} \quad \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \\ a_{31} & a_{32} \end{pmatrix}$$

$$\begin{pmatrix} a_{11} & 0 & 0 \\ 0 & a_{22} & 0 \\ 0 & 0 & a_{33} \end{pmatrix}$$

$$I_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

More Notation for Matrices

Continued

The **zero matrix** (of size $m \times n$) is the $m \times n$ matrix 0 with all zero entries.

$$0 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

More Notation for Matrices

Continued

The **zero matrix** (of size $m \times n$) is the $m \times n$ matrix 0 with all zero entries.

$$0 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

The **transpose** of an $m \times n$ matrix A is the $n \times m$ matrix A^T whose rows are the columns of A . In other words, the ij entry of A^T is a_{ji} .

$$\begin{matrix} & A & & A^T \\ \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{pmatrix} & \rightsquigarrow & \begin{pmatrix} a_{11} & a_{21} \\ a_{12} & a_{22} \\ a_{13} & a_{23} \end{pmatrix} \end{matrix}$$

Addition and Scalar Multiplication

You add two matrices component by component, like with vectors.

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{pmatrix} + \begin{pmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \end{pmatrix} = \begin{pmatrix} a_{11} + b_{11} & a_{12} + b_{12} & a_{13} + b_{13} \\ a_{21} + b_{21} & a_{22} + b_{22} & a_{23} + b_{23} \end{pmatrix}$$

Addition and Scalar Multiplication

You add two matrices component by component, like with vectors.

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{pmatrix} + \begin{pmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \end{pmatrix} = \begin{pmatrix} a_{11} + b_{11} & a_{12} + b_{12} & a_{13} + b_{13} \\ a_{21} + b_{21} & a_{22} + b_{22} & a_{23} + b_{23} \end{pmatrix}$$

Note you can only add two matrices *of the same size*.

Addition and Scalar Multiplication

You add two matrices component by component, like with vectors.

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{pmatrix} + \begin{pmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \end{pmatrix} = \begin{pmatrix} a_{11} + b_{11} & a_{12} + b_{12} & a_{13} + b_{13} \\ a_{21} + b_{21} & a_{22} + b_{22} & a_{23} + b_{23} \end{pmatrix}$$

Note you can only add two matrices *of the same size*.

You multiply a matrix by a scalar by multiplying each component, like with vectors.

$$c \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{pmatrix} = \begin{pmatrix} ca_{11} & ca_{12} & ca_{13} \\ ca_{21} & ca_{22} & ca_{23} \end{pmatrix}.$$

Addition and Scalar Multiplication

You add two matrices component by component, like with vectors.

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{pmatrix} + \begin{pmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \end{pmatrix} = \begin{pmatrix} a_{11} + b_{11} & a_{12} + b_{12} & a_{13} + b_{13} \\ a_{21} + b_{21} & a_{22} + b_{22} & a_{23} + b_{23} \end{pmatrix}$$

Note you can only add two matrices *of the same size*.

You multiply a matrix by a scalar by multiplying each component, like with vectors.

$$c \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{pmatrix} = \begin{pmatrix} ca_{11} & ca_{12} & ca_{13} \\ ca_{21} & ca_{22} & ca_{23} \end{pmatrix}.$$

These satisfy the expected rules, like with vectors:

Matrix Multiplication

Beware: matrix multiplication is more subtle than addition and scalar multiplication.

Matrix Multiplication


Beware: matrix multiplication is more subtle than addition and scalar multiplication.

Let A be an $m \times n$ matrix and let B be an $n \times p$ matrix with columns v_1, v_2, \dots, v_p :

$$B = \begin{pmatrix} | & | & & | \\ v_1 & v_2 & \cdots & v_p \\ | & | & & | \end{pmatrix}.$$

The **product** AB is the $m \times p$ matrix with columns Av_1, Av_2, \dots, Av_p :

The equality is
a definition


$$AB \stackrel{\text{def}}{=} \begin{pmatrix} | & | & & | \\ Av_1 & Av_2 & \cdots & Av_p \\ | & | & & | \end{pmatrix}.$$

Matrix Multiplication


Beware: matrix multiplication is more subtle than addition and scalar multiplication.

Let A be an $m \times n$ matrix and let B be an $n \times p$ matrix with columns v_1, v_2, \dots, v_p :

$$B = \begin{pmatrix} | & | & & | \\ v_1 & v_2 & \cdots & v_p \\ | & | & & | \end{pmatrix}.$$

The **product** AB is the $m \times p$ matrix with columns Av_1, Av_2, \dots, Av_p :

The equality is
a definition


$$AB \stackrel{\text{def}}{=} \begin{pmatrix} | & | & & | \\ Av_1 & Av_2 & \cdots & Av_p \\ | & | & & | \end{pmatrix}.$$

In order for Av_1, Av_2, \dots, Av_p to make sense, the number of _____ of A has to be the same as the number of _____ of B .

Matrix Multiplication

Beware: matrix multiplication is more subtle than addition and scalar multiplication.

Let A be an $m \times n$ matrix and let B be an $n \times p$ matrix with columns v_1, v_2, \dots, v_p :

$$B = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_p \\ | & | & \cdots & | \end{pmatrix}.$$

The **product** AB is the $m \times p$ matrix with columns Av_1, Av_2, \dots, Av_p :

The equality is a definition

$$AB \stackrel{\text{def}}{=} \begin{pmatrix} | & | & \cdots & | \\ Av_1 & Av_2 & \cdots & Av_p \\ | & | & \cdots & | \end{pmatrix}.$$

In order for Av_1, Av_2, \dots, Av_p to make sense, the number of **columns** of A has to be the same as the number of **rows** of B .

Matrix Multiplication

Beware: matrix multiplication is more subtle than addition and scalar multiplication.

Let A be an $m \times n$ matrix and let B be an $n \times p$ matrix with columns v_1, v_2, \dots, v_p :

$$B = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_p \\ | & | & \cdots & | \end{pmatrix}.$$

The **product** AB is the $m \times p$ matrix with columns Av_1, Av_2, \dots, Av_p :

The equality is a definition

$$AB \stackrel{\text{def}}{=} \begin{pmatrix} | & | & \cdots & | \\ Av_1 & Av_2 & \cdots & Av_p \\ | & | & \cdots & | \end{pmatrix}.$$

In order for Av_1, Av_2, \dots, Av_p to make sense, the number of **columns** of A has to be the same as the number of **rows** of B .

Example

$$\begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix} \begin{pmatrix} 1 & -3 \\ 2 & -2 \\ 3 & -1 \end{pmatrix} =$$

Composition of Transformations

Why is this the correct definition of matrix multiplication?

Composition of Transformations

Why is this the correct definition of matrix multiplication?

Definition

Let $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$ and $U: \mathbf{R}^p \rightarrow \mathbf{R}^n$ be transformations. The **composition** is the transformation

$$T \circ U: \mathbf{R}^p \rightarrow \mathbf{R}^m \quad \text{defined by} \quad T \circ U(x) = T(U(x)).$$

Composition of Transformations

Why is this the correct definition of matrix multiplication?

Definition

Let $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$ and $U: \mathbf{R}^p \rightarrow \mathbf{R}^n$ be transformations. The **composition** is the transformation

$$T \circ U: \mathbf{R}^p \rightarrow \mathbf{R}^m \quad \text{defined by} \quad T \circ U(x) = T(U(x)).$$

This makes sense because $U(x)$ (the output of U) is in ___, which is the domain of T (the inputs of T).

Composition of Transformations

Why is this the correct definition of matrix multiplication?

Definition

Let $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$ and $U: \mathbf{R}^p \rightarrow \mathbf{R}^n$ be transformations. The **composition** is the transformation

$$T \circ U: \mathbf{R}^p \rightarrow \mathbf{R}^m \quad \text{defined by} \quad T \circ U(x) = T(U(x)).$$

This makes sense because $U(x)$ (the output of U) is in \mathbf{R}^n , which is the domain of T (the inputs of T).

Composition of Transformations


Why is this the correct definition of matrix multiplication?

Definition

Let $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$ and $U: \mathbf{R}^p \rightarrow \mathbf{R}^n$ be transformations. The **composition** is the transformation

$$T \circ U: \mathbf{R}^p \rightarrow \mathbf{R}^m \quad \text{defined by} \quad T \circ U(x) = T(U(x)).$$

This makes sense because $U(x)$ (the output of U) is in \mathbf{R}^n , which is the domain of T (the inputs of T).



Composition of Transformations


Why is this the correct definition of matrix multiplication?

Definition

Let $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$ and $U: \mathbf{R}^p \rightarrow \mathbf{R}^n$ be transformations. The **composition** is the transformation

$$T \circ U: \mathbf{R}^p \rightarrow \mathbf{R}^m \quad \text{defined by} \quad T \circ U(x) = T(U(x)).$$

This makes sense because $U(x)$ (the output of U) is in \mathbf{R}^n , which is the domain of T (the inputs of T).



Fact: If T and U are linear then so is $T \circ U$.

Composition of Transformations


Why is this the correct definition of matrix multiplication?

Definition

Let $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$ and $U: \mathbf{R}^p \rightarrow \mathbf{R}^n$ be transformations. The **composition** is the transformation

$$T \circ U: \mathbf{R}^p \rightarrow \mathbf{R}^m \quad \text{defined by} \quad T \circ U(x) = T(U(x)).$$

This makes sense because $U(x)$ (the output of U) is in \mathbf{R}^n , which is the domain of T (the inputs of T).



Fact: If T and U are linear then so is $T \circ U$.

Guess: If A is the matrix for T , and B is the matrix for U , what is the matrix for $T \circ U$?

Composition of Linear Transformations

Let $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$ and $U: \mathbf{R}^p \rightarrow \mathbf{R}^n$ be *linear* transformations. Let A and B be their matrices:

Composition of Linear Transformations

Let $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$ and $U: \mathbf{R}^p \rightarrow \mathbf{R}^n$ be *linear* transformations. Let A and B be their matrices:

$$A = \left(\begin{array}{c|c|c|c} & & & \\ T(e_1) & T(e_2) & \cdots & T(e_n) \\ & & & \end{array} \right) \quad B = \left(\begin{array}{c|c|c|c} & & & \\ U(e_1) & U(e_2) & \cdots & U(e_p) \\ & & & \end{array} \right)$$

Composition of Linear Transformations

Let $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$ and $U: \mathbf{R}^p \rightarrow \mathbf{R}^n$ be *linear* transformations. Let A and B be their matrices:

$$A = \left(\begin{array}{c|c|c|c} & & & \\ T(e_1) & T(e_2) & \cdots & T(e_n) \\ & & & \end{array} \right) \quad B = \left(\begin{array}{c|c|c|c} & & & \\ U(e_1) & U(e_2) & \cdots & U(e_p) \\ & & & \end{array} \right)$$

Question

What is the matrix for $T \circ U$?

Composition of Linear Transformations

Let $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$ and $U: \mathbf{R}^p \rightarrow \mathbf{R}^n$ be *linear* transformations. Let A and B be their matrices:

$$A = \left(\begin{array}{c|c|c|c} & & & \\ T(e_1) & T(e_2) & \cdots & T(e_n) \\ & & & \end{array} \right) \quad B = \left(\begin{array}{c|c|c|c} & & & \\ U(e_1) & U(e_2) & \cdots & U(e_p) \\ & & & \end{array} \right)$$

Question

What is the matrix for $T \circ U$?

The matrix of the composition is the product of the matrices!

Composition of Linear Transformations

Example

Let $T: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ be rotation by 45° , and let $U: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ be projection onto the x -axis. Let's compute their standard matrices A and B :

Composition of Linear Transformations

Example

Let $T: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ be rotation by 45° , and let $U: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ be projection onto the x -axis. Let's compute their standard matrices A and B :

$$\Rightarrow \quad A = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} \quad B = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$$

Composition of Linear Transformations

Example, continued

So the matrix C for $T \circ U$ is

Composition of Linear Transformations

Example, continued

So the matrix C for $T \circ U$ is

Check:

Composition of Linear Transformations

Example, continued

So the matrix C for $T \circ U$ is

Check:

$$\Rightarrow C = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix} \quad \checkmark$$

Composition of Linear Transformations

Another example

Let

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix} \quad B = \begin{pmatrix} 1 & -3 \\ 2 & -2 \\ 3 & -1 \end{pmatrix}.$$

Let $T(x) = Ax$ and $U(y) = By$, so

$$T: \mathbf{R}^3 \longrightarrow \mathbf{R}^2 \quad U: \mathbf{R}^2 \longrightarrow \mathbf{R}^3 \quad T \circ U: \mathbf{R}^2 \longrightarrow \mathbf{R}^2.$$

Composition of Linear Transformations

Another example

Let

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix} \quad B = \begin{pmatrix} 1 & -3 \\ 2 & -2 \\ 3 & -1 \end{pmatrix}.$$

Let $T(x) = Ax$ and $U(y) = By$, so

$$T: \mathbf{R}^3 \longrightarrow \mathbf{R}^2 \quad U: \mathbf{R}^- \longrightarrow \mathbf{R}^- \quad T \circ U: \mathbf{R}^- \longrightarrow \mathbf{R}^-.$$

Composition of Linear Transformations

Another example

Let

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix} \quad B = \begin{pmatrix} 1 & -3 \\ 2 & -2 \\ 3 & -1 \end{pmatrix}.$$

Let $T(x) = Ax$ and $U(y) = By$, so

$$T: \mathbf{R}^3 \longrightarrow \mathbf{R}^2 \quad U: \mathbf{R}^2 \longrightarrow \mathbf{R}^3 \quad T \circ U: \mathbf{R}^2 \longrightarrow \mathbf{R}^2.$$

Composition of Linear Transformations

Another example

Let

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix} \quad B = \begin{pmatrix} 1 & -3 \\ 2 & -2 \\ 3 & -1 \end{pmatrix}.$$

Let $T(x) = Ax$ and $U(y) = By$, so

$$T: \mathbf{R}^3 \longrightarrow \mathbf{R}^2 \quad U: \mathbf{R}^2 \longrightarrow \mathbf{R}^3 \quad T \circ U: \mathbf{R}^2 \longrightarrow \mathbf{R}^2.$$

Composition of Linear Transformations

Another example

Let

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix} \quad B = \begin{pmatrix} 1 & -3 \\ 2 & -2 \\ 3 & -1 \end{pmatrix}.$$

Let $T(x) = Ax$ and $U(y) = By$, so

$$T: \mathbf{R}^3 \longrightarrow \mathbf{R}^2 \quad U: \mathbf{R}^2 \longrightarrow \mathbf{R}^3 \quad T \circ U: \mathbf{R}^2 \longrightarrow \mathbf{R}^2.$$

Let's find the matrix for $T \circ U$:

$$T \circ U(e_1) =$$

$$T \circ U(e_2) =$$

Composition of Linear Transformations

Another example

Let

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix} \quad B = \begin{pmatrix} 1 & -3 \\ 2 & -2 \\ 3 & -1 \end{pmatrix}.$$

Let $T(x) = Ax$ and $U(y) = By$, so

$$T: \mathbf{R}^3 \longrightarrow \mathbf{R}^2 \quad U: \mathbf{R}^2 \longrightarrow \mathbf{R}^3 \quad T \circ U: \mathbf{R}^2 \longrightarrow \mathbf{R}^2.$$

Let's find the matrix for $T \circ U$:

$$T \circ U(e_1) =$$

$$T \circ U(e_2) =$$

Before we computed $AB = \begin{pmatrix} 14 & -10 \\ 32 & -28 \end{pmatrix}$, so AB is the matrix of $T \circ U$.

Poll

Do there exist *nonzero* matrices A and B with $AB = 0$?

Poll

Do there exist *nonzero* matrices A and B with $AB = 0$?

Yes! Here's an example:

$$\begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 1 & 1 \end{pmatrix} = \left(\begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right) = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}.$$

The Row-Column Rule for Matrix Multiplication

Recall: A row vector of length n times a column vector of length n is a scalar:

$$\begin{pmatrix} a_1 & \cdots & a_n \end{pmatrix} \begin{pmatrix} b_1 \\ \vdots \\ b_n \end{pmatrix} = a_1 b_1 + \cdots + a_n b_n.$$

The Row-Column Rule for Matrix Multiplication

Recall: A row vector of length n times a column vector of length n is a scalar:

$$(a_1 \quad \cdots \quad a_n) \begin{pmatrix} b_1 \\ \vdots \\ b_n \end{pmatrix} = a_1 b_1 + \cdots + a_n b_n.$$

Another way of multiplying a matrix by a vector is:

$$Ax = \begin{pmatrix} -r_1- \\ \vdots \\ -r_m- \end{pmatrix} x = \begin{pmatrix} r_1 x \\ \vdots \\ r_m x \end{pmatrix}.$$

The Row-Column Rule for Matrix Multiplication

Recall: A row vector of length n times a column vector of length n is a scalar:

$$(a_1 \quad \cdots \quad a_n) \begin{pmatrix} b_1 \\ \vdots \\ b_n \end{pmatrix} = a_1 b_1 + \cdots + a_n b_n.$$

Another way of multiplying a matrix by a vector is:

$$Ax = \begin{pmatrix} -r_1- \\ \vdots \\ -r_m- \end{pmatrix} x = \begin{pmatrix} r_1 x \\ \vdots \\ r_m x \end{pmatrix}.$$

On the other hand, you multiply two matrices by

$$AB = A \begin{pmatrix} | & & | \\ c_1 & \cdots & c_p \\ | & & | \end{pmatrix} = \begin{pmatrix} | & & | \\ Ac_1 & \cdots & Ac_p \\ | & & | \end{pmatrix}.$$

The Row-Column Rule for Matrix Multiplication

Recall: A row vector of length n times a column vector of length n is a scalar:

$$(a_1 \quad \cdots \quad a_n) \begin{pmatrix} b_1 \\ \vdots \\ b_n \end{pmatrix} = a_1 b_1 + \cdots + a_n b_n.$$

Another way of multiplying a matrix by a vector is:

$$Ax = \begin{pmatrix} -r_1- \\ \vdots \\ -r_m- \end{pmatrix} x = \begin{pmatrix} r_1 x \\ \vdots \\ r_m x \end{pmatrix}.$$

On the other hand, you multiply two matrices by

$$AB = A \begin{pmatrix} | & & | \\ c_1 & \cdots & c_p \\ | & & | \end{pmatrix} = \begin{pmatrix} | & & | \\ Ac_1 & \cdots & Ac_p \\ | & & | \end{pmatrix}.$$

It follows that

$$AB = \begin{pmatrix} -r_1- \\ \vdots \\ -r_m- \end{pmatrix} \begin{pmatrix} | & & | \\ c_1 & \cdots & c_p \\ | & & | \end{pmatrix} = \begin{pmatrix} r_1 c_1 & r_1 c_2 & \cdots & r_1 c_p \\ r_2 c_1 & r_2 c_2 & \cdots & r_2 c_p \\ \vdots & \vdots & & \vdots \\ r_m c_1 & r_m c_2 & \cdots & r_m c_p \end{pmatrix}$$

The Row-Column Rule for Matrix Multiplication

The ij entry of $C = AB$ is the i th row of A times the j th column of B :

$$c_{ij} = (AB)_{ij} = a_{i1}b_{1j} + a_{i2}b_{2j} + \cdots + a_{in}b_{nj}.$$

The Row-Column Rule for Matrix Multiplication

The ij entry of $C = AB$ is the i th row of A times the j th column of B :

$$c_{ij} = (AB)_{ij} = a_{i1}b_{1j} + a_{i2}b_{2j} + \cdots + a_{in}b_{nj}.$$

This is how everybody on the planet actually computes AB .

The Row-Column Rule for Matrix Multiplication

The ij entry of $C = AB$ is the i th row of A times the j th column of B :

$$c_{ij} = (AB)_{ij} = a_{i1}b_{1j} + a_{i2}b_{2j} + \cdots + a_{in}b_{nj}.$$

This is how everybody on the planet actually computes AB . Diagram ($AB = C$):

$$\begin{pmatrix} a_{11} & \cdots & a_{1k} & \cdots & a_{1n} \\ \vdots & & \vdots & & \vdots \\ a_{i1} & \cdots & a_{ik} & \cdots & a_{in} \\ \vdots & & \vdots & & \vdots \\ a_{m1} & \cdots & a_{mk} & \cdots & a_{mn} \end{pmatrix} \cdot \begin{pmatrix} b_{11} & \cdots & b_{1j} & \cdots & b_{1p} \\ \vdots & & \vdots & & \vdots \\ b_{k1} & \cdots & b_{kj} & \cdots & b_{kp} \\ \vdots & & \vdots & & \vdots \\ b_{n1} & \cdots & b_{nj} & \cdots & b_{np} \end{pmatrix} = \begin{pmatrix} c_{11} & \cdots & c_{1j} & \cdots & c_{1p} \\ \vdots & & \vdots & & \vdots \\ c_{i1} & \cdots & c_{ij} & \cdots & c_{ip} \\ \vdots & & \vdots & & \vdots \\ c_{m1} & \cdots & c_{mj} & \cdots & c_{mp} \end{pmatrix}$$

j th column i th row i th row j th column ij entry

The Row-Column Rule for Matrix Multiplication

The ij entry of $C = AB$ is the i th row of A times the j th column of B :

$$c_{ij} = (AB)_{ij} = a_{i1}b_{1j} + a_{i2}b_{2j} + \cdots + a_{in}b_{nj}.$$

This is how everybody on the planet actually computes AB . Diagram ($AB = C$):

$$\begin{pmatrix} a_{11} & \cdots & a_{1k} & \cdots & a_{1n} \\ \vdots & & \vdots & & \vdots \\ a_{i1} & \cdots & a_{ik} & \cdots & a_{in} \\ \vdots & & \vdots & & \vdots \\ a_{m1} & \cdots & a_{mk} & \cdots & a_{mn} \end{pmatrix} \cdot \begin{pmatrix} b_{11} & \cdots & b_{1j} & \cdots & b_{1p} \\ \vdots & & \vdots & & \vdots \\ b_{k1} & \cdots & b_{kj} & \cdots & b_{kp} \\ \vdots & & \vdots & & \vdots \\ b_{n1} & \cdots & b_{nj} & \cdots & b_{np} \end{pmatrix} = \begin{pmatrix} c_{11} & \cdots & c_{1j} & \cdots & c_{1p} \\ \vdots & & \vdots & & \vdots \\ c_{i1} & \cdots & c_{ij} & \cdots & c_{ip} \\ \vdots & & \vdots & & \vdots \\ c_{m1} & \cdots & c_{mj} & \cdots & c_{mp} \end{pmatrix}$$

j th column ij entry

Example

Properties of Matrix Multiplication

Mostly matrix multiplication works like you'd expect. Suppose A has size $m \times n$, and that the other matrices below have the right size to make multiplication work.

Properties of Matrix Multiplication

Mostly matrix multiplication works like you'd expect. Suppose A has size $m \times n$, and that the other matrices below have the right size to make multiplication work.

Most of these are easy to verify.

Properties of Matrix Multiplication

Mostly matrix multiplication works like you'd expect. Suppose A has size $m \times n$, and that the other matrices below have the right size to make multiplication work.

Most of these are easy to verify.

Associativity is $A(BC) = (AB)C$. It is a pain to verify using the row-column rule!

Properties of Matrix Multiplication

Mostly matrix multiplication works like you'd expect. Suppose A has size $m \times n$, and that the other matrices below have the right size to make multiplication work.

Most of these are easy to verify.

Associativity is $A(BC) = (AB)C$. It is a pain to verify using the row-column rule! Much easier: use associativity of linear transformations:

$$S \circ (T \circ U) = (S \circ T) \circ U.$$

Properties of Matrix Multiplication

Mostly matrix multiplication works like you'd expect. Suppose A has size $m \times n$, and that the other matrices below have the right size to make multiplication work.

Most of these are easy to verify.

Associativity is $A(BC) = (AB)C$. It is a pain to verify using the row-column rule! Much easier: use associativity of linear transformations:

$$S \circ (T \circ U) = (S \circ T) \circ U.$$

This is a good example of an instance where having a conceptual viewpoint saves you a lot of work.

Properties of Matrix Multiplication

Mostly matrix multiplication works like you'd expect. Suppose A has size $m \times n$, and that the other matrices below have the right size to make multiplication work.

Most of these are easy to verify.

Associativity is $A(BC) = (AB)C$. It is a pain to verify using the row-column rule! Much easier: use associativity of linear transformations:

$$S \circ (T \circ U) = (S \circ T) \circ U.$$

This is a good example of an instance where having a conceptual viewpoint saves you a lot of work.

Recommended: Try to verify all of them on your own.

Properties of Matrix Multiplication

Caveats

Warnings!

- ▶ AB is usually not equal to BA .

Properties of Matrix Multiplication

Caveats

Warnings!

- ▶ AB is usually not equal to BA .

In fact, AB may be defined when BA is not.

Properties of Matrix Multiplication

Caveats

Warnings!

- ▶ AB is usually not equal to BA .

In fact, AB may be defined when BA is not.

- ▶ $AB = AC$ does not imply $B = C$, even if $A \neq 0$.

Properties of Matrix Multiplication

Caveats

Warnings!

- ▶ AB is usually not equal to BA .

In fact, AB may be defined when BA is not.

- ▶ $AB = AC$ does not imply $B = C$, even if $A \neq 0$.
- ▶ $AB = 0$ does not imply $A = 0$ or $B = 0$.

Read about powers of a matrix and multiplication of transposes in §2.1.