


Section 6.3

Orthogonal Projections

Idea Behind Orthogonal Projections

If x is not in a subspace W , then y in W is the closest to x if $x - y$ is in W^\perp :




Reformulation: Every vector x can be decomposed uniquely as

$$x = x_W + x_{W^\perp}$$

where $x_W = y$ is the closest vector to x in W , and $x_{W^\perp} = x - y$ is in W^\perp .

Example: Let $u = \begin{pmatrix} 3 \\ 2 \end{pmatrix}$ and let $L = \text{Span}\{u\}$. Let $x = \begin{pmatrix} -6 \\ 4 \end{pmatrix}$. Then the closest point to x in L is $\text{proj}_L(x) = \frac{x \cdot u}{u \cdot u} u$, so

$$x_L = \text{proj}_L(x) = -\frac{10}{13} \begin{pmatrix} 3 \\ 2 \end{pmatrix} \quad x_{L^\perp} = x - \text{proj}_L(x) = \begin{pmatrix} -6 \\ 4 \end{pmatrix} + \frac{10}{13} \begin{pmatrix} 3 \\ 2 \end{pmatrix}.$$



Orthogonal Projections

Definition

Let W be a subspace of \mathbf{R}^n , and let $\{u_1, u_2, \dots, u_m\}$ be an *orthogonal* basis for W . The **orthogonal projection** of a vector x onto W is

$$\text{proj}_W(x) \stackrel{\text{def}}{=} \sum_{i=1}^m \frac{x \cdot u_i}{u_i \cdot u_i} u_i.$$

Question: What is the difference between this and the formula for $[x]_B$ from before?

Theorem

Let W be a subspace of \mathbf{R}^n , and let x be a vector in \mathbf{R}^n . Then $\text{proj}_W(x)$ is the closest point to x in W . Therefore

$$x_W = \text{proj}_W(x) \quad x_{W^\perp} = x - \text{proj}_W(x).$$

Why? Let $y = \text{proj}_W(x)$. We need to show that $x - y$ is in W^\perp . In other words, $u_i \cdot (x - y) = 0$ for each i . Let's do u_1 :

$$u_1 \cdot (x - y) = u_1 \cdot \left(x - \sum_{i=1}^m \frac{x \cdot u_i}{u_i \cdot u_i} u_i \right) = u_1 \cdot x - \frac{x \cdot u_1}{u_1 \cdot u_1} (u_1 \cdot u_1) - 0 - \dots = 0.$$

Orthogonal Projections


Easy example

What is the projection of $x = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$ onto the xy -plane?

Answer: The xy -plane is $W = \text{Span}\{e_1, e_2\}$, and $\{e_1, e_2\}$ is an orthogonal basis.

$$x_W = \text{proj}_W \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} = \frac{x \cdot e_1}{e_1 \cdot e_1} e_1 + \frac{x \cdot e_2}{e_2 \cdot e_2} e_2 = \frac{1 \cdot 1}{1^2} e_1 + \frac{1 \cdot 2}{1^2} e_2 = \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix}.$$

So this is the same projection as before.



Orthogonal Projections


More complicated example

What is the projection of $x = \begin{pmatrix} -1.1 \\ 1.4 \\ 1.45 \end{pmatrix}$ onto $W = \text{Span} \left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1.1 \\ -0.2 \end{pmatrix} \right\}$?

Answer: The basis is orthogonal, so

$$\begin{aligned} x_W &= \text{proj}_W \begin{pmatrix} -1.1 \\ 1.4 \\ 1.45 \end{pmatrix} = \frac{x \cdot u_1}{u_1 \cdot u_1} u_1 + \frac{x \cdot u_2}{u_2 \cdot u_2} u_2 \\ &= \frac{(-1.1)(1)}{1^2} \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} + \frac{(1.4)(1.1) + (1.45)(-0.2)}{1.1^2 + (-0.2)^2} \begin{pmatrix} 0 \\ 1.1 \\ -0.2 \end{pmatrix} \end{aligned}$$

This turns out to be equal to $u_2 - 1.1u_1$.




Orthogonal Projections

Picture

Let W be a subspace of \mathbf{R}^n , and let $\{u_1, u_2, \dots, u_m\}$ be an orthogonal basis for W . Let $L_i = \text{Span}\{u_i\}$. Then

$$\text{proj}_W(x) = \sum_{i=1}^m \frac{x \cdot u_i}{u_i \cdot u_i} u_i = \sum_{i=1}^m \text{proj}_{L_i}(x).$$

So the orthogonal projection is formed by adding orthogonal projections onto perpendicular lines.



Orthogonal Projections

Properties

First we restate the property we've been using all along.

Best Approximation Theorem

Let W be a subspace of \mathbf{R}^n , and let x be a vector in \mathbf{R}^n . Then $y = \text{proj}_W(x)$ is the closest point in W to x , in the sense that

$$\text{dist}(x, y') \geq \text{dist}(x, y) \quad \text{for all } y' \text{ in } W.$$

We can think of orthogonal projection as a *transformation*:

$$\text{proj}_W: \mathbf{R}^n \longrightarrow \mathbf{R}^n \quad x \mapsto \text{proj}_W(x).$$

Theorem

Let W be a subspace of \mathbf{R}^n .

1. proj_W is a *linear* transformation.
2. For every x in W , we have $\text{proj}_W(x) = x$.
3. For every x in W^\perp , we have $\text{proj}_W(x) = 0$.
4. The range of proj_W is W .

Let W be a subspace of \mathbf{R}^n .

Poll

Let A be the matrix for proj_W . What is/are the eigenvalue(s) of A ?

A. 0 B. 1 C. -1 D. 0, 1 E. 1, -1 F. 0, -1 G. -1 , 0, 1

The 1-eigenspace is W .

The 0-eigenspace is W^\perp .

We have $\dim W + \dim W^\perp = n$, so that gives n linearly independent eigenvectors already.

So the answer is D.

Orthogonal Projections

Matrices

What is the matrix for $\text{proj}_W: \mathbf{R}^3 \rightarrow \mathbf{R}^3$, where

$$W = \text{Span} \left\{ \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right\}?$$

Answer: Recall how to compute the matrix for a linear transformation:

$$A = \begin{pmatrix} \left. \text{proj}_W(e_1) \right| & \left. \text{proj}_W(e_2) \right| & \left. \text{proj}_W(e_3) \right| \\ \left| \right. & \left| \right. & \left| \right. \end{pmatrix}.$$

We compute:

$$\text{proj}_W(e_1) = \frac{e_1 \cdot u_1}{u_1 \cdot u_1} u_1 + \frac{e_1 \cdot u_2}{u_2 \cdot u_2} u_2 = \frac{1}{2} \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} + \frac{1}{3} \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 5/6 \\ 1/3 \\ -1/6 \end{pmatrix}$$

$$\text{proj}_W(e_2) = \frac{e_2 \cdot u_1}{u_1 \cdot u_1} u_1 + \frac{e_2 \cdot u_2}{u_2 \cdot u_2} u_2 = 0 + \frac{1}{3} \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 1/3 \\ 1/3 \\ 1/3 \end{pmatrix}$$

$$\text{proj}_W(e_3) = \frac{e_3 \cdot u_1}{u_1 \cdot u_1} u_1 + \frac{e_3 \cdot u_2}{u_2 \cdot u_2} u_2 = -\frac{1}{2} \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} + \frac{1}{3} \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} -1/6 \\ 1/3 \\ 5/6 \end{pmatrix}$$

$$\text{Therefore } A = \begin{pmatrix} 5/6 & 1/3 & -1/6 \\ 1/3 & 1/3 & 1/3 \\ -1/6 & 1/3 & 5/6 \end{pmatrix}.$$

Orthogonal Projections

Matrix facts

Let W be an m -dimensional subspace of \mathbf{R}^n , let $\text{proj}_W: \mathbf{R}^n \rightarrow W$ be the projection, and let A be the matrix for proj_L .

Fact 1: A is diagonalizable with eigenvalues 0 and 1; it is similar to the diagonal matrix with m ones and $n - m$ zeros on the diagonal.

Why? Let v_1, v_2, \dots, v_m be a basis for W , and let $v_{m+1}, v_{m+2}, \dots, v_n$ be a basis for W^\perp . These are (linearly independent) eigenvectors with eigenvalues 1 and 0, respectively, and they form a basis for \mathbf{R}^n because there are n of them.

Example: If W is a plane in \mathbf{R}^3 , then A is similar to projection onto the xy -plane:

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

Fact 2: $A^2 = A$.

Why? Projecting twice is the same as projecting once:

$$\text{proj}_W \circ \text{proj}_W = \text{proj}_W \implies A \cdot A = A.$$

Orthogonal Projections

Minimum distance

What is the distance from e_1 to $W = \text{Span} \left\{ \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right\}$?

Answer: The closest point on W to e_1 is $\text{proj}_W(e_1) = \begin{pmatrix} 5/6 \\ 1/3 \\ -1/6 \end{pmatrix}$.

The distance from e_1 to this point is

$$\begin{aligned} \text{dist}(e_1, \text{proj}_W(e_1)) &= \|(e_1)_{W^\perp}\| \\ &= \left\| \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} - \begin{pmatrix} 5/6 \\ 1/3 \\ -1/6 \end{pmatrix} \right\| \\ &= \left\| \begin{pmatrix} 1/6 \\ -1/3 \\ -1/6 \end{pmatrix} \right\| \\ &= \sqrt{(1/6)^2 + (-1/3)^2 + (-1/6)^2} \\ &= \frac{1}{\sqrt{6}}. \end{aligned}$$

