

# Chapter 2

## Matrix Algebra

# Section 2.1

## Matrix Operations

## Motivation

Recall: we can turn any system of linear equations into a matrix equation

$$Ax = b.$$

# Motivation

Recall: we can turn any system of linear equations into a matrix equation

$$Ax = b.$$

This notation is suggestive. Can we solve the equation by “dividing by A”?

$$x \stackrel{??}{=} \frac{b}{A}$$

# Motivation

**Recall:** we can turn any system of linear equations into a matrix equation

$$Ax = b.$$

This notation is suggestive. Can we solve the equation by “dividing by A”?

$$x \stackrel{??}{=} \frac{b}{A}$$

**Answer:** Sometimes, but you have to know what you're doing.

# Motivation

**Recall:** we can turn any system of linear equations into a matrix equation

$$Ax = b.$$

This notation is suggestive. Can we solve the equation by “dividing by A”?

$$x \stackrel{??}{=} \frac{b}{A}$$

**Answer:** Sometimes, but you have to know what you're doing.

Today we'll study *matrix algebra*: adding and multiplying matrices.

## More Notation for Matrices

Let  $A$  be an  $m \times n$  matrix.

## More Notation for Matrices

Let  $A$  be an  $m \times n$  matrix.

We write  $a_{ij}$  for the entry in the  $i$ th row and the  $j$ th column. It is called the  **$ij$ th entry** of the matrix.

$$\begin{pmatrix} a_{11} & \cdots & a_{1j} & \cdots & a_{1n} \\ \vdots & & \vdots & & \vdots \\ a_{i1} & \cdots & a_{ij} & \cdots & a_{in} \\ \vdots & & \vdots & & \vdots \\ a_{m1} & \cdots & a_{mj} & \cdots & a_{mn} \end{pmatrix}$$

$j$ th column

$i$ th row



## More Notation for Matrices

Let  $A$  be an  $m \times n$  matrix.

We write  $a_{ij}$  for the entry in the  $i$ th row and the  $j$ th column. It is called the  **$ij$ th entry** of the matrix.

$$\begin{pmatrix} a_{11} & \cdots & a_{1j} & \cdots & a_{1n} \\ \vdots & & \vdots & & \vdots \\ a_{i1} & \cdots & a_{ij} & \cdots & a_{in} \\ \vdots & & \vdots & & \vdots \\ a_{m1} & \cdots & a_{mj} & \cdots & a_{mn} \end{pmatrix}$$

$j$ th column

The entries  $a_{11}, a_{22}, a_{33}, \dots$  are the **diagonal entries**; they form the **main diagonal** of the matrix.

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{pmatrix} \quad \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \\ a_{31} & a_{32} \end{pmatrix}$$

## More Notation for Matrices

Let  $A$  be an  $m \times n$  matrix.

We write  $a_{ij}$  for the entry in the  $i$ th row and the  $j$ th column. It is called the  **$ij$ th entry** of the matrix.

$$\begin{pmatrix} a_{11} & \cdots & a_{1j} & \cdots & a_{1n} \\ \vdots & & \vdots & & \vdots \\ a_{i1} & \cdots & a_{ij} & \cdots & a_{in} \\ \vdots & & \vdots & & \vdots \\ a_{m1} & \cdots & a_{mj} & \cdots & a_{mn} \end{pmatrix}$$

$j$ th column

$i$ th row

The entries  $a_{11}, a_{22}, a_{33}, \dots$  are the **diagonal entries**; they form the **main diagonal** of the matrix.

A **diagonal matrix** is a *square* matrix whose only nonzero entries are on the main diagonal.

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \quad \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \\ a_{31} & a_{32} \end{pmatrix}$$

$$\begin{pmatrix} a_{11} & 0 & 0 \\ 0 & a_{22} & 0 \\ 0 & 0 & a_{33} \end{pmatrix}$$

## More Notation for Matrices

Let  $A$  be an  $m \times n$  matrix.

We write  $a_{ij}$  for the entry in the  $i$ th row and the  $j$ th column. It is called the  **$ij$ th entry** of the matrix.

$$\begin{pmatrix} a_{11} & \cdots & a_{1j} & \cdots & a_{1n} \\ \vdots & & \vdots & & \vdots \\ a_{i1} & \cdots & a_{ij} & \cdots & a_{in} \\ \vdots & & \vdots & & \vdots \\ a_{m1} & \cdots & a_{mj} & \cdots & a_{mn} \end{pmatrix}$$

The entries  $a_{11}, a_{22}, a_{33}, \dots$  are the **diagonal entries**; they form the **main diagonal** of the matrix.

A **diagonal matrix** is a *square* matrix whose only nonzero entries are on the main diagonal.

The  $n \times n$  **identity matrix**  $I_n$  is the diagonal matrix with all diagonal entries equal to 1. It is special because  $I_n \mathbf{v} = \mathbf{v}$  for all  $\mathbf{v}$  in  $\mathbf{R}^n$ .

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \quad \begin{pmatrix} a_{11} \\ a_{22} \\ a_{33} \end{pmatrix}$$

$$\begin{pmatrix} a_{11} & 0 & 0 \\ 0 & a_{22} & 0 \\ 0 & 0 & a_{33} \end{pmatrix}$$

$$I_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

## More Notation for Matrices

Let  $A$  be an  $m \times n$  matrix.

We write  $a_{ij}$  for the entry in the  $i$ th row and the  $j$ th column. It is called the  **$ij$ th entry** of the matrix.

$$\begin{pmatrix} a_{11} & \cdots & a_{1j} & \cdots & a_{1n} \\ \vdots & & \vdots & & \vdots \\ a_{i1} & \cdots & a_{ij} & \cdots & a_{in} \\ \vdots & & \vdots & & \vdots \\ a_{m1} & \cdots & a_{mj} & \cdots & a_{mn} \end{pmatrix}$$

$j$ th column

$i$ th row

The entries  $a_{11}, a_{22}, a_{33}, \dots$  are the **diagonal entries**; they form the **main diagonal** of the matrix.

A **diagonal matrix** is a *square* matrix whose only nonzero entries are on the main diagonal.

The  $n \times n$  **identity matrix**  $I_n$  is the diagonal matrix with all diagonal entries equal to 1. It is special because  $I_n \mathbf{v} = \mathbf{v}$  for all  $\mathbf{v}$  in  $\mathbf{R}^n$ .

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{pmatrix} \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \\ a_{31} & a_{32} \end{pmatrix}$$

$$\begin{pmatrix} a_{11} & 0 & 0 \\ 0 & a_{22} & 0 \\ 0 & 0 & a_{33} \end{pmatrix}$$

$$I_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

# More Notation for Matrices

Continued

The **zero matrix** (of size  $m \times n$ ) is the  $m \times n$  matrix  $0$  with all zero entries.

$$0 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

# More Notation for Matrices

## Continued

The **zero matrix** (of size  $m \times n$ ) is the  $m \times n$  matrix  $0$  with all zero entries.

$$0 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

The **transpose** of an  $m \times n$  matrix  $A$  is the  $n \times m$  matrix  $A^T$  whose rows are the columns of  $A$ . In other words, the  $ij$  entry of  $A^T$  is  $a_{ji}$ .

$$\begin{matrix} & A & & A^T \\ \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{pmatrix} & \rightsquigarrow & \begin{pmatrix} a_{11} & a_{21} \\ a_{12} & a_{22} \\ a_{13} & a_{23} \end{pmatrix} \end{matrix}$$

## Addition and Scalar Multiplication

You add two matrices component by component, like with vectors.

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{pmatrix} + \begin{pmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \end{pmatrix} = \begin{pmatrix} a_{11} + b_{11} & a_{12} + b_{12} & a_{13} + b_{13} \\ a_{21} + b_{21} & a_{22} + b_{22} & a_{23} + b_{23} \end{pmatrix}$$

## Addition and Scalar Multiplication

You add two matrices component by component, like with vectors.

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{pmatrix} + \begin{pmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \end{pmatrix} = \begin{pmatrix} a_{11} + b_{11} & a_{12} + b_{12} & a_{13} + b_{13} \\ a_{21} + b_{21} & a_{22} + b_{22} & a_{23} + b_{23} \end{pmatrix}$$

Note you can only add two matrices *of the same size*.



## Addition and Scalar Multiplication

You add two matrices component by component, like with vectors.

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{pmatrix} + \begin{pmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \end{pmatrix} = \begin{pmatrix} a_{11} + b_{11} & a_{12} + b_{12} & a_{13} + b_{13} \\ a_{21} + b_{21} & a_{22} + b_{22} & a_{23} + b_{23} \end{pmatrix}$$

Note you can only add two matrices *of the same size*.

You multiply a matrix by a scalar by multiplying each component, like with vectors.

$$c \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{pmatrix} = \begin{pmatrix} ca_{11} & ca_{12} & ca_{13} \\ ca_{21} & ca_{22} & ca_{23} \end{pmatrix}.$$

## Addition and Scalar Multiplication

You add two matrices component by component, like with vectors.

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{pmatrix} + \begin{pmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \end{pmatrix} = \begin{pmatrix} a_{11} + b_{11} & a_{12} + b_{12} & a_{13} + b_{13} \\ a_{21} + b_{21} & a_{22} + b_{22} & a_{23} + b_{23} \end{pmatrix}$$

Note you can only add two matrices *of the same size*.

You multiply a matrix by a scalar by multiplying each component, like with vectors.

$$c \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{pmatrix} = \begin{pmatrix} ca_{11} & ca_{12} & ca_{13} \\ ca_{21} & ca_{22} & ca_{23} \end{pmatrix}.$$

These satisfy the expected rules, like with vectors:

# Matrix Multiplication

**Beware:** matrix multiplication is more subtle than addition and scalar multiplication.

# Matrix Multiplication


**Beware:** matrix multiplication is more subtle than addition and scalar multiplication.

Let  $A$  be an  $m \times n$  matrix and let  $B$  be an  $n \times p$  matrix with columns  $v_1, v_2, \dots, v_p$ :

$$B = \begin{pmatrix} | & | & & | \\ v_1 & v_2 & \cdots & v_p \\ | & | & & | \end{pmatrix}.$$

The **product**  $AB$  is the  $m \times p$  matrix with columns  $Av_1, Av_2, \dots, Av_p$ :

The equality is  
a definition


$$AB \stackrel{\text{def}}{=} \begin{pmatrix} | & | & & | \\ Av_1 & Av_2 & \cdots & Av_p \\ | & | & & | \end{pmatrix}.$$

# Matrix Multiplication

**Beware:** matrix multiplication is more subtle than addition and scalar multiplication.

Let  $A$  be an  $m \times n$  matrix and let  $B$  be an  $n \times p$  matrix with columns  $v_1, v_2, \dots, v_p$ :

$$B = \begin{pmatrix} | & | & & | \\ v_1 & v_2 & \cdots & v_p \\ | & | & & | \end{pmatrix}.$$

The **product**  $AB$  is the  $m \times p$  matrix with columns  $Av_1, Av_2, \dots, Av_p$ :

The equality is  
a definition

$$AB \stackrel{\text{def}}{=} \begin{pmatrix} | & | & & | \\ Av_1 & Av_2 & \cdots & Av_p \\ | & | & & | \end{pmatrix}.$$

In order for  $Av_1, Av_2, \dots, Av_p$  to make sense, the number of \_\_\_\_\_ of  $A$  has to be the same as the number of \_\_\_\_\_ of  $B$ .

# Matrix Multiplication

**Beware:** matrix multiplication is more subtle than addition and scalar multiplication.

Let  $A$  be an  $m \times n$  matrix and let  $B$  be an  $n \times p$  matrix with columns  $v_1, v_2, \dots, v_p$ :

$$B = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_p \\ | & | & \cdots & | \end{pmatrix}.$$

The **product**  $AB$  is the  $m \times p$  matrix with columns  $Av_1, Av_2, \dots, Av_p$ :

The equality is a definition

$$AB \stackrel{\text{def}}{=} \begin{pmatrix} | & | & \cdots & | \\ Av_1 & Av_2 & \cdots & Av_p \\ | & | & \cdots & | \end{pmatrix}.$$

In order for  $Av_1, Av_2, \dots, Av_p$  to make sense, the number of **columns** of  $A$  has to be the same as the number of **rows** of  $B$ .

# Matrix Multiplication

**Beware:** matrix multiplication is more subtle than addition and scalar multiplication.

Let  $A$  be an  $m \times n$  matrix and let  $B$  be an  $n \times p$  matrix with columns  $v_1, v_2, \dots, v_p$ :

$$B = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_p \\ | & | & \cdots & | \end{pmatrix}.$$

The **product**  $AB$  is the  $m \times p$  matrix with columns  $Av_1, Av_2, \dots, Av_p$ :

The equality is a definition

$$AB \stackrel{\text{def}}{=} \begin{pmatrix} | & | & \cdots & | \\ Av_1 & Av_2 & \cdots & Av_p \\ | & | & \cdots & | \end{pmatrix}.$$

In order for  $Av_1, Av_2, \dots, Av_p$  to make sense, the number of **columns** of  $A$  has to be the same as the number of **rows** of  $B$ .

**Example**

$$\begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix} \begin{pmatrix} 1 & -3 \\ 2 & -2 \\ 3 & -1 \end{pmatrix} =$$

## Composition of Transformations

Why is this the correct definition of matrix multiplication?



# Composition of Transformations

Why is this the correct definition of matrix multiplication?

## Definition

Let  $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$  and  $U: \mathbf{R}^p \rightarrow \mathbf{R}^n$  be transformations. The **composition** is the transformation

$$T \circ U: \mathbf{R}^p \rightarrow \mathbf{R}^m \quad \text{defined by} \quad T \circ U(x) = T(U(x)).$$

# Composition of Transformations

Why is this the correct definition of matrix multiplication?

## Definition

Let  $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$  and  $U: \mathbf{R}^p \rightarrow \mathbf{R}^n$  be transformations. The **composition** is the transformation

$$T \circ U: \mathbf{R}^p \rightarrow \mathbf{R}^m \quad \text{defined by} \quad T \circ U(x) = T(U(x)).$$

This makes sense because  $U(x)$  (the output of  $U$ ) is in \_\_, which is the domain of  $T$  (the inputs of  $T$ ).

# Composition of Transformations

Why is this the correct definition of matrix multiplication?

## Definition

Let  $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$  and  $U: \mathbf{R}^p \rightarrow \mathbf{R}^n$  be transformations. The **composition** is the transformation

$$T \circ U: \mathbf{R}^p \rightarrow \mathbf{R}^m \quad \text{defined by} \quad T \circ U(x) = T(U(x)).$$

This makes sense because  $U(x)$  (the output of  $U$ ) is in  $\mathbf{R}^n$ , which is the domain of  $T$  (the inputs of  $T$ ).

# Composition of Transformations


Why is this the correct definition of matrix multiplication?

## Definition

Let  $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$  and  $U: \mathbf{R}^p \rightarrow \mathbf{R}^n$  be transformations. The **composition** is the transformation

$$T \circ U: \mathbf{R}^p \rightarrow \mathbf{R}^m \quad \text{defined by} \quad T \circ U(x) = T(U(x)).$$

This makes sense because  $U(x)$  (the output of  $U$ ) is in  $\mathbf{R}^n$ , which is the domain of  $T$  (the inputs of  $T$ ).



# Composition of Transformations


Why is this the correct definition of matrix multiplication?

## Definition

Let  $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$  and  $U: \mathbf{R}^p \rightarrow \mathbf{R}^n$  be transformations. The **composition** is the transformation

$$T \circ U: \mathbf{R}^p \rightarrow \mathbf{R}^m \quad \text{defined by} \quad T \circ U(x) = T(U(x)).$$

This makes sense because  $U(x)$  (the output of  $U$ ) is in  $\mathbf{R}^n$ , which is the domain of  $T$  (the inputs of  $T$ ).



**Fact:** If  $T$  and  $U$  are linear then so is  $T \circ U$ .

# Composition of Transformations


**Why** is this the correct definition of matrix multiplication?

## Definition

Let  $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$  and  $U: \mathbf{R}^p \rightarrow \mathbf{R}^n$  be transformations. The **composition** is the transformation

$$T \circ U: \mathbf{R}^p \rightarrow \mathbf{R}^m \quad \text{defined by} \quad T \circ U(x) = T(U(x)).$$

This makes sense because  $U(x)$  (the output of  $U$ ) is in  $\mathbf{R}^n$ , which is the domain of  $T$  (the inputs of  $T$ ).



**Fact:** If  $T$  and  $U$  are linear then so is  $T \circ U$ .

**Guess:** If  $A$  is the matrix for  $T$ , and  $B$  is the matrix for  $U$ , what is the matrix for  $T \circ U$ ?

## Composition of Linear Transformations

Let  $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$  and  $U: \mathbf{R}^p \rightarrow \mathbf{R}^n$  be *linear* transformations. Let  $A$  and  $B$  be their matrices:

## Composition of Linear Transformations

Let  $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$  and  $U: \mathbf{R}^p \rightarrow \mathbf{R}^n$  be *linear* transformations. Let  $A$  and  $B$  be their matrices:

$$A = \left( \begin{array}{c|c|c|c} & & & \\ T(e_1) & T(e_2) & \cdots & T(e_n) \\ & & & \end{array} \right) \quad B = \left( \begin{array}{c|c|c|c} & & & \\ U(e_1) & U(e_2) & \cdots & U(e_p) \\ & & & \end{array} \right)$$



## Composition of Linear Transformations

Let  $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$  and  $U: \mathbf{R}^p \rightarrow \mathbf{R}^n$  be *linear* transformations. Let  $A$  and  $B$  be their matrices:

$$A = \left( \begin{array}{c|c|c|c} & & & \\ T(e_1) & T(e_2) & \cdots & T(e_n) \\ & & & \end{array} \right) \quad B = \left( \begin{array}{c|c|c|c} & & & \\ U(e_1) & U(e_2) & \cdots & U(e_p) \\ & & & \end{array} \right)$$

### Question

What is the matrix for  $T \circ U$ ?

## Composition of Linear Transformations

Let  $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$  and  $U: \mathbf{R}^p \rightarrow \mathbf{R}^n$  be *linear* transformations. Let  $A$  and  $B$  be their matrices:

$$A = \left( \begin{array}{c|c|c|c} & & & \\ T(e_1) & T(e_2) & \cdots & T(e_n) \\ & & & \end{array} \right) \quad B = \left( \begin{array}{c|c|c|c} & & & \\ U(e_1) & U(e_2) & \cdots & U(e_p) \\ & & & \end{array} \right)$$

### Question

What is the matrix for  $T \circ U$ ?

The matrix of the composition is the product of the matrices!

# Composition of Linear Transformations

## Example

Let  $T: \mathbf{R}^2 \rightarrow \mathbf{R}^2$  be rotation by  $45^\circ$ , and let  $U: \mathbf{R}^2 \rightarrow \mathbf{R}^2$  be projection onto the  $x$ -axis. Let's compute their standard matrices  $A$  and  $B$ :

# Composition of Linear Transformations

## Example

Let  $T: \mathbf{R}^2 \rightarrow \mathbf{R}^2$  be rotation by  $45^\circ$ , and let  $U: \mathbf{R}^2 \rightarrow \mathbf{R}^2$  be projection onto the  $x$ -axis. Let's compute their standard matrices  $A$  and  $B$ :

$$\Rightarrow \quad A = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} \quad B = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$$

# Composition of Linear Transformations

Example, continued

So the matrix  $C$  for  $T \circ U$  is

# Composition of Linear Transformations

Example, continued

So the matrix  $C$  for  $T \circ U$  is

Check:

# Composition of Linear Transformations

Example, continued

So the matrix  $C$  for  $T \circ U$  is

Check:

$$\Rightarrow C = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix} \quad \checkmark$$

# Composition of Linear Transformations

Another example

Let

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix} \quad B = \begin{pmatrix} 1 & -3 \\ 2 & -2 \\ 3 & -1 \end{pmatrix}.$$

Let  $T(x) = Ax$  and  $U(y) = By$ , so

$$T: \mathbf{R}^3 \longrightarrow \mathbf{R}^2 \quad U: \mathbf{R}^2 \longrightarrow \mathbf{R}^3 \quad T \circ U: \mathbf{R}^2 \longrightarrow \mathbf{R}^2.$$



# Composition of Linear Transformations

Another example

Let

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix} \quad B = \begin{pmatrix} 1 & -3 \\ 2 & -2 \\ 3 & -1 \end{pmatrix}.$$

Let  $T(x) = Ax$  and  $U(y) = By$ , so

$$T: \mathbf{R}^3 \longrightarrow \mathbf{R}^2 \quad U: \mathbf{R}^- \longrightarrow \mathbf{R}^- \quad T \circ U: \mathbf{R}^- \longrightarrow \mathbf{R}^-.$$

# Composition of Linear Transformations

Another example

Let

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix} \quad B = \begin{pmatrix} 1 & -3 \\ 2 & -2 \\ 3 & -1 \end{pmatrix}.$$

Let  $T(x) = Ax$  and  $U(y) = By$ , so

$$T: \mathbf{R}^3 \longrightarrow \mathbf{R}^2 \quad U: \mathbf{R}^2 \longrightarrow \mathbf{R}^3 \quad T \circ U: \mathbf{R}^2 \longrightarrow \mathbf{R}^2.$$

# Composition of Linear Transformations

Another example

Let

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix} \quad B = \begin{pmatrix} 1 & -3 \\ 2 & -2 \\ 3 & -1 \end{pmatrix}.$$

Let  $T(x) = Ax$  and  $U(y) = By$ , so

$$T: \mathbf{R}^3 \longrightarrow \mathbf{R}^2 \quad U: \mathbf{R}^2 \longrightarrow \mathbf{R}^3 \quad T \circ U: \mathbf{R}^2 \longrightarrow \mathbf{R}^2.$$

# Composition of Linear Transformations

Another example

Let

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix} \quad B = \begin{pmatrix} 1 & -3 \\ 2 & -2 \\ 3 & -1 \end{pmatrix}.$$

Let  $T(x) = Ax$  and  $U(y) = By$ , so

$$T: \mathbf{R}^3 \longrightarrow \mathbf{R}^2 \quad U: \mathbf{R}^2 \longrightarrow \mathbf{R}^3 \quad T \circ U: \mathbf{R}^2 \longrightarrow \mathbf{R}^2.$$

Let's find the matrix for  $T \circ U$ :

$$T \circ U(e_1) =$$

$$T \circ U(e_2) =$$

# Composition of Linear Transformations

Another example

Let

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix} \quad B = \begin{pmatrix} 1 & -3 \\ 2 & -2 \\ 3 & -1 \end{pmatrix}.$$

Let  $T(x) = Ax$  and  $U(y) = By$ , so

$$T: \mathbf{R}^3 \longrightarrow \mathbf{R}^2 \quad U: \mathbf{R}^2 \longrightarrow \mathbf{R}^3 \quad T \circ U: \mathbf{R}^2 \longrightarrow \mathbf{R}^2.$$

Let's find the matrix for  $T \circ U$ :

$$T \circ U(e_1) =$$

$$T \circ U(e_2) =$$

Before we computed  $AB = \begin{pmatrix} 14 & -10 \\ 32 & -28 \end{pmatrix}$ , so  $AB$  is the matrix of  $T \circ U$ .

Poll

Do there exist *nonzero* matrices  $A$  and  $B$  with  $AB = 0$ ?

## Poll

Do there exist *nonzero* matrices  $A$  and  $B$  with  $AB = 0$ ?

Yes! Here's an example:

$$\begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 1 & 1 \end{pmatrix} = \left( \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right) = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}.$$

# The Row-Column Rule for Matrix Multiplication

**Recall:** A row vector of length  $n$  times a column vector of length  $n$  is a scalar:

$$(a_1 \quad \cdots \quad a_n) \begin{pmatrix} b_1 \\ \vdots \\ b_n \end{pmatrix} = a_1 b_1 + \cdots + a_n b_n.$$



# The Row-Column Rule for Matrix Multiplication

**Recall:** A row vector of length  $n$  times a column vector of length  $n$  is a scalar:

$$(a_1 \quad \cdots \quad a_n) \begin{pmatrix} b_1 \\ \vdots \\ b_n \end{pmatrix} = a_1 b_1 + \cdots + a_n b_n.$$

Another way of multiplying a matrix by a vector is:

$$Ax = \begin{pmatrix} -r_1- \\ \vdots \\ -r_m- \end{pmatrix} x = \begin{pmatrix} r_1 x \\ \vdots \\ r_m x \end{pmatrix}.$$

# The Row-Column Rule for Matrix Multiplication

**Recall:** A row vector of length  $n$  times a column vector of length  $n$  is a scalar:

$$(a_1 \quad \cdots \quad a_n) \begin{pmatrix} b_1 \\ \vdots \\ b_n \end{pmatrix} = a_1 b_1 + \cdots + a_n b_n.$$

Another way of multiplying a matrix by a vector is:

$$Ax = \begin{pmatrix} -r_1- \\ \vdots \\ -r_m- \end{pmatrix} x = \begin{pmatrix} r_1 x \\ \vdots \\ r_m x \end{pmatrix}.$$

On the other hand, you multiply two matrices by

$$AB = A \begin{pmatrix} | & & | \\ c_1 & \cdots & c_p \\ | & & | \end{pmatrix} = \begin{pmatrix} | & & | \\ Ac_1 & \cdots & Ac_p \\ | & & | \end{pmatrix}.$$

# The Row-Column Rule for Matrix Multiplication

**Recall:** A row vector of length  $n$  times a column vector of length  $n$  is a scalar:

$$(a_1 \quad \cdots \quad a_n) \begin{pmatrix} b_1 \\ \vdots \\ b_n \end{pmatrix} = a_1 b_1 + \cdots + a_n b_n.$$

Another way of multiplying a matrix by a vector is:

$$A\mathbf{x} = \begin{pmatrix} -r_1- \\ \vdots \\ -r_m- \end{pmatrix} \mathbf{x} = \begin{pmatrix} r_1 x \\ \vdots \\ r_m x \end{pmatrix}.$$

On the other hand, you multiply two matrices by

$$AB = A \left( \begin{array}{c|c|c} | & & | \\ c_1 & \cdots & c_p \\ | & & | \end{array} \right) = \left( \begin{array}{c|c|c} | & & | \\ Ac_1 & \cdots & Ac_p \\ | & & | \end{array} \right).$$

It follows that

$$AB = \begin{pmatrix} -r_1- \\ \vdots \\ -r_m- \end{pmatrix} \left( \begin{array}{c|c|c} | & & | \\ c_1 & \cdots & c_p \\ | & & | \end{array} \right) = \begin{pmatrix} r_1 c_1 & r_1 c_2 & \cdots & r_1 c_p \\ r_2 c_1 & r_2 c_2 & \cdots & r_2 c_p \\ \vdots & \vdots & & \vdots \\ r_m c_1 & r_m c_2 & \cdots & r_m c_p \end{pmatrix}$$

## The Row-Column Rule for Matrix Multiplication

The  $ij$  entry of  $C = AB$  is the  $i$ th row of  $A$  times the  $j$ th column of  $B$ :

$$c_{ij} = (AB)_{ij} = a_{i1}b_{1j} + a_{i2}b_{2j} + \cdots + a_{in}b_{nj}.$$

## The Row-Column Rule for Matrix Multiplication

The  $ij$  entry of  $C = AB$  is the  $i$ th row of  $A$  times the  $j$ th column of  $B$ :

$$c_{ij} = (AB)_{ij} = a_{i1}b_{1j} + a_{i2}b_{2j} + \cdots + a_{in}b_{nj}.$$

This is how everybody on the planet actually computes  $AB$ .

# The Row-Column Rule for Matrix Multiplication

The  $ij$  entry of  $C = AB$  is the  $i$ th row of  $A$  times the  $j$ th column of  $B$ :

$$c_{ij} = (AB)_{ij} = a_{i1}b_{1j} + a_{i2}b_{2j} + \cdots + a_{in}b_{nj}.$$

This is how everybody on the planet actually computes  $AB$ . Diagram  $(AB = C)$ :

$$\begin{pmatrix} a_{11} & \cdots & a_{1k} & \cdots & a_{1n} \\ \vdots & & \vdots & & \vdots \\ a_{i1} & \cdots & a_{ik} & \cdots & a_{in} \\ \vdots & & \vdots & & \vdots \\ a_{m1} & \cdots & a_{mk} & \cdots & a_{mn} \end{pmatrix} \cdot \begin{pmatrix} b_{11} & \cdots & b_{1j} & \cdots & b_{1p} \\ \vdots & & \vdots & & \vdots \\ b_{k1} & \cdots & b_{kj} & \cdots & b_{kp} \\ \vdots & & \vdots & & \vdots \\ b_{n1} & \cdots & b_{nj} & \cdots & b_{np} \end{pmatrix} = \begin{pmatrix} c_{11} & \cdots & c_{1j} & \cdots & c_{1p} \\ \vdots & & \vdots & & \vdots \\ c_{i1} & \cdots & c_{ij} & \cdots & c_{ip} \\ \vdots & & \vdots & & \vdots \\ c_{m1} & \cdots & c_{mj} & \cdots & c_{mp} \end{pmatrix}$$

$j$ th column                       $i$ th row                       $i$ th row                       $j$ th column                       $ij$  entry

# The Row-Column Rule for Matrix Multiplication

The  $ij$  entry of  $C = AB$  is the  $i$ th row of  $A$  times the  $j$ th column of  $B$ :

$$c_{ij} = (AB)_{ij} = a_{i1}b_{1j} + a_{i2}b_{2j} + \cdots + a_{in}b_{nj}.$$

This is how everybody on the planet actually computes  $AB$ . Diagram ( $AB = C$ ):

$$\begin{pmatrix} a_{11} & \cdots & a_{1k} & \cdots & a_{1n} \\ \vdots & & \vdots & & \vdots \\ a_{i1} & \cdots & a_{ik} & \cdots & a_{in} \\ \vdots & & \vdots & & \vdots \\ a_{m1} & \cdots & a_{mk} & \cdots & a_{mn} \end{pmatrix} \cdot \begin{pmatrix} b_{11} & \cdots & b_{1j} & \cdots & b_{1p} \\ \vdots & & \vdots & & \vdots \\ b_{k1} & \cdots & b_{kj} & \cdots & b_{kp} \\ \vdots & & \vdots & & \vdots \\ b_{n1} & \cdots & b_{nj} & \cdots & b_{np} \end{pmatrix} = \begin{pmatrix} c_{11} & \cdots & c_{1j} & \cdots & c_{1p} \\ \vdots & & \vdots & & \vdots \\ c_{i1} & \cdots & c_{ij} & \cdots & c_{ip} \\ \vdots & & \vdots & & \vdots \\ c_{m1} & \cdots & c_{mj} & \cdots & c_{mp} \end{pmatrix}$$

$j$ th column                       $ij$  entry

Example

## Properties of Matrix Multiplication

Mostly matrix multiplication works like you'd expect. Suppose  $A$  has size  $m \times n$ , and that the other matrices below have the right size to make multiplication work.



## Properties of Matrix Multiplication

Mostly matrix multiplication works like you'd expect. Suppose  $A$  has size  $m \times n$ , and that the other matrices below have the right size to make multiplication work.

Most of these are easy to verify.

## Properties of Matrix Multiplication

Mostly matrix multiplication works like you'd expect. Suppose  $A$  has size  $m \times n$ , and that the other matrices below have the right size to make multiplication work.

Most of these are easy to verify.

**Associativity** is  $A(BC) = (AB)C$ . It is a pain to verify using the row-column rule!

## Properties of Matrix Multiplication

Mostly matrix multiplication works like you'd expect. Suppose  $A$  has size  $m \times n$ , and that the other matrices below have the right size to make multiplication work.

Most of these are easy to verify.

**Associativity** is  $A(BC) = (AB)C$ . It is a pain to verify using the row-column rule! Much easier: use associativity of linear transformations:

$$S \circ (T \circ U) = (S \circ T) \circ U.$$

## Properties of Matrix Multiplication

Mostly matrix multiplication works like you'd expect. Suppose  $A$  has size  $m \times n$ , and that the other matrices below have the right size to make multiplication work.

Most of these are easy to verify.

**Associativity** is  $A(BC) = (AB)C$ . It is a pain to verify using the row-column rule! Much easier: use associativity of linear transformations:

$$S \circ (T \circ U) = (S \circ T) \circ U.$$

This is a good example of an instance where having a conceptual viewpoint saves you a lot of work.

## Properties of Matrix Multiplication

Mostly matrix multiplication works like you'd expect. Suppose  $A$  has size  $m \times n$ , and that the other matrices below have the right size to make multiplication work.

Most of these are easy to verify.

**Associativity** is  $A(BC) = (AB)C$ . It is a pain to verify using the row-column rule! Much easier: use associativity of linear transformations:

$$S \circ (T \circ U) = (S \circ T) \circ U.$$

This is a good example of an instance where having a conceptual viewpoint saves you a lot of work.

**Recommended:** Try to verify all of them on your own.

# Properties of Matrix Multiplication

## Caveats

### Warnings!

- ▶  $AB$  is usually not equal to  $BA$ .

# Properties of Matrix Multiplication

## Caveats

### Warnings!

- ▶  $AB$  is usually not equal to  $BA$ .

In fact,  $AB$  may be defined when  $BA$  is not.

# Properties of Matrix Multiplication

## Caveats

### Warnings!

- ▶  $AB$  is usually not equal to  $BA$ .

In fact,  $AB$  may be defined when  $BA$  is not.

- ▶  $AB = AC$  does not imply  $B = C$ , even if  $A \neq 0$ .



# Properties of Matrix Multiplication

## Caveats

### Warnings!

- ▶  $AB$  is usually not equal to  $BA$ .

In fact,  $AB$  may be defined when  $BA$  is not.

- ▶  $AB = AC$  does not imply  $B = C$ , even if  $A \neq 0$ .
- ▶  $AB = 0$  does not imply  $A = 0$  or  $B = 0$ .

Read about powers of a matrix and multiplication of transposes in §2.1.