

Section 1.9

The Matrix of a Linear Transformation



Unit Coordinate Vectors

Definition

The **unit coordinate vectors** in \mathbf{R}^n are

This is what e_1, e_2, \dots mean,
for the rest of the class.

$$e_1 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad \dots, \quad e_{n-1} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ 0 \end{pmatrix}, \quad e_n = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}.$$



Note: if A is an $m \times n$ matrix with columns v_1, v_2, \dots, v_n , then $Ae_i = v_i$ for $i = 1, 2, \dots, n$: multiplying a matrix by e_i gives you the i th column.

$$\begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 4 \\ 7 \end{pmatrix} \quad \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 2 \\ 5 \\ 8 \end{pmatrix} \quad \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix} \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 3 \\ 6 \\ 9 \end{pmatrix}$$

Linear Transformations are Matrix Transformations

Recall: A matrix A defines a linear transformation T by $T(x) = Ax$.

Theorem

Let $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$ be a linear transformation. Let

$$A = \begin{pmatrix} \left| \begin{array}{c} T(e_1) \\ \vdots \end{array} \right| & \left| \begin{array}{c} T(e_2) \\ \vdots \end{array} \right| & \cdots & \left| \begin{array}{c} T(e_n) \\ \vdots \end{array} \right| \end{pmatrix}.$$

This is an $m \times n$ matrix, and T is the matrix transformation for A : $T(x) = Ax$.

The matrix A is called the **standard matrix** for T .

Take-Away

Linear transformations are the same as matrix transformations.

Dictionary

Linear transformation $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$ \rightsquigarrow $m \times n$ matrix $A = \begin{pmatrix} \left| \begin{array}{c} T(e_1) \\ \vdots \end{array} \right| & \left| \begin{array}{c} T(e_2) \\ \vdots \end{array} \right| & \cdots & \left| \begin{array}{c} T(e_n) \\ \vdots \end{array} \right| \end{pmatrix}$

$T(x) = Ax$ \longleftarrow $m \times n$ matrix A

$T: \mathbf{R}^n \rightarrow \mathbf{R}^m$

Linear Transformations are Matrix Transformations

Continued

Why is a linear transformation a matrix transformation?

Suppose for simplicity that $T: \mathbf{R}^3 \rightarrow \mathbf{R}^2$.

$$\begin{aligned} T \begin{pmatrix} x \\ y \\ z \end{pmatrix} &= T \left(x \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} + y \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} + z \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \right) \\ &= T(x\mathbf{e}_1 + y\mathbf{e}_2 + z\mathbf{e}_3) \\ &= xT(\mathbf{e}_1) + yT(\mathbf{e}_2) + zT(\mathbf{e}_3) \\ &= \begin{pmatrix} | & | & | \\ T(\mathbf{e}_1) & T(\mathbf{e}_2) & T(\mathbf{e}_3) \\ | & | & | \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} \\ &= A \begin{pmatrix} x \\ y \\ z \end{pmatrix}. \end{aligned}$$

Linear Transformations are Matrix Transformations

Example

Before, we defined a **dilation** transformation $T: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ by $T(x) = 1.5x$.
What is its standard matrix?

$$\left. \begin{aligned} T(e_1) &= 1.5e_1 = \begin{pmatrix} 1.5 \\ 0 \end{pmatrix} \\ T(e_2) &= 1.5e_2 = \begin{pmatrix} 0 \\ 1.5 \end{pmatrix} \end{aligned} \right\} \Rightarrow A = \begin{pmatrix} 1.5 & 0 \\ 0 & 1.5 \end{pmatrix}.$$

Check:

$$\begin{pmatrix} 1.5 & 0 \\ 0 & 1.5 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1.5x \\ 1.5y \end{pmatrix} = 1.5 \begin{pmatrix} x \\ y \end{pmatrix} = T \begin{pmatrix} x \\ y \end{pmatrix}.$$

Linear Transformations are Matrix Transformations



Example

Question

What is the matrix for the linear transformation $T: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ defined by

$$T(x) = x \text{ rotated counterclockwise by an angle } \theta?$$

(Check linearity...)




$$\left. \begin{aligned} T(e_1) &= \begin{pmatrix} \cos(\theta) \\ \sin(\theta) \end{pmatrix} \\ T(e_2) &= \begin{pmatrix} -\sin(\theta) \\ \cos(\theta) \end{pmatrix} \end{aligned} \right\} \Rightarrow A = \begin{pmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{pmatrix} \quad \left(\begin{array}{l} \theta = 90^\circ \Rightarrow \\ A = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \\ \text{(from before)} \end{array} \right)$$

Linear Transformations are Matrix Transformations

Example

Question

What is the matrix for the linear transformation $T: \mathbf{R}^3 \rightarrow \mathbf{R}^3$ that reflects through the xy -plane and then projects onto the yz -plane?




$$T(e_1) = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}.$$

Linear Transformations are Matrix Transformations

Example, continued

Question

What is the matrix for the linear transformation $T: \mathbf{R}^3 \rightarrow \mathbf{R}^3$ that reflects through the xy -plane and then projects onto the yz -plane?




$$T(e_2) = e_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}.$$

Linear Transformations are Matrix Transformations

Example, continued

Question

What is the matrix for the linear transformation $T: \mathbf{R}^3 \rightarrow \mathbf{R}^3$ that reflects through the xy -plane and then projects onto the yz -plane?



$$T(e_3) = \begin{pmatrix} 0 \\ 0 \\ -1 \end{pmatrix}.$$

Linear Transformations are Matrix Transformations

Example, continued

Question

What is the matrix for the linear transformation $T: \mathbf{R}^3 \rightarrow \mathbf{R}^3$ that reflects through the xy -plane and then projects onto the yz -plane?



$$\left. \begin{aligned} T(e_1) &= \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \\ T(e_2) &= \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \\ T(e_3) &= \begin{pmatrix} 0 \\ 0 \\ -1 \end{pmatrix} \end{aligned} \right\} \implies A = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}.$$

There is a long list of geometric transformations of \mathbf{R}^2 in §1.9 of Lay. (Reflections over the diagonal, contractions and expansions along different axes, shears, projections, ...) Please look them over.

Onto Transformations

Definition

A transformation $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$ is **onto** (or **surjective**) if the range of T is equal to \mathbf{R}^m (its codomain). In other words, each b in \mathbf{R}^m is the image of *at least one* x in \mathbf{R}^n : every possible output has an input. Note that *not* onto means there is some b in \mathbf{R}^m which is not the image of any x in \mathbf{R}^n .



Characterization of Onto Transformations

Theorem

Let $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$ be a linear transformation with matrix A . Then the following are equivalent:

- ▶ T is onto
- ▶ $T(x) = b$ has a solution for every b in \mathbf{R}^m
- ▶ $Ax = b$ is consistent for every b in \mathbf{R}^m
- ▶ The columns of A span \mathbf{R}^m
- ▶ A has a pivot in every row

Question

If $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$ is onto, what can we say about the relative sizes of n and m ?

Answer: T corresponds to an $m \times n$ matrix A . In order for A to have a pivot in every row, it must have *at least as many* columns as rows: $m \leq n$.



$$\begin{pmatrix} \color{red}{1} & 0 & \star & 0 & \star \\ 0 & \color{red}{1} & \star & 0 & \star \\ 0 & 0 & 0 & \color{red}{1} & \star \end{pmatrix}$$

For instance, \mathbf{R}^2 is “too small” to map *onto* \mathbf{R}^3 .

One-to-one Transformations

Definition

A transformation $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$ is **one-to-one** (or **into**, or **injective**) if different vectors in \mathbf{R}^n map to different vectors in \mathbf{R}^m . In other words, each b in \mathbf{R}^m is the image of *at most one* x in \mathbf{R}^n : different inputs have different outputs. Note that *not* one-to-one means different vectors in \mathbf{R}^n have the same image.



Characterization of One-to-One Transformations

Theorem

Let $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$ be a linear transformation with matrix A . Then the following are equivalent:

- ▶ T is one-to-one
- ▶ $T(x) = b$ has one or zero solutions for every b in \mathbf{R}^m
- ▶ $Ax = b$ has a unique solution or is inconsistent for every b in \mathbf{R}^m
- ▶ $Ax = 0$ has a unique solution
- ▶ The columns of A are linearly independent
- ▶ A has a pivot in every column.

Question

If $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$ is one-to-one, what can we say about the relative sizes of n and m ?

Answer: T corresponds to an $m \times n$ matrix A . In order for A to have a pivot in every column, it must have *at least as many rows as columns*: $n \leq m$.

$$\begin{pmatrix} \color{red}{1} & 0 & 0 \\ 0 & \color{red}{1} & 0 \\ 0 & 0 & \color{red}{1} \\ 0 & 0 & 0 \end{pmatrix}$$

For instance, \mathbf{R}^3 is “too big” to map *into* \mathbf{R}^2 .