

Molecular Geometry and Rotational Constant Analysis

Dr. Yukio Yamaguchi

Center for Computational Quantum Chemistry, University of Georgia

with additions and modifications by T. Daniel Crawford, Virginia Tech

1. Read the molecular cartesian coordinates and atomic numbers from the given file.

2. Calculate all possible interatomic distances, R_{ij} .

$$R_{ij} = \sqrt{(X_i - X_j)^2 + (Y_i - Y_j)^2 + (Z_i - Z_j)^2} \quad (1)$$

3. Calculate all possible bond angles. For example, the angle, ϕ_{ijk} , between atoms $\mathbf{i} - \mathbf{j} - \mathbf{k}$, where \mathbf{j} is the central atom is given by:

$$\cos \phi_{ijk} = \mathbf{e}_{ji} \cdot \mathbf{e}_{jk} \quad (2)$$

where the \mathbf{e}_{ij} are unit vectors between the atoms, e.g.,

$$e_{ij}^X = -(X_i - X_j) / R_{ij} \quad e_{ij}^Y = -(Y_i - Y_j) / R_{ij} \quad e_{ij}^Z = -(Z_i - Z_j) / R_{ij} \quad (3)$$

4. Calculate all possible out-of-plane angles. For example, the angle θ_{ijkl} for atom \mathbf{i} out of the plane containing atoms $\mathbf{j} - \mathbf{k} - \mathbf{l}$ (with \mathbf{k} as the central atom) is given by:

$$\sin \theta_{ijkl} = \frac{\mathbf{e}_{kj} \times \mathbf{e}_{kl}}{\sin \phi_{jkl}} \cdot \mathbf{e}_{ki} \quad (4)$$

5. Calculate all possible torsional angles. For example, the torsional angle τ_{ijkl} for the atom connectivity $\mathbf{i} - \mathbf{j} - \mathbf{k} - \mathbf{l}$ is given by:

$$\cos \tau_{ijkl} = \frac{(\mathbf{e}_{ij} \times \mathbf{e}_{jk}) \cdot (\mathbf{e}_{jk} \times \mathbf{e}_{kl})}{\sin \phi_{ijk} \sin \phi_{jkl}} \quad (5)$$

Can you also determine the sign of the torsional angle?

6. Find the center of mass of the molecule.

$$X_{c.m.} = \frac{\sum_i m_i X_i}{\sum_i m_i} \quad Y_{c.m.} = \frac{\sum_i m_i Y_i}{\sum_i m_i} \quad Z_{c.m.} = \frac{\sum_i m_i Z_i}{\sum_i m_i}, \quad (6)$$

where m_i is the mass of atom i and the summation runs over all atoms in the molecule.

7. Calculate elements of the moment of inertia tensor.

$$I_{\alpha\alpha} = \sum_i m_i (\beta_i^2 + \gamma_i^2) \quad (7)$$

$$I_{\alpha\beta} = \sum_i m_i \alpha_i \beta_i, \quad (8)$$

where α , β , and γ correspond to choices of x , y , and z (e.g., I_{xy} is one choice of $I_{\alpha\beta}$).

8. Diagonalize the inertia tensor to obtain the principal moments of inertia.

$$I_a \leq I_b \leq I_c \quad (9)$$

9. Determine the molecular type:

- diatomic
- linear polyatomic
- asymmetric top
- symmetric top (prolate or oblate)
- spherical top

10. Determine the moments of inertia in $\text{amu}\cdot\text{\AA}^2$ and $\text{g}\cdot\text{cm}^2$ and determine the rotational constants in cm^{-1} and MHz.

$$A \geq B \geq C \quad (10)$$

$$A = \frac{h}{8\pi^2 I_a} \quad B = \frac{h}{8\pi^2 I_b} \quad C = \frac{h}{8\pi^2 I_c} \quad (11)$$

For more information see E.B. Wilson, J.C. Decius, and P.C. Cross, '**Molecular Vibrations**', McGraw-Hill, 1955.