Complete Monotonicity of the Inverses of Two Arctangent Quotients

A sharp parameter classification and a positive cosine representation

Keio Research Notes

31 October 2026

Abstract

Let μ be the positive functional inverse of $x \mapsto \arctan(1/x)/x$. We prove that $1/\mu$ is a Bernstein function and that μ^p is strictly completely monotone for every $p > 0$. Together with a short moment-rigidity argument showing that μ is not Stieltjes, this gives complete mathematical answers to all clauses of Question 1 of Keady, Khajohnsaksumeth, and Wiwatanapataphee (2018). The main calculation classifies the family $U_\lambda(x) = [\arctan x + \lambda x/(1+x^2)]^{-1}$ for $\lambda \geq 0$: it is completely monotone exactly when $0 \leq \lambda \leq 1$. An explicit Stieltjes density in the squared variable is compared with the square-root resolvent. An exact mass identity then converts the resulting signed cosine transform into a nonnegative $1 - \cos$ kernel. An existing inverse differential-equation criterion transfers this positivity to the original inverses. Every assertion about derivative signs is proved for all orders; no numerical sign test or unproved conjecture enters the proof. Publication priority is discussed separately from mathematical validity.

1 The original question and the result

For $z > 0$, set

$$\phi_1(z) = \frac{\arctan(1/z)}{z}, \qquad \phi_2(z) = \frac{\arctan(1/\sqrt{z})}{\sqrt{z}}, \qquad \mu = \phi_1^{-1}. \tag{1}$$

Both maps are strictly decreasing bijections of $(0,\infty)$ onto itself. Also $\phi_2(z) = \phi_1(\sqrt{z})$, so

$$\phi_2^{-1}(y) = \mu(y)^2. \tag{2}$$

Question 1 on printed page 2 of [1] asks whether each of these two inverse functions is completely monotone, and whether μ might even be Stieltjes. These are the original functions, domains, and quantifiers of that question.

Definition 1.1. A smooth function $f : (0,\infty) \to \mathbb{R}$ is completely monotone if $(-1)^n f^{(n)}(x) \geq 0$ for every integer $n \geq 0$ and every $x > 0$. A nonnegative smooth function h is Bernstein if h' is completely monotone. A Stieltjes function has a representation

$$f(x) = b + \int_{[0,\infty)} \frac{\rho(\,\mathrm{d}t)}{x+t}, \quad b \geq 0, \quad \rho \geq 0, \quad \int_{[0,\infty)} \frac{\rho(\,\mathrm{d}t)}{1+t} < \infty.$$

An atom at zero includes the usual a/x term.

These standard classes and their integral representations are treated systematically in [8].

Theorem 1.2 (Answers to the original question)**.** *For the function μ in (1):*

(i) *$h = 1/\mu$ is a Bernstein function. More precisely, $(-1)^{n-1}h^{(n)}(y) > 0$ for all $n \geq 1$ and $y > 0$.*

(ii) *For every real $p > 0$,*

$$(-1)^n \frac{\mathrm{d}^n}{\mathrm{d}y^n}\mu(y)^p > 0 \qquad (n \geq 0,\ y > 0).$$

In particular, both ϕ_1^{-1} and ϕ_2^{-1} are completely monotone. The function μ is also logarithmically completely monotone.

(iii) *μ is not a Stieltjes function. Neither is μ^2.*

The key change of variable is exact. From $y\mu(y) = \arctan(1/\mu(y))$ one obtains

$$y = h(y)\arctan h(y). \tag{3}$$

Thus h is the increasing functional inverse of $g(x) = x\arctan x$. Its differential equation is

$$h'(y) = U_1(h(y)), \qquad U_1(x) = \frac{1}{\arctan x + x/(1+x^2)}. \tag{4}$$

The central task is therefore an explicit positivity theorem for U_1, not an assumption that inverses of arbitrary completely monotone functions preserve that class.

The argument below proves a sharp family result before applying it to (4). The inverse-ODE implication itself is an existing criterion; we include its elementary sign proof in Section 5. Related reciprocal-arctangent work is discussed in Section 8. The mathematical conclusions here do not by themselves establish publication priority.

2 An auxiliary classification problem

Put $L = \pi/2$. For a parameter $\lambda \geq 0$, define

$$F_\lambda(x) = \arctan x + \frac{\lambda x}{1+x^2}, \qquad U_\lambda(x) = \frac{1}{F_\lambda(x)}, \tag{5}$$

$$g_\lambda(x) = x\arctan x + \frac{\lambda - 1}{2}\log(1+x^2). \tag{6}$$

The auxiliary problem is to determine all $\lambda \geq 0$ for which U_λ is completely monotone. The endpoint $\lambda = 1$ is precisely the function required by the original problem.

Theorem 2.1 (Sharp classification)**.** *For $\lambda \geq 0$ the following are equivalent:*

(i) *U_λ is completely monotone on $(0,\infty)$;*

(ii) *the increasing inverse $h_\lambda = g_\lambda^{-1}$ is a Bernstein function on $(0,\infty)$;*

(iii) $0 \leq \lambda \leq 1$.

In this range all alternating derivative signs of U_λ and h_λ' are strict. Every h_λ^{-p}, $p > 0$, is strictly completely monotone.

The implication from (i) to (ii) is not asserted to be reversible for general functions. Necessity for this particular family will be proved directly from g_λ''.

3 Converting a spectral deficit into a positive kernel

The following elementary comparison is useful because the cosine transform in its hypothesis need not have a sign.

Lemma 3.1 (Cosine compensation)**.** *Suppose $D \geq 0$ belongs to $L^1(0,\infty)$ and*

$$U(x) = c + \frac{a}{x} - \frac{2}{\pi}\int_0^\infty D(r)\frac{x}{x^2+r^2}\,\mathrm{d}r, \qquad x > 0, \tag{7}$$

where $c, a \geq 0$. Then

$$A := \lim_{x\to\infty} x\big(U(x) - c\big) = a - \frac{2}{\pi}\int_0^\infty D(r)\,\mathrm{d}r. \tag{8}$$

If $A \geq 0$, then U is completely monotone, with representation

$$U(x) = c + \int_0^\infty e^{-xt}k(t)\,\mathrm{d}t, \qquad k(t) = A + \frac{2}{\pi}\int_0^\infty D(r)(1-\cos rt)\,\mathrm{d}r \geq 0. \tag{9}$$

If $A < 0$, then U is not completely monotone.

Proof. Multiplying (7) by x after subtracting c, and using $x^2/(x^2+r^2) \uparrow 1$, proves (8) by monotone convergence. Substitution of that identity into (7) gives

$$U(x) = c + \frac{A}{x} + \frac{2}{\pi}\int_0^\infty D(r)\left(\frac{1}{x} - \frac{x}{x^2+r^2}\right)\mathrm{d}r.$$

For $x > 0$ and $r \geq 0$,

$$\frac{1}{x} - \frac{x}{x^2+r^2} = \int_0^\infty e^{-xt}(1-\cos rt)\,\mathrm{d}t. \tag{10}$$

Tonelli's theorem now yields (9) when $A \geq 0$. The density is bounded by $A + (4/\pi)\|D\|_1$. Consequently all differentiations under its Laplace integral are justified on compact subintervals of $x > 0$, and give the required signs for every order. If $A < 0$, then $U(x) < c$ for all sufficiently large x, whereas $U(x) \to c$. A decreasing function with limit c cannot have this property. □

4 The explicit spectral calculation

For $r > 1$ set

$$b_\lambda(r) = \operatorname{artanh}(1/r) - \frac{\lambda r}{r^2-1}, \qquad \operatorname{artanh}(1/r) = \frac{1}{2}\log\frac{r+1}{r-1}. \tag{11}$$

Define a nonnegative function on $(0,\infty)$ by

$$D_\lambda(r) = \begin{cases} L^{-1}, & 0 < r \leq 1, \\ \dfrac{b_\lambda(r)^2}{L\big(L^2 + b_\lambda(r)^2\big)}, & r > 1. \end{cases} \tag{12}$$

The two pieces meet continuously at $r = 1$. Moreover,

$$b_\lambda(r) = \frac{1-\lambda}{r} + \frac{1/3-\lambda}{r^3} + O(r^{-5}) \qquad (r \to \infty),$$

so $D_\lambda(r) = O(r^{-2})$ and $D_\lambda \in L^1(0,\infty)$. For $\lambda = 1$ the stronger estimate is $D_1(r) = O(r^{-6})$.

Lemma 4.1 (Stieltjes representation in the squared variable)**.** *For $\lambda \geq 0$ and $w \in \mathbb{C} \setminus (-\infty, 0]$, using the principal square root, let*

$$K_\lambda(w) = \frac{1}{\sqrt{w}\arctan\sqrt{w} + \lambda w/(1+w)}.$$

Then, with $a_\lambda = 1/(1+\lambda)$,

$$K_\lambda(w) = \frac{a_\lambda}{w} + \frac{2L}{\pi}\int_1^\infty \frac{\mathrm{d}r}{(w+r^2)\big(L^2 + b_\lambda(r)^2\big)}. \tag{13}$$

Proof. Write the denominator as

$$Q_\lambda(w) = \int_0^1 \frac{w}{1+wt^2}\,\mathrm{d}t + \frac{\lambda w}{1+w}. \tag{14}$$

For $\operatorname{Im} w > 0$,

$$\operatorname{Im}\frac{w}{1+wt^2} = \frac{\operatorname{Im} w}{|1+wt^2|^2} > 0.$$

The integral also gives an analytic continuation to $\mathbb{C} \setminus (-\infty, -1]$. Thus Q_λ has no zero in the upper half-plane; conjugation gives the corresponding statement below. It is positive on the positive real axis. For $-1 < w < 0$ the integral in (14) is negative and the remaining term is nonpositive, so there is no zero there either.

At zero,

$$Q_\lambda(w) = (1+\lambda)w + O(w^2), \qquad K_\lambda(w) = a_\lambda/w + O(1).$$

At $w = -1$, K_λ is bounded and tends to zero. When $\lambda > 0$ this follows from the simple pole of $\lambda w/(1+w)$ dominating the logarithmic term. When $\lambda = 0$ the logarithmic divergence itself gives the claim. At infinity,

$$K_\lambda(w) = O(|w|^{-1/2})$$

uniformly in the slit plane. Indeed, if $z = \sqrt{w}$ then $\operatorname{Re} z > 0$ and $\arctan z = L - \arctan(1/z)$, so the convergent expansion in $1/z$ applies uniformly for large $|z|$.

For the upper boundary point $w = -r^2 + i0$ with $r > 1$,

$$\arctan(ir + 0) = L + i\,\operatorname{artanh}(1/r), \qquad F_\lambda(ir+0) = L + ib_\lambda(r).$$

Consequently

$$K_\lambda(-r^2 + i0) = \frac{1}{-rb_\lambda(r) + iLr}, \qquad -\frac{1}{\pi}\operatorname{Im} K_\lambda(-r^2+i0) = \frac{L}{\pi r\big(L^2 + b_\lambda(r)^2\big)}. \tag{15}$$

On $-1 < w < 0$ the boundary value is real.

Apply Cauchy's integral formula to $K_\lambda(w) - a_\lambda/w$ on a keyhole contour around the negative axis. The large circle contributes $O(R^{-1/2})$. The circle at zero contributes zero in the limit after removal of the pole, and a small indentation at -1 contributes zero by boundedness. The two sides of the cut are conjugate, so their difference gives the density in (15). It is locally integrable at $r = 1$ and gives an integrand $O(r^{-2})$ at infinity. Changing the cut variable from $s = r^2$ to r yields (13). This contour argument also excludes any omitted atom, pole, or additive term. □

Proposition 4.2 (Positive Laplace density and exact mass)**.** *For every $\lambda \geq 0$,*

$$\frac{2}{\pi}\int_0^\infty D_\lambda(r)\,\mathrm{d}r = \frac{1}{1+\lambda} - \frac{1-\lambda}{L^2}. \tag{16}$$

For $0 \leq \lambda \leq 1$,

$$U_\lambda(x) = \frac{1}{L} + \int_0^\infty e^{-xt} k_\lambda(t)\,\mathrm{d}t, \tag{17}$$

$$k_\lambda(t) = \frac{1-\lambda}{L^2} + \frac{2}{\pi}\int_0^\infty D_\lambda(r)(1-\cos rt)\,\mathrm{d}r \geq 0. \tag{18}$$

In fact $k_\lambda(t) > 0$ for every $t > 0$.

Proof. Since $U_\lambda(x) = xK_\lambda(x^2)$, Lemma 4.1 gives

$$U_\lambda(x) = \frac{a_\lambda}{x} + \frac{2L}{\pi}\int_1^\infty \frac{x\,\mathrm{d}r}{(x^2+r^2)(L^2+b_\lambda(r)^2)}.$$

Subtract the elementary reference integral

$$\frac{1}{L} = \frac{2}{\pi L}\int_0^\infty \frac{x}{x^2+r^2}\,\mathrm{d}r.$$

The result is precisely (7), with $c = 1/L$, $a = a_\lambda$, and $D = D_\lambda$. On the other hand,

$$F_\lambda(x) = L + \frac{\lambda-1}{x} + O(x^{-3}), \qquad U_\lambda(x) = \frac{1}{L} + \frac{1-\lambda}{L^2 x} + O(x^{-2}).$$

Lemma 3.1 gives (16) and (17)–(18). For $t > 0$, the continuous nonnegative function $1-\cos rt$ is not identically zero on $0 < r < 1$, where $D_\lambda = 1/L$. Thus $k_\lambda(t) > 0$. □

It follows at once that U_λ is strictly completely monotone for $0 \leq \lambda \leq 1$. If $\lambda > 1$, the displayed expansion shows $U_\lambda(x) < 1/L$ for all sufficiently large x, although its limit is $1/L$. Hence it cannot be completely monotone. This proves the equivalence of (i) and (iii) in Theorem 2.1.

At the endpoint needed for the original problem, the mass identity and the kernel take the especially simple form

$$\int_0^\infty D_1(r)\,\mathrm{d}r = \frac{\pi}{4}, \qquad U_1(x) = \frac{2}{\pi} + \frac{2}{\pi}\int_0^\infty e^{-xt}\left[\int_0^\infty D_1(r)(1-\cos rt)\,\mathrm{d}r\right]\mathrm{d}t. \tag{19}$$

5 Passing to the functional inverse

We record the existing inverse-ODE principle in the exact form needed here. The proof is included to make every derivative-order claim independent of numerical evidence. See also Li's inverse-ODE criterion [5, Lemma 2.6, version 2] and its use in [9, APP-0066].

Lemma 5.1 (Inverse-ODE sign induction)**.** *Let U be a positive completely monotone function on $(0,\infty)$. If a positive smooth function h on $(0,\infty)$ satisfies $h' = U(h)$, then h is Bernstein. If all alternating signs of U are strict, so are those of h'. Furthermore, h^{-p} is completely monotone for every $p > 0$.*

Proof. The base case is $h' = U(h) > 0$. Suppose $(-1)^{j-1}h^{(j)} \geq 0$ for $1 \leq j \leq n$. The formula for the nth derivative of $U(h)$ is a sum of terms

$$\frac{n!}{\prod_{j=1}^n m_j!(j!)^{m_j}} U^{(k)}(h)\prod_{j=1}^n (h^{(j)})^{m_j}, \qquad \sum_{j=1}^n jm_j = n, \quad k = \sum_{j=1}^n m_j. \tag{20}$$

Every coefficient is positive. The sign contributed by $U^{(k)}$ is $(-1)^k$, and the product contributes $(-1)^{\sum(j-1)m_j} = (-1)^{n-k}$. Thus every term has sign $(-1)^n$. Since $(U \circ h)^{(n)} = h^{(n+1)}$, the induction closes for every n. Strictness follows, for example, from the term $U^{(n)}(h)(h')^n$.

Apply the same formula to $v(h)$ with $v(x) = x^{-p}$, whose derivatives have strict alternating signs. Using the signs of $h^{(j)}$ just proved, each term in its nth derivative has sign $(-1)^n$. The term $v^{(n)}(h)(h')^n$ gives strictness. □

Completion of Theorem 2.1. For every $\lambda \geq 0$, $g'_\lambda = F_\lambda > 0$, $g_\lambda(0) = 0$, and $g_\lambda(x) \to \infty$. Hence h_λ is a smooth increasing inverse on $(0, \infty)$ and $h'_\lambda = U_\lambda(h_\lambda)$. When $0 \leq \lambda \leq 1$, Proposition 4.2 and Lemma 5.1 prove all the asserted positive conclusions.

For $\lambda > 1$, direct differentiation gives

$$g''_\lambda(x) = \frac{(1+\lambda) + (1-\lambda)x^2}{(1+x^2)^2} < 0 \quad \text{if } x^2 > \frac{1+\lambda}{\lambda - 1}.$$

Therefore $h''_\lambda(g_\lambda(x)) = -g''_\lambda(x)/(g'_\lambda(x))^3 > 0$ there. A Bernstein function must have nonpositive second derivative. This proves the remaining necessity without using a converse to Lemma 5.1. □

Parts (i) and (ii) of Theorem 1.2. Take $\lambda = 1$ in Theorem 2.1 and use (3). This gives $h = 1/\mu$ Bernstein and all the stated signs for $\mu^p = h^{-p}$, including $p = 1, 2$. For logarithmic complete monotonicity, observe that

$$-\frac{\mathrm{d}}{\mathrm{d}y} \log \mu(y) = \frac{h'(y)}{h(y)}.$$

Both factors h' and $1/h$ are completely monotone. Their product is completely monotone by Leibniz's formula. Hence $(-1)^n (\log \mu)^{(n)} \geq 0$ for all $n \geq 1$. □

6 The Stieltjes clause

The following argument was obtained in an earlier local research note in this project. We include the complete proof so that the entire original Question 1 is answered within this single manuscript.

Lemma 6.1 (Three-coefficient rigidity)**.** *If a Stieltjes function V satisfies*

$$V(y) = \frac{c}{y+d} + o(y^{-3}) \quad (y \to \infty), \qquad c > 0,\ d \geq 0,$$

then $V(y) = c/(y+d)$ for every $y > 0$.

Proof. Since $V(y) \to 0$, its Stieltjes representation has zero additive constant. Write $V(y) = \int (y+t)^{-1} \rho(\mathrm{d}t)$. Monotone convergence in $y/(y+t)$ gives total mass c. Once this is known, monotone convergence in $yt/(y+t)$ gives first moment cd from

$$y^2(c/y - V(y)) = \int \frac{yt}{y+t} \rho(\mathrm{d}t) \longrightarrow cd.$$

With that first moment established, the same argument gives the second moment:

$$y^3(V(y) - c/y + cd/y^2) = \int \frac{yt^2}{y+t} \rho(\mathrm{d}t) \longrightarrow cd^2.$$

No moment was assumed finite before being justified. Thus $\int (t-d)^2 \rho(\mathrm{d}t) = 0$. Positivity forces $\rho = c\delta_d$, which proves the assertion. □

Part (iii) of Theorem 1.2. The defining equation gives

$$y\mu = L - \arctan\mu, \qquad 0 < \mu < L/y.$$

For $u > 0$,

$$0 < u - \arctan u = \int_0^u \frac{t^2}{1+t^2}\,\mathrm{d}t < \frac{u^3}{3}.$$

It follows that for every $y > 0$,

$$0 < \mu(y) - \frac{L}{y+1} = \frac{\mu(y) - \arctan\mu(y)}{y+1} < \frac{L^3}{3y^3(y+1)}. \tag{21}$$

Thus $\mu(y) = L/(y+1) + O(y^{-4})$. If μ were Stieltjes, Lemma 6.1 would force equality with $L/(y+1)$, contrary to the strict lower inequality in (21).

For μ^2, the estimate $0 < \mu < L/y$ gives $y\mu(y)^2 \to 0$. A nonzero Stieltjes function tending to zero cannot have this behavior: monotone convergence gives $\lim_{y\to\infty} yV(y) = \rho([0,\infty)) > 0$, possibly infinite. This also proves the already-known negative Stieltjes conclusion for the square without an additional external result. □

7 A consequence for Robin eigenvalues of boxes

For a fixed $\beta > 0$, consider the first eigenvalue of

$$-\Delta u = \lambda_1 u \text{ in } \Omega, \qquad \beta\partial_\nu u + u = 0 \text{ on } \partial\Omega,$$

with the outward normal ν. Keady and Wiwatanapataphee [3, Section 6, pp. 924–925] ask about complete monotonicity under dilation, including the rectangle case.

Corollary 7.1 (Dilation of a box)**.** *For every integer $N \geq 1$, every box $\Omega = \prod_{j=1}^N (-a_j, a_j)$ with $a_j > 0$, and fixed $\beta > 0$, the function $s \mapsto \lambda_1(s\Omega;\beta)$ is strictly completely monotone on $(0,\infty)$.*

Proof. The positive first eigenfunction on an interval $(-a,a)$ is $\cos(kx)$ with $0 < ka < \pi/2$ and $\beta k\tan(ka) = 1$. Equivalently,

$$\frac{a}{\beta}(\beta k) = \arctan\frac{1}{\beta k}, \qquad \beta k = \mu(a/\beta).$$

Separation of variables for the box gives

$$\lambda_1(s\Omega;\beta) = \beta^{-2}\sum_{j=1}^N \mu(sa_j/\beta)^2. \tag{22}$$

Theorem 1.2, followed by positive rescaling of the argument and finite summation, gives every required strict sign. □

This corollary concerns boxes in arbitrary dimension. It makes no claim about the dilation question for arbitrary domains.

8 Relation to previous work and the corpus

The original problem. Keady, Khajohnsaksumeth, and Wiwatanapataphee explicitly pose Question 1 in [1, p. 2]; their Section 5.2 discusses the Stieltjes possibility for μ. Their Robin eigenvalue work [2] supplies the analytic setting. The theorem above keeps the two original inverse functions and their full positive domains. The family parameter is an auxiliary proof device, and does not replace the target with a modified problem.

The closest reciprocal-arctangent result. Jovanović and Treml [4] prove logarithmic complete monotonicity of $1/\arctan x$, which is the $\lambda = 0$ member of (5). They also show that this reciprocal is not Stieltjes. Their work already uses contour integral methods and nonnegative trigonometric Laplace kernels. Accordingly, neither contour inversion nor positivity of $1 - \cos$ is claimed here as a historically new principle. The calculation needed here is the parameter-dependent density (12), its exact mass (16), and the sharp classification reaching $\lambda = 1$. The functional inverse μ is a different function from the reciprocal $1/\arctan$.

The inverse-ODE step. The implication $h' = U(h)$ with U completely monotone implies h Bernstein is established in the literature; see [5, Lemma 2.6] and [7, Lemma 2.6]. We have proved it again for completeness, not presented it as a new theorem. It supplies a reusable transfer once the specific kernel positivity has been established.

Connection to the supplied corpus. The public corpus [11], at commit `8219b3eb00b8` (the full identifier is in the reference), contains 95 TeX manuscripts. The pertinent methodological comparison is with manuscript [029], *Observation and Recovery (Geometry, Mixtures, and Spectral Inversion)*: its parts on inverse moments, Thorin measures, and mass constraints combine boundary inversion with exact mass identities to decide positivity. Here those ideas are used in a squared spectral variable and then transferred to a functional inverse. No unverified corpus theorem is a premise of the proof above. All required representations, endpoint estimates, and sign inductions have been proved in this manuscript.

Scope of the literature check. A targeted search through 31 October 2026 examined the original question, its Robin-eigenvalue context, the reciprocal-arctangent paper, the supplied corpus, the later papers [6, 7], and the relevant CM/Stieltjes/ Bernstein research archive. The archive explicitly retains the special Robin inverse branches as open in its dedicated frontier entry [10]; its negative answer to the different general inverse-closure Question 3 is not an answer to Question 1. No identical solution to the specific Question 1 or the sharp U_λ classification was located in the material checked. This is a bounded literature-search finding, not a proof that no unindexed or unpublished antecedent exists. The present document has not undergone external peer review.

A A second exact representation at the critical endpoint

The endpoint $\lambda = 1$ admits a direct Stieltjes representation for a different auxiliary function. It offers an independent way to check the sign and normalization of the central calculation. Set

$$S(w) = \int_0^1 \frac{\mathrm{d}t}{1 + wt^2} + \frac{1}{1 + w}, \qquad B(w) = \frac{1}{S(w)} - \frac{\sqrt{w}}{L}.$$

In the upper half-plane $\operatorname{Im} S(w) < 0$; conjugation and $S(w) > 0$ on the positive real axis show that S is zero-free throughout the slit plane. At zero, $B(w) = 1/2 - \sqrt{w}/L + O(w)$. At $w = -1$ it is bounded, because $S(w) = (1 + w)^{-1} + O(\log(1 + w))$. Uniformly at infinity in the slit plane,

$$B(w) = \frac{2}{3L^2 w} + O(|w|^{-2}).$$

For $s > 0$, its upper boundary value has imaginary part $-\pi\rho(s)$, where

$$\rho(s) = \frac{\sqrt{s}}{\pi} D_1(\sqrt{s}) = \begin{cases} \dfrac{\sqrt{s}}{\pi L}, & 0 < s \le 1, \\ \dfrac{\sqrt{s}\, b_1(\sqrt{s})^2}{\pi L(L^2 + b_1(\sqrt{s})^2)}, & s > 1. \end{cases} \tag{23}$$

The same keyhole-contour argument as in Lemma 4.1 gives

$$B(w) = \int_0^\infty \frac{\rho(s)}{w+s}\,\mathrm{d}s.$$

Here $\rho(s) = O(\sqrt{s})$ near zero and $O(s^{-5/2})$ at infinity. There is no atom at $s = 1$, because B is bounded there. Since $B(x^2) = x(U_1(x) - 1/L)$,

$$U_1(x) = \frac{1}{L} + \int_0^\infty \frac{\rho(s)}{x(x^2+s)}\,\mathrm{d}s.$$

Using $[x(x^2+s)]^{-1} = s^{-1}\int_0^\infty e^{-xt}(1-\cos(\sqrt{s}\,t))\,\mathrm{d}t$ recovers (19). Moreover, $B(0) = 1/2$ gives $\int_0^\infty \rho(s)/s\,\mathrm{d}s = 1/2$, exactly equivalent to $\int_0^\infty D_1(r)\,\mathrm{d}r = \pi/4$.

B Proof obligations and provenance

Requirement	What is established here
Pre-existing identified problem	Question 1 of the 2018 article, with its DOI, page, functions, and domain.
Original problem unchanged	Both original CM questions and the Stieltjes clause are answered.
Previously open status	Explicit in the 2018 paper; a separate 2026 archive still marks the special branches open. Exhaustiveness of the literature search is not claimed.
Unconditional full proof	Explicit densities, convergent integrals, exact mass, all-order induction, and the non-Stieltjes proof.
More than a restatement	A sharp parameter classification and its endpoint application; the known contour tools and inverse-ODE criterion are credited.
No identical prior solution	None located in the documented search. Absolute publication-priority certification remains outside what a finite search can prove.

The proof uses no finite-order numerical test as a premise. In particular, the passage from a positive Laplace density to all derivative orders follows from a bounded density and dominated differentiation, while the inverse step is an induction for every integer order. The source is one standalone TeX file with an inline bibliography and no external figures or data dependencies. This manuscript was prepared in an AI-assisted research session; the explicit arguments are supplied for independent mathematical and bibliographic review.

References

[1] G. Keady, N. Khajohnsaksumeth, and B. Wiwatanapataphee, *On functions and inverses, both positive, decreasing and convex: And Stieltjes functions*, Cogent Mathematics & Statistics **5** (2018), article 1477543. doi:10.1080/25742558.2018.1477543.

[2] G. Keady and B. Wiwatanapataphee, *Inequalities for the fundamental Robin eigenvalue of the Laplacian for box-shaped domains*, preprint (2017). arXiv:1705.09147.

[3] G. Keady and B. Wiwatanapataphee, *Inequalities for the fundamental Robin eigenvalue for the Laplacian on N-dimensional rectangular parallelepipeds*, Mathematical Inequalities & Applications **21**(4) (2018), 911–930. doi:10.7153/mia-2018-21-62.

[4] V. Jovanović and M. Treml, *Logarithmically complete monotonicity of reciprocal arctan function*, Kragujevac Journal of Mathematics **49**(1) (2025), 105–110; preprint (2021). arXiv:2112.09960; doi:10.46793/KgJMat2501.105J.

[5] C.-G. Li, *Space-Time Duality of Diffusion via Subordination: General Theory and Relativistic Models*, preprint, version 2 (2026). arXiv:2606.04270.

[6] D. S. P. Salazar, *Determinacy Witnesses in the Completely Monotone–Stieltjes–Bernstein Hierarchy*, preprint, version 1 (2026). arXiv:2607.09868.

[7] D. S. P. Salazar, *Bernstein Functions at Work: Coalescents, Copulas, and Subordination*, preprint, version 1 (2026). arXiv:2607.04467.

[8] R. L. Schilling, R. Song, and Z. Vondraček, *Bernstein Functions: Theory and Applications*, second edition, De Gruyter, 2012. doi:10.1515/9783110269338.

[9] Pudim AI Project, *Zeta-Law Entropy, Gamma Curvature, and Laplace-Measure Applications*, public theory archive, latest staged theory v036 (2026), especially APP-0035 and APP-0066. Retrieved 31 October 2026. Public repository.

[10] Pudim AI Project, *Keady Robin special inverse branch complete monotonicity remains open*, public frontier entry, repository commit `0867904e9718c33537197541da62c383b8734321`. Versioned frontier entry. Retrieved 31 October 2026.

[11] QEEDDD, *Verifiable Mathematical Reasoning Corpus*, public repository, commit `8219b3eb00b83a9bb81bc7b7afbc3c49b4538c9c` (29 October 2026). In particular manuscript [029], *Observation and Recovery (Geometry, Mixtures, and Spectral Inversion)*. Repository.