

# State-space models

FW 891

[Click here to view presentation online](#)

Christopher Cahill

6 November 2023



Quantitative Fisheries Center  
MICHIGAN STATE UNIVERSITY

# Purpose

- Goal
- Background
  - Some applications and history
- Dynamic linear model and some extensions
- Identifying nonidentifiability
- Future directions and other extensions
- R and Stan demo

# A key reference on ecological state space models:


## REVIEW

*Ecological Monographs*, 91(4), 2021, e01470

© 2021 The Authors. *Ecological Monographs* published by Wiley Periodicals LLC on behalf of Ecological Society of America

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

### A guide to state–space modeling of ecological time series

MARIE AUGER-MÉTHÉ <sup>1,2,13</sup> KEN NEWMAN,<sup>3,4</sup> DIANA COLE,<sup>5</sup> FANNY EMPACHER,<sup>6</sup> ROWENNA GRYBA,<sup>1,2</sup>  
AARON A. KING,<sup>7</sup> VIANEY LEOS-BARAJAS,<sup>8,9</sup> JOANNA MILLS FLEMMING,<sup>10</sup> ANDERS NIELSEN,<sup>11</sup> GIOVANNI PETRIS,<sup>12</sup>  
AND LEN THOMAS<sup>6</sup>

# What is a state space model

- State space models are a popular modeling framework for analyzing time-series data

# What is a state space model

- State space models are a popular modeling framework for analyzing time-series data
- Useful for modeling population dynamics, metapopulation dynamics, fisheries stock assessments, integrated population models, capture recapture data, animal movement, and biodiversity data

# What is a state space model

- State space models are a popular modeling framework for analyzing time-series data
- Useful for modeling population dynamics, metapopulation dynamics, fisheries stock assessments, integrated population models, capture recapture data, animal movement, and biodiversity data
- Quite popular because they directly model temporal autocorrelation in a way that helps differentiate process variation vs. observation error.

# What is a state space model

- A type of hierarchical model

# What is a state space model

- A type of hierarchical model
  - Hierarchical structure accomodates modeling two time series:



# What is a state space model

- A type of hierarchical model
  - Hierarchical structure accomodates modeling two time series:
    1. A state or process time series that is unobserved and attempts to reflect the true, but hidden, state of nature

# What is a state space model

- A type of hierarchical model
  - Hierarchical structure accomodates modeling two time series:
    1. A state or process time series that is unobserved and attempts to reflect the true, but hidden, state of nature
    2. An observation time series that consists of observations related to the state time series

# What is a state space model

- A type of hierarchical model
  - Hierarchical structure accomodates modeling two time series:
    1. A state or process time series that is unobserved and attempts to reflect the true, but hidden, state of nature
    2. An observation time series that consists of observations related to the state time series
- An example

# A historical aside

- The Kalman filter

# A historical aside

- The Kalman filter
  - A mathematical technique that removes “noise” from data

# A historical aside

- The Kalman filter
  - A mathematical technique that removes “noise” from data
  - Used to estimate and control the gradually changing condition of complex systems optimally from incomplete information

# A historical aside

- The Kalman filter
  - A mathematical technique that removes “noise” from data
  - Used to estimate and control the gradually changing condition of complex systems optimally from incomplete information
    - Think flight control systems

# A historical aside

- The Kalman filter
  - A mathematical technique that removes “noise” from data
  - Used to estimate and control the gradually changing condition of complex systems optimally from incomplete information
    - Think flight control systems
- Importance




# A historical aside

- The Kalman filter
  - A mathematical technique that removes “noise” from data
  - Used to estimate and control the gradually changing condition of complex systems optimally from incomplete information
    - Think flight control systems
  - Importance
  - Remarkably simple and elegant

# A historical aside



# A historical aside



<https://www.lancaster.ac.uk/stor-i-student-sites/jack-trainer/how-nasa-used-the-kalman-filter-in-the-apollo-program/>

# A historical aside



Rudolf Kalman receiving the National Medal of Science from Barack Obama: Photo NAE


# Other applications in ecology and resource management?

# Other applications in ecology and resource management?


- Hint



# Other applications in ecology and resource management?



# Other applications in ecology and resource management?





# A toy example: normal dynamic linear model

- Time series of univariate observations  $y_t$ 
  - Evenly spaced points in time  $t(t = 1, 2, \dots, T)$

# A toy example: normal dynamic linear model

- Time series of univariate observations  $y_t$ 
  - Evenly spaced points in time  $t(t = 1, 2, \dots, T)$
- Call the time series of states  $z_t$

# A toy example: normal dynamic linear model

- Time series of univariate observations  $y_t$ 
  - Evenly spaced points in time  $t(t = 1, 2, \dots, T)$
- Call the time series of states  $z_t$
- Process variance and observation error modeled using Gaussian distributions and both are modeled with linear equations

# A toy example: normal dynamic linear model

- Time series of univariate observations  $y_t$ 
  - Evenly spaced points in time  $t(t = 1, 2, \dots, T)$
- Call the time series of states  $z_t$
- Process variance and observation error modeled using Gaussian distributions and both are modeled with linear equations
- This is a normal dynamic linear model

# The toy example in math:

$$z_t = \alpha + z_{t-1} + \varepsilon_t, \quad \varepsilon_t \sim \text{N}(0, \sigma_p^2), \quad [\text{process equation}]$$

$$y_t = z_t + \eta_t, \quad \eta_t \sim \text{N}(0, \sigma_o^2). \quad [\text{observation equation}]$$

# The toy example in math:

$$z_t = \alpha + z_{t-1} + \varepsilon_t, \quad \varepsilon_t \sim \text{N}(0, \sigma_p^2), \quad [\text{process equation}]$$

$$y_t = z_t + \eta_t, \quad \eta_t \sim \text{N}(0, \sigma_o^2). \quad [\text{observation equation}]$$

- The time-series follows a random walk with drift or trend term  $\alpha$

# The toy example in math:

$$z_t = \alpha + z_{t-1} + \varepsilon_t, \quad \varepsilon_t \sim \text{N}(0, \sigma_p^2), \quad [\text{process equation}]$$

$$y_t = z_t + \eta_t, \quad \eta_t \sim \text{N}(0, \sigma_o^2). \quad [\text{observation equation}]$$

- The time-series follows a random walk with drift or trend term  $\alpha$
- $\sigma_p^2$  represents process variance
- $\sigma_o^2$  represents observation variance

# The toy example in math:

$$z_t = \alpha + z_{t-1} + \varepsilon_t, \quad \varepsilon_t \sim \text{N}(0, \sigma_p^2), \quad [\text{process equation}]$$

$$y_t = z_t + \eta_t, \quad \eta_t \sim \text{N}(0, \sigma_o^2). \quad [\text{observation equation}]$$

- The time-series follows a random walk with drift or trend term  $\alpha$
- $\sigma_p^2$  represents process variance
- $\sigma_o^2$  represents observation variance
- $\varepsilon_t$  represent process errors
- $\eta_t$  represent observation errors



# The toy example in pictures




FIG. 1. The dependence structure and evolution of the two time series comprising a simple univariate state-space model. (a) Dependence relationships with arrows, demonstrating that once the dependence of the observations  $y_t$  on the states  $z_t$  is accounted for, the observations are assumed independent. (b) Our toy model (Eqs. 1, 2). The blue and red dots are the simulated observations and states, respectively. The black line and gray band are the estimated states and associated 95% confidence intervals. The true states, but not the observations, usually fall in the 95% confidence intervals. This demonstrates that the state estimates can be a closer approximation of the truth than the observations.

# Two main assumptions

- State space models (SSMs) assume states evolve as a Markov process, generally as a first order process

# Two main assumptions

- State space models (SSMs) assume states evolve as a Markov process, generally as a first order process
  - For toy model, this means that the state at time  $t$ ,  $z_t$  depends only on the state at the previous time step  $z_{t-1}$

# Two main assumptions

- State space models (SSMs) assume states evolve as a Markov process, generally as a first order process
  - For toy model, this means that the state at time  $t$ ,  $z_t$  depends only on the state at the previous time step  $z_{t-1}$
- Second, SSMs assume observations are independent of one another after we account for their dependence on the states

# Two main assumptions

- State space models (SSMs) assume states evolve as a Markov process, generally as a first order process
  - For toy model, this means that the state at time  $t$ ,  $z_t$  depends only on the state at the previous time step  $z_{t-1}$
- Second, SSMs assume observations are independent of one another after we account for their dependence on the states
  - More formally, given some state  $z_t$ , each observation  $y_t$  is conditionally independent of all other observations

# Two main assumptions

- State space models (SSMs) assume states evolve as a Markov process, generally as a first order process
  - For toy model, this means that the state at time  $t$ ,  $z_t$  depends only on the state at the previous time step  $z_{t-1}$
- Second, SSMs assume observations are independent of one another after we account for their dependence on the states
  - More formally, given some state  $z_t$ , each observation  $y_t$  is conditionally independent of all other observations
  - Phrased differently, dependence between observations results from dependence between hidden states  $z_t$

# Two main assumptions

- State space models (SSMs) assume states evolve as a Markov process, generally as a first order process
  - For toy model, this means that the state at time  $t$ ,  $z_t$  depends only on the state at the previous time step  $z_{t-1}$
- Second, SSMs assume observations are independent of one another after we account for their dependence on the states
  - More formally, given some state  $z_t$ , each observation  $y_t$  is conditionally independent of all other observations
  - Phrased differently, dependence between observations results from dependence between hidden states  $z_t$
- In a pop dy context, this could be interpreted to mean that  $y_t$  values are autocorrelated because the underlying process driving them is autocorrelated through time

# The toy example needs one more thing:

$$z_t = \alpha + z_{t-1} + \varepsilon_t, \quad \varepsilon_t \sim \text{N}(0, \sigma_p^2), \quad [\text{process equation}]$$

$$y_t = z_t + \eta_t, \quad \eta_t \sim \text{N}(0, \sigma_o^2). \quad [\text{observation equation}]$$

- We must define the state at  $z_{t=0}$



# The toy example needs one more thing:

$$z_t = \alpha + z_{t-1} + \varepsilon_t, \quad \varepsilon_t \sim \text{N}(0, \sigma_p^2), \quad [\text{process equation}]$$

$$y_t = z_t + \eta_t, \quad \eta_t \sim \text{N}(0, \sigma_o^2). \quad [\text{observation equation}]$$

- We must define the state at  $z_{t=0}$ 
  - Typically referred to as the initialization equation

# The toy example needs one more thing:

$$z_t = \alpha + z_{t-1} + \varepsilon_t, \quad \varepsilon_t \sim \text{N}(0, \sigma_p^2), \quad [\text{process equation}]$$

$$y_t = z_t + \eta_t, \quad \eta_t \sim \text{N}(0, \sigma_o^2). \quad [\text{observation equation}]$$

- We must define the state at  $z_{t=0}$ 
  - Typically referred to as the initialization equation
- We'll set  $z_{t=0} = z_0$ , where  $z_0$  is an estimated parameter

# Handling nonlinearity

- Our toy example is useful for teaching but rarely helpful in practice

# Handling nonlinearity

- Our toy example is useful for teaching but rarely helpful in practice
- Some useful extensions are

# Handling nonlinearity

- Our toy example is useful for teaching but rarely helpful in practice
- Some useful extensions are

$$z_t = z_{t-1} \exp(\beta_0 + \beta_1 z_{t-1} + \varepsilon_t), \quad \varepsilon_t \sim \text{N}(0, \sigma_p^2) \quad [\text{process equation}]$$
$$y_t = z_t + \eta_t, \quad \eta_t \sim \text{N}(0, \sigma_o^2) . \quad [\text{observation equation}]$$

# Handling nonlinearity

- Our toy example is useful for teaching but rarely helpful in practice
- Some useful extensions are

$$z_t = z_{t-1} \exp(\beta_0 + \beta_1 z_{t-1} + \varepsilon_t), \quad \varepsilon_t \sim \text{N}(0, \sigma_p^2) \quad [\text{process equation}]$$
$$y_t = z_t + \eta_t, \quad \eta_t \sim \text{N}(0, \sigma_o^2) . \quad [\text{observation equation}]$$

- stochastic logistic model

# Handling nonlinearity

- Our toy example is useful for teaching but rarely helpful in practice
- Some useful extensions are

$$w_t = w_{t-1} + \beta_0 + \beta_1 \exp(w_{t-1}) + \varepsilon_t, \quad \varepsilon_t \sim \text{N}(0, \sigma_p^2) \quad [\text{process eq.}]$$

$$y_t = \exp(w_t) + \eta_t, \quad \eta_t \sim \text{N}(0, \sigma_o^2) \quad [\text{observation eq.}]$$

$$\text{where } w_t = \log(z_t)$$

- playing with math and modeling population on logarithmic scale

# Handling nonlinearity

- Our toy example is useful for teaching but rarely helpful in practice
- Some useful extensions are

$$z_t = z_{t-1} \exp(\beta_0 + \beta_1 \log(z_{t-1}) + \varepsilon_t), \quad \varepsilon_t \sim \text{N}(0, \sigma_p^2) \quad [\text{process eq.}]$$

- Gompertz model, which assumes that per-unit-abundance growth rate depends on log abundance



# Handling nonlinearity


- Our toy example is useful for teaching but rarely helpful in practice
- Some useful extensions are

$$w_t = \beta_0 + (1 + \beta_1) w_{t-1} + \varepsilon_t, \quad \varepsilon_t \sim \text{N}(0, \sigma_p^2), \quad [\text{process equation}]$$
$$g_t = w_t + \eta_t, \quad \eta_t \sim \text{N}(0, \sigma_o^2), \quad [\text{observation equation}]$$

where  $w_t = \log(z_t)$  and  $g_t = \log(y_t)$

- Linearized Gompertz model

# What's the point?



# State space models are complex

- e.g., see Thorson et al. 2016

# State space models are complex

- e.g., see Thorson et al. 2016
- Are there simpler alternatives?
  - SSMs superior to simpler models when both the process variance and the observation error are large (e.g., de Valpine and Hastings 2002)


# State space models are complex

- e.g., see Thorson et al. 2016
- Are there simpler alternatives?
  - SSMs superior to simpler models when both the process variance and the observation error are large (e.g., de Valpine and Hastings 2002)
  - Many studies have shown that SSMs provided better inference than simpler models

# State space models are complex

- e.g., see Thorson et al. 2016
- Are there simpler alternatives?
  - SSMs superior to simpler models when both the process variance and the observation error are large (e.g., de Valpine and Hastings 2002)
  - Many studies have shown that SSMs provided better inference than simpler models
- Should SSMs be the default for many ecological time series?

# Nonidentifiability



The Hunt-Lenox Globe: here be dragons

# Nonidentifiability

- Identifiability refers to whether there is a unique representation of the model




FIG. 3. Examples of log-likelihood profiles for a parameter  $\theta$  under various identifiability scenarios: (a) globally identifiable, (b) locally identifiable, and (c) nonidentifiable model.



# Nonidentifiability

- Identifiability refers to whether there is a unique representation of the model
  - Intrinsic vs. extrinsic nonidentifiability

# Nonidentifiability

- Identifiability refers to whether there is a unique representation of the model
  - Intrinsic vs. extrinsic nonidentifiability
- Misusing informed priors may hide identifiability issues

# Nonidentifiability

- Identifiability refers to whether there is a unique representation of the model
  - Intrinsic vs. extrinsic nonidentifiability
- Misusing informed priors may hide identifiability issues
- Diagnostics in the Bayesian framework

# Nonidentifiability

- Identifiability refers to whether there is a unique representation of the model
  - Intrinsic vs. extrinsic nonidentifiability
- Misusing informed priors may hide identifiability issues
- Diagnostics in the Bayesian framework
  - Correlation between parameters

# Nonidentifiability

- Identifiability refers to whether there is a unique representation of the model
  - Intrinsic vs. extrinsic nonidentifiability
- Misusing informed priors may hide identifiability issues
- Diagnostics in the Bayesian framework
  - Correlation between parameters
  - Simulation
    - If your model is non-identifiable, parameters are usually biased with large variances

# Nonidentifiability

- Identifiability refers to whether there is a unique representation of the model
  - Intrinsic vs. extrinsic nonidentifiability
- Misusing informed priors may hide identifiability issues
- Diagnostics in the Bayesian framework
  - Correlation between parameters
  - Simulation
    - If your model is non-identifiable, parameters are usually biased with large variances
- *One should choose priors with great care for SSMs*

# Remedies for nonidentifiability

- Reformulate the model
- Make simplifying assumptions when data are limited
- Estimate measurement errors externally
- Integrate additional data
- Try to use replicated observations
- Match temporal resolution between states and observations
  - If one has a data set with locations every 8 h, it would be challenging to estimate behavioral states lasting <16–24 h

# Wrap up

- SSMs are flexible models for time series that can be used to answer a broad range of ecological questions
- They can be used to model univariate or multivariate time series
- SSMs can be linear or nonlinear, and have discrete or continuous time steps
- Key point:



# Wrap up

- SSMs are flexible models for time series that can be used to answer a broad range of ecological questions
- They can be used to model univariate or multivariate time series
- SSMs can be linear or nonlinear, and have discrete or continuous time steps
- Key point:
  - Recognize that this is a crash course, and there are a great many extensions of SSMs in ecology

# To the example

# References

- Aeberhard et al. 2018. Review of state–space models for fisheries science. *Annual Review of Statistics and Its Application* 5:215–235.
- Auger-Methe et al. 2021. A guide to state-space modeling of ecological time series. *Ecological Monographs*.
- de Valpine and Hastings 2002. Fitting population models incorporating process noise and observation error. *Ecological Monographs*.
- Jamieson and Brooks 2004. Density dependence in North American ducks. *Animal Biodiversity and Conservation* 27:113–128.
- Thorson et al. 2016. Joint dynamic species distribution models: a tool for community ordination and spatio-temporal monitoring. *Global Ecology and Biogeography* 25:1144–1158.
- Walters 1986. *Adaptive Management of Renewable Resources*.