

Article
Talk

Read
View source
View history

Search

Heart

From Wikipedia, the free encyclopedia

*This article is about the internal organ. For other uses, see *Heart (disambiguation)*.*

The **heart** is a **muscular organ** in humans and other **animals**, which pumps **blood** through the **blood vessels** of the **circulatory system**.^[1] Blood provides the body with **oxygen** and **nutrients**, and also assists in the removal of **metabolic wastes**.^[2] The heart is located in the middle compartment of the **mediastinum** in the **chest**.^[3]

In humans, other mammals, and birds, the heart is divided into four chambers: upper left and right **atria**; and lower left and right **ventricles**.^{[4][5]} Commonly the right atrium and ventricle are referred together as the *right heart* and their left counterparts as the *left heart*.^[6] Fish in contrast have two chambers, an atrium and a ventricle, while reptiles have three chambers.^[5] In a healthy heart blood flows one way through the heart due to **heart valves**, which prevent **backflow**.^[3] The heart is enclosed in a protective sac, the **pericardium**, which also contains a small amount of **fluid**. The wall of the heart is made up of three layers: **epicardium**, **myocardium**, and **endocardium**.^[7]

The heart pumps blood through the body. Blood low in oxygen from the **systemic circulation** enters the right atrium from the **superior** and **inferior venae cavae** and passes to the right ventricle. From here it is pumped into the **pulmonary circulation**, through the **lungs** where it receives oxygen and gives off carbon dioxide. Oxygenated blood then returns to the left atrium, passes through the left ventricle and is pumped out through the **aorta** to the systemic circulation—where the oxygen is used and **metabolized** to **carbon dioxide**.^[8] In addition the blood carries nutrients from the **digestive tract** to various organs of the body, while transporting **waste** to the liver and **kidneys**. Normally with each **heartbeat** the right ventricle pumps the same amount of blood into the lungs as the left ventricle pumps to the body. **Veins** transport blood to the heart and carry deoxygenated blood - except for the pulmonary and **portal veins**. **Arteries** transport blood away from the heart, and apart from the pulmonary artery hold oxygenated blood. Their increased distance from the heart cause veins to have lower **pressures** than arteries.^{[2][3]} The heart contracts at a resting **rate** close to 72 beats per minute.^[9] **Exercise** temporarily increases the rate, but lowers **resting heart rate** in the long term, and is good for heart health.^[10]

Cardiovascular diseases (CVD) are the most common cause of death globally as of 2008, accounting for 30% of deaths.^{[11][12]} Of these more than three quarters follow **coronary artery disease** and **stroke**.^[11] Risk factors include: **smoking**, being **overweight**, little exercise, **high cholesterol**, **high blood pressure**, and poorly controlled **diabetes**, among others.^[13] Diagnosis of CVD is often done by **listening** to the **heart-sounds** with a **stethoscope**, **ECG** or by **ultrasound**.^[3] Specialists who focus on diseases of the heart are called **cardiologists**, although many specialties of medicine may be involved in treatment.^[12]

Contents

- 1

Structure
- 1.1

Location and shape
- 1.2

Chambers
- 1.3

Heart wall
- 1.4

Coronary circulation
- 1.5

Nerve supply
- 2

Development
- 3

Physiology
- 3.1

Blood flow
- 3.2

Electrical conduction
- 3.3

Heart rate
- 3.4

Heart sounds
- 4

Clinical significance
- 4.1

Disease
- 4.2

Diagnosis
- 4.3

Treatment
- 5

History
- 5.1

Ancient
- 5.2

Pre-modern
- 5.3

Modern
- 6

Society and culture
- 6.1

Symbolism
- 6.2

Food
- 7

Other animals
- 7.1

Double circulatory systems
- 7.2

The fully divided heart
- 7.3

Fish
- 7.4

Invertebrates
- 8

Additional images
- 9

Notes
- 10

References

Heart

The human heart

Details

System
Circulatory

Artery
Aorta,^[a] **pulmonary trunk** and right and left **pulmonary arteries**^[b]
Right coronary artery, left main coronary artery^[c]

Vein
Superior vena cava , inferior vena cava,^[d] right and left **pulmonary veins**,^[e] great cardiac vein , middle cardiac vein , small cardiac vein , anterior cardiac veins.^[f]

Nerve
Accelerans nerve , Vagus nerve

Identifiers

Latin
cor

Greek
kardía (*καρδία*)

MeSH
A07.541

TA
A12.1.00.001

FMA
7088

Anatomical terminology

[edit on Wikidata]

Normal heart sounds

Normal heart sounds as heard with a **stethoscope**

*Problems playing this file? See *media help*.*

Main page
Current events
Random article
Donate to Wikipedia
Wikipedia store

Interaction

Help
About Wikipedia
Community portal
Recent changes
Contact page

Tools

What links here
Related changes
Upload file
Special pages
Permanent link
Page information
Wikidata item
Cite this page

Print/export

Create a book
Download as PDF
Printable version

In other projects

Wikimedia Commons
Wikiquote

Languages

Afrikaans
Alemannisch
አማርኛ
/English
العربية
Aragonés
ܐܪܡܝܐ
Armãneashti
অসমীয়া
Asturianu
Avañe'ê
Awap
Aymar aru
Azərbaycanca
Башҡортса
Беларуская
Беларуская (тарашкевіца)
Bikol Central
Български
Bosanski
Brezhoneg
Буряад
Català
Чӕвашла
Cebuano
Čeština
Chavacano de Zamboanga
ChiShona
Corsu
Cymraeg
Dansk
Deutsch
𐌆𐌿𐌺𐌹𐌸𐌰
Eesti
Ελληνικά
Emiliàn e rumagnòl
Español
Esperanto
Euskara
فارسی
Fiji Hindi
Føroyskt
Français
Frysk
Gaeilge
Galego
客家語/Hak-kâ-ngî
Хальмг
한국어
Հայերեն
हिन्दी

- Hrvatski
- Ido
- Ilokano
- Bahasa Indonesia
- Interlingua
- Interlingue
- Íslenska
- Italiano
- עברית
- Basa Jawa
- ■

■

■

■
- Kapampangan
- ქართული
- Қазақша
- Kiswahili
- Кomi
- Kreyòl ayisyen
- Кыргызча
- Кырык мары
- Лакку
- Latina
- Latviešu
- Lëtzebuergesch
- Lietuvių
- Limburgs
- Lingála
- La .lojban.
- Lumbaart
- Magyar
- Македонски
- മലയാളം
- Malti
- मराठी
- مصرى
- Bahasa Melayu
- Mìng-dĕ̍ng-ngṳ̂
- Монгол
- მწკჷმაშუალო
- Nāhuatl
- Nederlands
- नेपाली
- नेपाल भाषा
- 日本語
- Napulitano
- Нохчийн
- Norsk bokmål
- Norsk nynorsk
- Nouormand
- Occitan
- Олык марий
- ଓଡ଼ିଆ
- Oʻzbekcha/Ўзбекча
- ਪੰਜਾਬੀ
- Pangasinan
- پنجابی
- پښتو
- Plattdüütsch
- Polski
- Português
- Română
- Runa Simi
- Русиньскый
- Русский
- Саха тыла
- संस्कृतम्
- Sardu
- Scots
- Shqip
- Sicilianu
- සිංහල
- Simple English
- Slovenčina
- Slovenščina
- Ślůnski
- Soomaaliga
- کوردی ناوەندی
- Српски / srpski
- Srpskohrvatski / српскохрватски
- Basa Sunda
- Suomi
- Svenska
- Tagalog
- தமிழ்
- Татарча/tatarça
- ■

■

■

■
- ᩋᩣ᩠᩵ᨶᩣ᩠᩵ᨶ
- ትግርኛ
- Тоҷикӣ
- Türkçe
- Українська
- اردو
- ئۇيغۇرچە / Uyghurche
- Vahcuengh
- Vèneto
- Vepsän kel'
- Tiếng Việt
- Võro
- West-Vlams
- Winaray
- יידיש
- Yorùbá


- 11 Bibliography
- 12 External links

Structure


Location and shape

The human heart is situated in the **middle mediastinum** , at the level of **thoracic vertebrae T5–T8**. A double-membraned sac called the **pericardium** surrounds the heart and attaches to the mediastinum.^[15] The back surface of the heart lies near the **vertebral column**, and the front surface sits behind to the **sternum** and **rib cartilages**.^[7] The upper part of the heart is the attachment point for several large blood vessels - the **venae cavae**, **aorta** and **pulmonary trunk** . The upper part of the heart is located at the level of the third costal cartilage.^[7] The lower tip of the heart, the apex, lies to the left of the sternum (8 to 9 cm from the **midsternal line**) between the junction of the fourth and fifth ribs near their **articulation** with the costal cartilages.^[7]


The largest part of the heart is usually slightly offset to the left side of the chest (though occasionally it may be **offset to the right** ) and is felt to be on the left because the **left heart** is stronger and larger, since it pumps to all body parts. Because the heart is between the **lungs**, the left lung is smaller than the right lung and has a cardiac notch in its border to accommodate the heart.^[7] The heart is cone-shaped, with its base positioned upwards and tapering down to the apex.^[7] An adult heart has a mass of 250–350 grams (9–12 oz).^[16] The heart is typically the size of a fist: 12 cm (5 in) in length, 8 cm (3.5 in) wide, and 6 cm (2.5 in) in thickness.^[7] Well-trained **athletes** can have much larger hearts due to the effects of exercise on the heart muscle, similar to the response of skeletal muscle.^[7]



The human heart is in the middle of the **thorax**, with its apex pointing to the left.^[14]



Computer generated animation of a beating human heart



Play media
Real-time MRI of the human heart

Chambers

The heart has four chambers, two upper **atria**, the receiving chambers, and two lower **ventricles**, the discharging chambers. The atria open into the ventricles via the atrioventricular valves, present in the **atrioventricular septum**. This distinction is visible also on the surface of the heart as the **coronary sulcus**.^[17] There is an ear-shaped structure in the upper right atrium called the right atrial appendage, or auricle, and another in the upper left atrium, the left atrial appendage.^[18] The right atrium and the right ventricle together are sometimes referred to as the *right heart*. Similarly, the left atrium and the left ventricle together are sometimes referred to as the *left heart*.^[6] The ventricles are separated from each other by the **interventricular septum**, visible on the surface of the heart as the **anterior longitudinal sulcus** and the **posterior interventricular sulcus**.^[17]

The **cardiac skeleton** is made of **dense connective tissue** and this gives structure to the heart. It forms the **atrioventricular septum** which separates the atria from the ventricles, and the fibrous rings which serve as bases for the four **heart valves**.^[19] The cardiac skeleton also provides an important boundary in the heart's electrical conduction system since collagen cannot conduct electricity. The interatrial septum separates the atria and the interventricular septum separates the ventricles.^[7] The interventricular septum is much thicker than the interatrial septum, since the ventricles need to generate greater pressure when they contract.^[7]

Valves

*Main article: **Heart valves***

The heart has four valves, which separate its chambers.^[7] The valves ensure blood flows in the correct direction through the heart and prevents backflow. Each valve consists of two to three cusps. The valves between the atria and ventricles connected to cartilaginous strings called **chordae tendinae** which in turn connect to muscles on the heart wall called **papillary muscles**.


The valves between the atria and ventricles are called the atrioventricular valves. Between the right atrium and the right ventricle is the **tricuspid valve**. The tricuspid valve has three cusps,^[20] which connect to chordae tendinae and three papillary muscles named the anterior, posterior, and septal muscles, after their relative positions.^[20] The **mitral valve** lies between the left atrium and left ventricle. It is also known as the bicuspid valve due to its having two cusps, an anterior and a posterior cusp. These cusps are also attached via chordae tendinae to two papillary muscles projecting from the ventricular wall.^[21]

The **papillary muscles** extends from the walls of the heart to the chordae tendinae of valves. These muscles prevent the valves from falling too far back when they close.^[22] During the relaxation phase of the cardiac cycle, the papillary muscles are also relaxed and the tension on the chordae tendineae is slight. As the heart chambers contract, so do the papillary muscles. This creates tension on the chordae tendineae, helping to hold the cusps of the atrioventricular valves in place and preventing them from being blown back into the atria.^[7] ^[g]^[20]

Two additional semilunar valves sit at the exit of each of the ventricles. The **pulmonary valve** is located at the base of the **pulmonary artery**. This has three cusps which are not attached to any papillary muscles. When the ventricle relaxes blood flows back into the ventricle from the artery and this



With the atria and major vessels removed, all four valves are clearly visible.^[7]



The heart, showing valves, arteries and veins. The white arrows shows the normal direction of blood flow.

flow of blood fills the pocket-like valve, pressing against the cusps which close to seal the valve. The semilunar **aortic valve** is at the base of the **aorta** and also is not attached to papillary muscles. This too has three cusps which close with the pressure of the blood flowing back from the aorta.^[7]

Right heart

The right heart consists of two chambers, the right atrium and the right ventricle, separated by a valve, the **tricuspid valve**.^[7]

The right atrium receives blood almost continuously from the body's two major veins, the **superior** and **inferior venae cavae**. A small amount of blood from the coronary circulation also drains into the right atrium via the **coronary sinus**, which is immediately above and to the middle of the opening of the inferior vena cava. ^[7] In the wall of the right atrium is an oval-shaped depression known as the **fossa ovalis**, which is a remnant of an opening in the fetal heart known as the **foramen ovale**.^[7] Most of the internal surface of the right atrium is smooth, the depression of the fossa ovalis is medial, and the anterior surface has prominent ridges of **pectinate muscles**, which are also present in the **right atrial appendage**.^[7]

The right atrium is connected to the right ventricle by the **tricuspid valve**.^[7] The walls of the right ventricle are lined with **trabeculae carneae**, ridges of cardiac muscle covered by endocardium. In addition to these muscular ridges, a band of cardiac muscle, also covered by endocardium, known as the **moderator band** reinforces the thin walls of the right ventricle and plays a crucial role in cardiac conduction. It arises from the lower part of the interventricular septum and crosses the interior space of the right ventricle to connect with the inferior papillary muscle.^[7] The right ventricle tapers into the **pulmonary trunk**, into which it ejects blood when contracting. The pulmonary trunk branches into the left and right pulmonary arteries that carry the blood to each lung. The pulmonary valve lies between the right heart and the pulmonary trunk.^[7]

Left heart

The left heart has two chambers: the left atrium, and the left ventricle, separated by the **mitral valve**.^[7]

The left atrium receives oxygenated blood back from the lungs via one of the four **pulmonary veins**. The left atrium has an outpouching called the **left atrial appendage**. Like the right atrium, the left atrium is lined by **pectinate muscles**.^[23] The left atrium is connected to the left ventricle by the **mitral valve**.^[7]

The left ventricle is much thicker as compared with the right, due to the greater force needed to pump blood to the entire body. Like the right ventricle, the left also has **trabeculae carneae**, but there is no **moderator band**. The left ventricle pumps blood to the body through the **aortic valve** and into the **aorta**. Two small openings above the aortic valve carry blood to the heart itself, the **left main coronary artery** and the **right coronary artery**.^[7]

Heart wall

Further information: **Cardiac muscle**

The heart wall is made up of three layers: the inner **endocardium**, middle **myocardium** and outer **epicardium**. These are surrounded by a double-membraned sac called the **pericardium**.

The innermost layer of the heart is called the endocardium. It is made up of a lining of **simple squamous epithelium**, and covers heart chambers and valves. It is continuous with the **endothelium** of the veins and arteries of the heart, and is joined to the myocardium with a thin layer of connective tissue.^[7] The endocardium, by secreting **endothelins**, may also play a role in regulating the contraction of the myocardium.^[7]

The middle layer of the heart wall is the myocardium, which is the **cardiac muscle** – a layer of involuntary **striated muscle tissue** surrounded by a framework of **collagen**. The cardiac muscle pattern is elegant and complex, as the muscle cells swirl and spiral around the chambers of the heart, with the outer muscles forming a figure 8 pattern around the atria and around the bases of the great vessels, and inner muscles forming a figure 8 around the two ventricles and proceed toward the apex. This complex swirling pattern allows the heart to pump blood more effectively.^[7]

There are two types of cells in cardiac muscle: **muscle cells** which have the ability to contract easily, and **pacemaker cells** of the conducting system. The muscle cells make up the bulk (99%) of cells in the atria and ventricles. These contractile cells are connected by **intercalated discs** which allow a rapid response to impulses of **action potential** from the pacemaker cells. The intercalated discs allow the cells to act as a **syncytium** and enable the contractions that pump blood through the heart and into the **major arteries**.^[7] The pacemaker cells make up 1% of cells and form the conduction system of the heart. They are generally much smaller than the contractile cells and have few **myofibrils** which gives them limited contractibility. Their function is similar in many respects to **neurons**.^[7] Cardiac muscle tissue has **autorhythmicity**, the unique ability to initiate a **cardiac action potential** at a fixed rate – spreading the impulse rapidly from cell to cell to trigger the contraction of the entire heart.^[7]

The pericardium surrounds the heart. It consists of two membranes: an inner **serous membrane** called the epicardium, and an outer fibrous membrane. Blood vessels and nerves reach the cardiac muscle from the epicardium.^[7] These help influence the **heart rate**.^[7] These enclose the **pericardial cavity** which contains the **pericardial fluid** that lubricates the surface of the heart. ^[24]

Coronary circulation

Main article: **Coronary circulation**

Heart tissue, like all cells in the body, needs to be supplied with **oxygen**, **nutrients** and a way of removing **metabolic wastes**. This is achieved by the **coronary circulation**, which includes **arteries**, **veins**, and **lymphatic vessels**. Blood circulates through the coronary circulation cyclically, in peaks and troughs relating to the heart muscle's relaxation or contraction.^[7]

Heart tissue receives blood from two arteries which arise just above the aortic valve. These are the **left main coronary artery** and the **right coronary artery**. The left main coronary artery splits shortly after leaving the **aorta** into two vessels, the **left anterior descending** and the **left circumflex artery**. The left anterior descending artery supplies heart tissue and the front, outer side, and the septum of the left ventricle. It does this by smaller branching arteries - diagonal and septal branches. The left circumflex supplies the back and underneath of the left ventricle. The right coronary artery supplies the right atrium, right ventricle, and lower posterior sections of the left ventricle. The right coronary artery also supplies blood to the atrioventricular node (in about 90% of people) and the sinoatrial node (in about 60% of people). The right coronary artery runs in a

Frontal section showing papillary muscles attached to the tricuspid valve on the right and to the mitral valve on the left via chordae tendineae.^[7]

Layers of the heart wall, including visceral and parietal pericardium.

The swirling pattern of myocardium helps the heart pump effectively

Arterial supply to the heart (red), with other areas labelled (blue).

groove at the back of the heart and the left anterior descending artery runs in a groove at the front. There is significant variation between people in the anatomy of the arteries that supply the heart ^[25] The arteries divide at their furthest reaches into smaller branches that join together at the edges of each arterial distribution. ^[7]

The **coronary sinus** is a large vein that drains into the right atrium, and receives most of the venous drainage of the heart. It receives blood from the **great cardiac vein** (receiving the left atrium and both ventricles), the **posterior cardiac vein** (draining the back of the left ventricle), the **middle cardiac vein** (draining the bottom of the left and right ventricles), and **small cardiac veins**.^[26] The **anterior cardiac veins** drain the front of the right ventricle and drain directly into the right atrium. ^[7]

Small lymphatic networks called **plexuses** exist beneath each of the three layers of the heart. These networks collect into a main left and a main right trunk, which travel up the groove between the ventricles that exists on the heart's surface, receiving smaller vessels as they travel up. These vessels then travel into the atrioventricular groove, and receive a third vessel which drains the section of the left ventricle sitting on the diaphragm. The left vessel joins with this third vessel, and travels along the pulmonary artery and left atrium, ending in the **inferior tracheobronchial node**. The right vessel travels along the right atrium and the part of the right ventricle sitting on the diaphragm. It usually then travels in front of the ascending aorta and then ends in a **brachiocephalic node**.^[27]

Nerve supply

The heart receives nerves from the **sympathetic trunk** and the **vagus nerve**. These nerves act to influence, but not control, the heart rate. **Sympathetic nerves** also influence the force of heart contraction. ^[28] Signals that travel along these nerves arise from two paired **cardiovascular centres** in the **medulla oblongata**. The **vagus nerve** of the parasympathetic nervous system acts to decrease the heart rate, and nerves from the **sympathetic trunk** act to increase the heart rate. ^[7] These nerves form a network of nerves that lies over the heart called the **cardiac plexus**.^{[7][27]}

The vagus nerve is a long, wandering nerve that emerges from the **brainstem** and provides parasympathetic stimulation to a large number of organs in the thorax and abdomen, including the heart.^[29] The nerves from the sympathetic trunk emerge through the T1-T4 **thoracic ganglia** and travel to both the sinoatrial and atrioventricular nodes, as well as to the atria and ventricles. The ventricles are more richly innervated by sympathetic fibers than parasympathetic fibers. Sympathetic stimulation causes the release of the neurotransmitter **norepinephrine** (also known as **noradrenaline**) at the **neuromuscular junction** of the cardiac nerves. This shortens the repolarization period, thus speeding the rate of depolarization and contraction, which results in an increased heart rate. It opens chemical or ligand-gated sodium and calcium ion channels, allowing an influx of **positively charged ions**.^[7] Norepinephrine binds to the **beta-1 receptor**.^[7]

Development

Main articles: [Heart development](#) and [Human embryogenesis](#)

The heart is the first functional organ to develop and starts to beat and pump blood at about three weeks into **embryogenesis**. This early start is crucial for subsequent embryonic and **prenatal development**.

The heart derives from **splanchnopleuric mesenchyme** in the neural plate which forms the cardiogenic region.

Two **endocardial tubes** form here that fuse to form a primitive heart tube known as the **tubular heart**.^[30] Between the third and fourth week, the heart tube lengthens, and begins to fold to form an S-shape within the pericardium. This places the chambers and major vessels into the correct alignment for the developed heart. Further development will include the septa and valves formation and remodelling of the heart chambers. By the end of the fifth week the septa are complete and the heart valves are completed by the ninth week. ^[7]

Before the fifth week, there is an opening in the fetal heart known as the **foramen ovale**. The foramen ovale allowed blood in the fetal heart to pass directly from the right atrium to the left atrium, allowing some blood to bypass the lungs. Within seconds after birth, a flap of tissue known as the **septum primum** that previously acted as a valve closes the foramen ovale and establishes the typical cardiac circulation pattern. A depression in the surface of the right atrium remains where the foramen ovale once walls, called the fossa ovalis.^[7]

The **embryonic** heart begins beating at around 22 days after conception (5 weeks after the last normal menstrual period, LMP). It starts to beat at a rate near to the mother's which is about 75–80 **beats per minute** (bpm). The embryonic heart rate then accelerates and reaches a peak rate of 165–185 bpm early in the early 7th week (early 9th week after the LMP).^{[31][32]} After 9 weeks (start of the **fetal** stage) it starts to decelerate, slowing to around 145 (±25) bpm at birth. There is no difference in female and male heart rates before birth.^[33]

Physiology

Main article: [Cardiac physiology](#)

Blood flow

The heart functions as a pump in the **circulatory system** to provide a continuous **flow of blood** throughout the body. This circulation consists of the **systemic circulation** to and from the body and the **pulmonary circulation** to and from the lungs. Blood in the pulmonary circulation exchanges carbon dioxide for oxygen in the lungs through the process of **respiration**. The systemic circulation then transports oxygen to the body and returns carbon dioxide and relatively deoxygenated blood to the heart for transfer to the lungs. ^[7]

The **right heart** collects deoxygenated blood from two large veins, the **superior** and **inferior venae cavae**. Blood collects in the right and left atrium continuously. ^[7] The superior vena cava drains blood from above the **diaphragm** and empties into the upper back part of the right atrium. The inferior vena cava drains the blood from below the diaphragm and empties into the back part of the atrium below the opening for the superior vena cava.

Autonomic innervation of the heart

Development of the human heart during the first eight weeks (top), and the formation of the heart chambers (bottom).^[7]

Blood flow through the valves

Immediately above and to the middle of the opening of the inferior vena cava is the opening of the thin-walled coronary sinus.^[7] Additionally, the [coronary sinus](#) returns deoxygenated blood from the myocardium to the right atrium. The blood collects in the right atrium. When the right atrium contracts, the blood is pumped through the [tricuspid valve](#) into the right ventricle. As the right ventricle contracts, the tricuspid valve closes and the blood is pumped into the pulmonary trunk through the [pulmonary valve](#). The pulmonary trunk divides into pulmonary arteries and progressively smaller arteries throughout the lungs, until it reaches [capillaries](#). As these pass by [alveoli](#) carbon dioxide is [exchanged](#) for oxygen. This happens through the passive process of [diffusion](#).

In the **left heart**, oxygenated blood is returned to the left atrium via the [pulmonary veins](#). It is then pumped into the left ventricle through the [mitral valve](#) and into the aorta through the [aortic valve](#) for systemic circulation. The aorta is a large artery that branches into many smaller arteries, [arterioles](#), and ultimately [capillaries](#). In the capillaries, oxygen and nutrients from blood are supplied to body cells for metabolism, and exchanged for carbon dioxide and waste products.^[7] Capillary blood, now deoxygenated, travels into [venules](#) and [veins](#) that ultimately collect in the superior and inferior vena cavae, and into the right heart.

Cardiac cycle

Main articles: [Cardiac cycle](#), [Systole](#), and [Diastole](#)

The **cardiac cycle** refers to a complete heartbeat which includes [systole](#) and [diastole](#) and the intervening pause.^[9] The cycle begins with contraction of the atria and ends with relaxation of the ventricles. Systole refers to contraction of the atria or ventricles of the heart. Diastole is when the atria or ventricles relax and fill with blood. The atria and ventricles work in concert, so in systole when the ventricles are contracting, the atria are relaxed and collecting blood. When the ventricles are relaxed in diastole, the atria contract to pump blood to the ventricles. This coordination ensures blood is pumped efficiently to the body.^[7]

At the beginning of the cardiac cycle, in early diastole, both the atria and ventricles are relaxed. Since blood moves from areas of high pressure to areas of low pressure, when the chambers are relaxed, blood will flow into the atria (through the [coronary sinus](#) and the pulmonary veins). As the atria begin to fill, the pressure will rise so that the blood will move from the atria into the ventricles. In late diastole the atria contract, pumping more blood into the ventricles. This causes a rise in pressure in the ventricles. As the ventricles reach systole, blood will be pumped into the pulmonary artery (right ventricle) or aorta (left ventricle).^[9]

When the atrioventricular valves (tricuspid and mitral) are open, during blood flow to the ventricles, the aortic and pulmonary valves are closed to prevent [backflow](#) into the ventricles. When the ventricular pressure is greater than the atrial pressure the tricuspid and mitral valves will shut. When the ventricles contract the pressure forces the aortic and pulmonary valves open. As the ventricles relax, the aortic and pulmonary valves will close in response to decreased pressure.^[9]

Cardiac output

Main article: [Cardiac output](#)

Cardiac output (CO) is a measurement of the amount of blood pumped by each ventricle (stroke volume) in one minute. This is calculated by multiplying the stroke volume (SV) by the beats per minute of the heart rate (HR). So that: $CO = SV \times HR$.^[7] The cardiac output is normalized to body size through [body surface area](#) and is called the [cardiac index](#).

The average cardiac output, using an average stroke volume of about 70mL, is 5.25 L/min, with a normal range of 4.0–8.0 L/min.^[7] The stroke volume is normally measured using an [echocardiogram](#) and can be influenced by the size of the heart, physical and mental condition of the individual, [sex](#), [contractility](#), duration of contraction, [preload](#) and [afterload](#).^[7]

Preload refers to the filling pressure of the atria at the end of diastole, when they are at their fullest. A main factor is how long it takes the ventricles to fill—if the ventricles contract faster, then there is less time to fill and the preload will be less.^[7] Preload can also be affected by a person's blood volume. The force of each contraction of the heart muscle is proportional to the preload, described as the [Frank-Starling mechanism](#).

This states that the force of contraction is directly proportional to the initial length of muscle fiber, meaning a ventricle will contract more forcefully, the more it is stretched.^{[7][34]}

Afterload, or how much pressure the heart must generate to eject blood at systole, is influenced by [vascular resistance](#). It can be influenced by narrowing of the heart valves ([stenosis](#)) or contraction or relaxation of the peripheral blood vessels.^[7]

The strength of heart muscle contractions controls the stroke volume. This can be influenced positively or negatively by agents termed [inotropes](#). These can be either conditions or drugs. Positive inotropes that cause stronger contractions include [high blood calcium](#) and drugs such as [Digoxin](#), which will act to stimulate the [sympathetic nerves](#) in the [fight-or-flight](#) response. Negative inotropes causing weaker contractions include [high blood potassium](#), [hypoxia](#), [acidosis](#), and drugs such as [beta blockers](#) and [calcium channel blockers](#).

Electrical conduction

Main articles: [Electrical conduction system of the heart](#) and [Heart rate](#)

The normal rhythmical heart beat, called [sinus rhythm](#), is established by the [sinoatrial node](#), the heart's [pacemaker](#). Here an electrical signal is created that travels through the heart, causing the heart muscle to contract.

The sinoatrial node is found in the upper part of the [right atrium](#) near to the junction with the superior vena cava.^[35] The electrical signal generated by the sinoatrial node travels through the right atrium in a radial way that is not completely understood. It travels to the left atrium via [Bachmann's bundle](#), such that the muscles of the left and right atria contract together. ^{[36][37][38]} The signal then travels to the [atrioventricular node](#). This is found at the bottom of the right atrium in the [atrioventricular septum](#)—the boundary between the right atrium and the left ventricle. The septum is part of the [cardiac skeleton](#), tissue within the heart that the electrical signal cannot pass through, which forces the signal to pass through the atrioventricular node only.^[7] The signal then travels along the [bundle of His](#) to left and right [bundle branches](#) through to the ventricles of the heart. In the ventricles the signal is carried by specialized tissue called the [Purkinje fibers](#) which then transmit the electric charge to the heart muscle.^[39]

Heart rate

[Play media](#)

Blood flow through the heart from the [Khan academy](#)

The cardiac cycle as correlated to the ECG

The x-axis reflects time with a recording of the heart sounds. The y-axis represents pressure.^[7]

Main article: [Heart rate](#)

The normal resting heart rate is called the [sinus rhythm](#), created and sustained by the [sinoatrial node](#), a group of pacemaking cells found in the wall of the right atrium. Cells in the sinoatrial node do this by creating an [action potential](#). The [cardiac action potential](#) is created by the movement of specific [electrolytes](#) into and out of the pacemaker cells. The action potential then spreads to nearby cells.^[40]

When the sinoatrial cells are resting, they have a negative charge on their membranes. However a rapid influx of [sodium](#) ions causes the membrane's charge to become positive. This is called [depolarisation](#) and occurs spontaneously.^[7] Once the cell has a sufficiently high charge, the sodium channels close and [calcium](#) ions then begin to enter the cell, shortly after which [potassium](#) begins to leave it. All the ions travel through [ion channels](#) in the membrane of the sinoatrial cells. The potassium and calcium only start to move out of and into the cell once it has a sufficiently high charge, and so are called [voltage-gated](#). Shortly after this, the calcium channels close and [potassium channels](#) open, allowing potassium to leave the cell. This causes the cell to have a negative resting charge and is called [repolarization](#). When the membrane potential reaches approximately −60 mV, the potassium channels close and the process may begin again.^[7]

The ions move from areas where they are concentrated to where they are not. For this reason sodium moves into the cell from outside, and potassium moves from within the cell to outside the cell. Calcium also plays a critical role. Their influx through slow channels means that the sinoatrial cells have a prolonged "plateau" phase when they have a positive charge. A part of this is called the [absolute refractory period](#). Calcium ions also combine with the regulatory protein [troponin C](#) in the [troponin complex](#) to enable [contraction](#) of the cardiac muscle, and separate from the protein to allow relaxation.^[41]

The adult resting heart rate ranges from 60 to 100 bpm. The resting heart rate of a [newborn](#) can be 129 beats per minute (bpm) and this gradually decreases until maturity.^[42] An athlete's heart rate can be lower than 60 bpm. During exercise the rate can be 150 bpm with maximum rates reaching from 200 to 220 bpm.^[7]

Influences

The normal [sinus rhythm](#) of the heart, giving the resting [heart rate](#), is influenced a number of factors. The cardiovascular centres in the brainstem that control the sympathetic and parasympathetic influences to the heart through the vagus nerve and sympathetic trunk.^[43] These cardiovascular centres receive input from a series of receptors including [baroreceptors](#), sensing stretch the stretching of blood vessels and [chemoreceptors](#), sensing the amount of oxygen and carbon dioxide in the blood and its pH. Through a series of reflexes these help regulate and sustain blood flow.^[7]

Baroreceptors are stretch receptors located in the [aortic sinus](#), [carotid bodies](#), the venae cavae, and other locations, including pulmonary vessels and the right side of the heart itself. Baroreceptors fire at a rate determined by how much they are stretched,^[44] which is influenced by blood pressure, level of physical activity, and the relative distribution of blood. With increased pressure and stretch, the rate of baroreceptor firing increases, and the cardiac centers decrease sympathetic stimulation and increase parasympathetic stimulation. As pressure and stretch decrease, the rate of baroreceptor firing decreases, and the cardiac centers increase sympathetic stimulation and decrease parasympathetic stimulation.^[7] There is a similar reflex, called the atrial reflex or [Bainbridge reflex](#), associated with varying rates of blood flow to the atria. Increased venous return stretches the walls of the atria where specialized baroreceptors are located. However, as the atrial baroreceptors increase their rate of firing and as they stretch due to the increased blood pressure, the cardiac center responds by increasing sympathetic stimulation and inhibiting parasympathetic stimulation to increase heart rate. The opposite is also true.^[7] Chemoreceptors present in the carotid body or adjacent to the aorta in an aortic body respond to the blood's oxygen, carbon dioxide levels. Low oxygen or high carbon dioxide will stimulate firing of the receptors.^[45]

Exercise and fitness levels, age, body temperature, [basal metabolic rate](#), and even a person's emotional state can all affect the heart rate. High levels of the hormones [epinephrine](#), norepinephrine, and [thyroid hormones](#) can increase the heart rate. The levels of electrolytes including calcium, potassium, and sodium can also influence the speed and regularity of the heart rate; [low blood oxygen](#), low [blood pressure](#) and [dehydration](#) may increase it.^[7]

Heart sounds

Main article: [Heart sounds](#)

One of the simplest methods of assessing the heart's condition is to [listen](#) to it using a [stethoscope](#).^[7] Typically, healthy hearts have only two audible [heart sounds](#), called S1 and S2. The first heart sound S1, is the sound created by the closing of the atrioventricular valves during ventricular contraction and is normally described as "lub". The second heart sound, S2, is the sound of the semilunar valves closing during ventricular diastole and is described as "dub".^[7] Each sound consists of two components, reflecting the slight difference in time as the two valves close.^[46] S2 may [split](#) into two distinct sounds, either as a result of inspiration or different valvular or cardiac problems.^[46] Additional heart sounds may also be present and these give rise to [gallop rhythms](#). A [third heart sound](#), S3 usually indicates an increase in ventricular blood volume. A [fourth heart sound](#) S4 is referred to as an atrial gallop and is produced by the sound of blood being forced into a stiff ventricle. The combined presence of S3 and S4 give a quadruple gallop.^[7]

[Heart murmurs](#) are abnormal heart sounds which can be either pathological or benign.^[47] One example of a murmur is [Still's murmur](#), which presents a musical sound in children, has no symptoms and disappears in adolescence.^[48]

A different type of sound, a [pericardial friction rub](#) can be heard in cases of pericarditis where the inflamed membranes can rub together.^[49]

Clinical significance

Disease

Transmission of a [cardiac action potential](#) through the heart's conduction system

Conduction system of the heart

The prepotential is due to a slow influx of sodium ions until the threshold is reached followed by a rapid depolarization and repolarization. The prepotential accounts for the membrane reaching threshold and initiates the spontaneous depolarization and contraction of the cell; there is no resting potential.^[7]

3D echocardiogram showing the mitral valve (right), tricuspid and mitral valves (top left) and aortic valve (top right). The closure of the heart valves causes the [heart sounds](#).

Cardiovascular diseases, which include diseases of the heart, are the leading cause of death worldwide.^[50] The majority of cardiovascular disease is noncommunicable and related to lifestyle and other factors, becoming more prevalent with ageing.^[50] Heart disease is a major cause of death, accounting for an average of 30% of all deaths in 2008, globally.^[11] This rate varies from a lower 28% to a high 40% in **high-income countries**.^[12] Doctors that specialise in the heart are called **cardiologists**. Many other medical professionals are involved in treating diseases of the heart, including **doctors** such as **general practitioners**, **cardiothoracic surgeons** and **intensivists**, and **allied health** practitioners including **physiotherapists** and **dieticians**.^[51]

Coronary artery disease is also known as ischemic heart disease, is caused by **atherosclerosis** – a build-up of **plaque** along the inner walls of the arteries which **narrows** them, reducing the blood flow to the heart.^[52] A stable plaque may cause chest pain (**angina**) or breathlessness during exercise or at rest, or no symptoms at all. A ruptured plaque can block a blood vessel and lead to ischaemia of the heart muscle, causing **unstable angina** or a **heart attack**.^[53] In the worst case this may cause cardiac arrest, a sudden and utter loss of output from the heart.^[54] **Obesity**, **high blood pressure**, uncontrolled **diabetes**, smoking and high **cholesterol** can all increase the risk of developing atherosclerosis and coronary artery disease.^{[50][52]}

Heart failure is where the heart can't beat enough blood to meet the needs of the body.^[52] It is generally a chronic condition, associated with age, that progresses gradually.^[55] Each side of the heart can fail independently of the other, resulting in heart failure of the right heart or the left heart. Left heart failure can also lead to right heart failure (**cor pulmonale**) by increasing **strain on the right heart**. If the heart is unable to pump sufficient blood, it may accumulate throughout the body, causing breathlessness in the lungs (**pulmonary congestion**; **pulmonary edema**), swelling (**edema**) of the feet or other gravity-dependent areas, decrease exercise tolerance, or cause other clinical **signs** such as an enlarged liver, cardiac murmurs, or a raised **jugular venous pressure**. Common causes of heart failure include coronary artery disease, **valve disorders** and diseases of cardiac muscle.^[56]

Cardiomyopathy is a noticeable deterioration of the heart muscle's ability to contract, which can lead to **heart failure**. The causes of many types of cardiomyopathy are poorly understood; some identified causes include **alcohol**, toxins, systemic disease such as **sarcoidosis**, and congenital conditions such as **HOCM**. The types of cardiomyopathy are described according to how they affect heart muscle. Cardiomyopathy can cause the heart to become enlarged (**hypertrophic cardiomyopathy**), constrict the outflow tracts of the heart (**restrictive cardiomyopathy**), or cause the heart to dilate and impact on the efficiency of its beating (**dilated cardiomyopathy**).^[57] HOCM is often **undiagnosed** and can cause **sudden death** in young athletes.^[7]

Heart murmurs are abnormal heart sounds which can be either related to disease or benign, and there are several kinds.^[58] There are normally two heart sounds, and abnormal heart sounds can either be extra sounds, or "murmurs" related to the flow of blood between the sounds. Murmurs are graded by volume, from 1) the quietest, to 6) the loudest, and evaluated by their relationship to the heart sounds, position in the cardiac cycle, and additional features such as their radiation to other sites, changes with a person's position, the frequency of the sound as determined by the side of the **stethoscope** by which they are heard, and site at which they are heard loudest.^[58] **Phonocardiograms** can record these sounds,^[7] and echocardiograms are generally required for their diagnosis.^[58] Murmurs can result from **valvular heart diseases** due to narrowing (**stenosis**), or regurgitation of any of the main heart valves, such as **aortic stenosis**, **mitral regurgitation** or **mitral valve prolapse**. They can also result from a number of other disorders, including **atrial** and **ventricular septal defects**.^[58] Two common and infective causes of heart murmurs, are **infective endocarditis** and **rheumatic fever**, particularly in **developing countries**. Infective endocarditis involves colonisation of a heart valve,^[59] and rheumatic fever involves an initial bacterial infection by **Group A streptococcus** followed by a reaction against heart tissue that resembles the streptococcal **antigen**.^[60]

Abnormalities in the normal **sinus rhythm** of the heart can prevent the heart from effectively pumping blood, and are generally identified by ECG. These **cardiac arrhythmias** can cause an abnormal but regular heart rhythm, such as a rapid heart rate (**tachycardia**, classified as **arising from above the ventricles** or **from the ventricles**) or a slow heart rate (**bradycardia**); or may result in irregular rhythms. Tachycardia is generally defined as a heart rate faster than 100 beats per minute, and bradycardia as a heart rate slower than 60.^[61] **Asystole** is the cessation of heart rhythm. A random and varying rhythm is classified as **atrial** or **ventricular fibrillation** depending if the electrical activity originates in the atria or the ventricles.^[61] Abnormal conduction can cause a delay or unusual order of contraction of the heart muscle. This can be a result of a disease process, such as **heart block**, or congenital, such as **Wolff-Parkinson-White syndrome**.^[61]

Diseases may also affect the pericardium which surrounds the heart, which when inflamed is called **pericarditis**. This may result from infective causes (such as **glandular fever**, **cytomegalovirus**, **coxsackievirus**, **tuberculosis** or **Q fever**), systemic disorders such as **amyloidosis** or **sarcoidosis**, tumours, high uric acid levels, and other causes. This inflammation affects the ability of the heart to pump effectively. When fluid builds up in the pericardium this is called **pericardial effusion**, which when it causes acute heart failure is called **cardiac tamponade**. This may be blood from a traumatic injury or fluid from an effusion.^[62] This can compress the heart and adversely affect the function of the heart. The fluid can be removed from the pericardial sac using a syringe in a procedure called **pericardiocentesis**.^[63]

The heart can be affected by congenital diseases. These include failure of the developmental **foramen ovale** to close, present in up to 25% of people;^[64] **ventricular** or **atrial septal defects**, congenital diseases of the heart valves (e.g. congenital **aortic stenosis**) or disease relating to blood vessels or blood flow from the heart (such as a **patent ductus arteriosus** or **aortic coarctation**).^[65] **Harrisons 1458–1465** These may cause symptoms at a variety of ages. If unoxygenated blood travels directly from the right to the left side of the heart, it may be noticed at birth, as it may cause a baby to become blue (**cyanotic**) such as **Tetralogy of Fallot**. A heart problem may impact a child's ability to grow.^[65] Some causes rectify with time and are regarded as benign. Other causes may be incidentally detected on a cardiac examination. These disorders are often diagnosed on an **echocardiogram**.^[66]

Diagnosis

Heart disease is diagnosed by the taking of a **medical history**, a **cardiac examination**, and further investigations, including **blood tests**, **echocardiograms**, **ECGs** and **imaging**. Other invasive procedures such as **cardiac catheterisation** can also play a role.^[67]

Examination

*Main article: **Cardiac examination***

The cardiac examination includes inspection, feeling the chest with the hands (**palpation**) and listening with a stethoscope (**auscultation**).^{[68][69]} It involves assessment of **signs** that may be visible on a person's hands (such as **splinter haemorrhages**), joints and other areas. A person's pulse is taken, usually at the **radial artery** near the wrist, in order to assess for the rhythm and strength of the pulse. The **blood pressure** is taken, using either a manual or automatic **sphygmomanometer** or using **a more invasive measurement** from within the artery. Any elevation of the **jugular venous pressure** is noted. A person's **chest** is felt for any transmitted vibrations from the heart, and then listened to with a stethoscope. A normal heart has two hearts

The stethoscope is used for **auscultation** of the heart, and is one of the most iconic symbols for **medicine**. A number of diseases can be detected primarily by listening for **heart murmurs**.

Atherosclerosis is a condition affecting the **circulatory system**. If the **coronary arteries** are affected **angina pectoris** may result or at worse a **heart attack**.


sounds - additional heart sounds or [heart murmurs](#) may also be able to be heard and may point to disease. Additional tests may be conducted to assess a person's heart murmurs if they are present, and peripheral signs of heart disease such as [swollen feet](#) or [fluid in the lungs](#) may be assessed.^[69]

Electrocardiogram

Main article: [Electrocardiography](#)

Using surface electrodes on the body, it is possible to record the electrical activity of the heart. This tracing of the electrical signal is the electrocardiogram (ECG) or (EKG). An ECG is a [bedside test](#) and usually requires the placement of ten leads on the body. This produces a "12 lead" ECG (three extra leads are calculated mathematically, and one lead is a [ground](#)).^[69]

There are five prominent features on the ECG: the P wave (atrial depolarisation), the QRS complex (atrial repolarisation and ventricular depolarisation) and the T wave (ventricular repolarisation).^[7] These reflect the summed [action potential](#) of the heart's muscle cells as they contract. A downward deflection on the ECG implies cells are becoming more negative in charge ("depolarising"), whereas an upward inflection implies cells are becoming more positive ("repolarising"). The ECG is a useful tool in detecting rhythm disturbances and in detecting insufficient blood supply to the heart.^[68] Sometimes abnormalities are not immediately visible on the ECG. [Testing when exercising](#) can be used to provoke an abnormality, or an ECG can be worn for a longer period such as a 24-hour [Holter monitor](#) if a suspected rhythm abnormality is not present at the time of assessment.^[69]



Imaging

Main article: [Cardiac imaging](#)

Several [imaging methods](#) can be used to assess the anatomy and function of the heart, including [ultrasound \(echocardiography\)](#), [angiography](#), [CT scans](#), [MRI](#) and [PET](#). An echocardiogram is an ultrasound of the heart used to measure the heart's function, assess for valve disease, and look for any abnormalities. Echocardiography can be conducted by a probe on the chest ("transthoracic") or by a probe in the [esophagus](#) ("transoesophageal"). A typical echocardiography report will include information about the width of the valves noting any [stenosis](#), whether there is any backflow of blood ([regurgitation](#)) and information about the blood volumes at the end of systole and diastole, including an [ejection fraction](#), which describes how much blood is ejected from the left and right ventricles after systole. Ejection fraction can then be obtained by dividing the volume ejected by the heart (stroke volume) by the volume of the filled heart (end-diastolic volume).^[70] Echocardiograms can also be conducted under circumstances when the body is more stressed, in order to examine for signs of lack of blood supply. This [cardiac stress test](#) involves either direct exercise, or where this is not possible, injection of a drug such as [dobutamine](#).^[69]

CT scans, [chest X-rays](#) and other forms of [imaging](#) can help evaluate the heart's size, evaluate for signs of [pulmonary oedema](#), and indicate whether there is [fluid around the heart](#). They are also useful for evaluating the aorta, the major blood vessel which leaves the heart. ^[69]

Treatment

A number of medications are used to treat diseases of the heart, or ameliorate symptoms.

For diseases of the heart rate or rhythm, a number of different [antiarrhythmic agents](#) are used. These may interfere with electrolyte channels and thus the cardiac action potential (such as [calcium channel blockers](#), [sodium channel blockers](#)), interfere with stimulation of the heart by the sympathetic nervous system ([beta blockers](#)), or interfere with the movement of [sodium and potassium across the cell membrane](#), such as [digoxin](#).^[71] Other examples include [atropine](#) for slow rhythms, and [amiodarone](#) for irregular rhythms. Such medications are not the only way of treating diseases of heart rate or rhythm. In the context of a new-onset irregular heart rhythm ([atrial fibrillation](#)), immediate electrical [cardioversion](#) may be attempted. For a slow heartbeat or [heart block](#), a [pacemaker](#) or [defibrillator](#) may be inserted.^[72] The acuity of onset often affects how a rhythm disturbance is managed, as does whether a rhythm causes hemodynamic instability, such as low blood pressure or symptoms. An instigating cause is investigated for, such as a heart attack, medication, or metabolic problem.^[72]

For ischaemic heart disease, treatment also includes amelioration of symptoms. This includes [GTN](#), [beta blockers](#) and, in the context of an acute event, stronger pain relief such as [morphine](#) and other [opiates](#). Many of these drugs have additional protective benefits, by decreasing the sympathetic tone on the heart that occurs with the pain, or by dilating blood vessels (GTN).^[72]

Treatment of heart disease includes [primary](#) and [secondary prevention](#) to prevent the occurrence or worsening of symptoms and [atherosclerosis](#). This includes recommendations to cease smoking, decrease alcohol consumption, increase exercise, and make modifications to their diet to decrease the consumption of fats and sugars. Medications may also be given to help better control concurrent [diabetes](#). [Statins](#) or other drugs such as [fibrates](#) may also be given to decrease a person's [cholesterol](#) levels. [Blood pressure](#) medication may also be commenced or modified.^[72]

For many diseases of the heart, including atrial fibrillation and valvular disease, and after a heart operation, anticoagulation in the form of [aspirin](#), [warfarin](#), [clopidogrel](#) or [novel oral anticoagulants](#) is often given simultaneously, because of an increased risk of stroke or, in the context of a clotted heart vessel, rethrombosis.^[72]

Surgery

Main articles: [Cardiac surgery](#), [Coronary artery bypass surgery](#), and [Coronary stent](#)

Surgery, when considered necessary for diseases of the heart, can take place via an [open operation](#) or via small guidewires inserted via peripheral arteries ("[percutaneous coronary intervention](#)"). Percutaneous coronary intervention is usually used in the context of an [acute coronary syndrome](#), and may be used to insert a [stent](#).^[73]

[Coronary artery bypass surgery](#) is one such operation. In this operation, one or more arteries surrounding the heart that have become narrowed are bypassed. This is done by taking blood vessels harvested from another part of the body. Commonly harvested veins include the [saphenous veins](#) or the [internal mammary artery](#). Because this operation involves the heart tissue, [a machine is used](#) so that blood can bypass the heart during the operation.^[74]

[Heart valve repair](#) or [valve replacement](#) are options for diseases of the heart valves. ^[73]

History

Ancient

Humans have known about the heart since ancient times, although its precise function and anatomy were not clearly understood. ^[75] From the primarily religious views of earlier societies towards the heart, [ancient Greeks](#) are considered to have been the primary seat of scientific understanding of the heart in the ancient world.^{[76][77][78]} [Aristotle](#) considered the heart to be organ responsible for creating blood; [Plato](#) considered the heart as the source of circulating blood and [Hippocrates](#) noted blood circulating cyclically from the body through the heart to the lungs. ^{[76][78]} [Erasistratos](#) (304-250 BC) noted the heart as a pump, causing dilation of blood vessels, and noted that arteries and veins both radiate from the heart,

becoming progressively smaller with distance, although he believed they were filled with air and not blood. He also discovered the heart valves.^[76]

The Greek physician [Galen](#) (2nd century AD) knew blood vessels carried blood and identified venous (dark red) and arterial (brighter and thinner) blood, each with distinct and separate functions.^[76] Galen, noting the heart as the hottest organ in the body, concluded that it provided heat to the body.^[78] The heart did not pump blood around, the heart's motion sucked blood in during diastole and the blood moved by the pulsation of the arteries themselves.^[78] Galen believed the arterial blood was created by venous blood passing from the left ventricle to the right through 'pores' between the ventricles.^[75] Air from the lungs passed from the lungs via the pulmonary artery to the left side of the heart and created arterial blood.^[78]

These ideas went unchallenged for almost a thousand years.^{[75][78]}

Pre-modern

The earliest descriptions of the [coronary](#) and [pulmonary circulation](#) systems can be found in the *Commentary on Anatomy in Avicenna's Canon*, published in 1242 by [Ibn al-Nafis](#).^[79] In his manuscript, al-Nafis wrote that blood passes through the pulmonary circulation instead of moving from the right to the left ventricle as previously believed by Galen.^[80] His work was later translated into [Latin](#) by [Andrea Alpago](#).^[81]

In Europe, the teachings of Galen continued to dominate the academic community and his doctrines were adopted as the official canon of the Church. [Andreas Vesalius](#) questioned some of Galen's beliefs of the heart in *De humani corporis fabrica* (1543), but his *magnum opus* was interpreted as a challenge to the authorities and he was subjected to a number of attacks.^[82] [Michael Servetus](#) wrote in *Christianismi Restitutio* (1553) that blood flows from one side of the heart to the other via the lungs.^[82]

Modern

The breakthrough came with the publication of *De Motu Cordis* (1628) by the English physician [William Harvey](#). Harvey's book completely describes the [systemic circulation](#) and the mechanical force of the heart, leading to an overhaul of the Galenic doctrines. ^[83] [Otto Frank](#) (1865–1944) was a German physiologist; among his many published works are detailed studies of this important heart relationship. [Ernest Starling](#) (1866–1927) was an important English physiologist who also studied the heart. Although they worked largely independently, their combined efforts and similar conclusions have been recognized in the name "[Frank–Starling mechanism](#)".^[7]

Although [Purkinje fibers](#) and the [bundle of His](#) were discovered as early as the 19th century, their specific role in the [electrical conduction system of the heart](#) remained unknown until [Sunao Tawara](#) published his monograph, titled *Das Reizleitungssystem des Säugetierherzens*, in 1906. Tawara's discovery of the [atrioventricular node](#) prompted [Arthur Keith](#) and [Martin Flack](#) to look for similar structures in the heart, leading to their discovery of the [sinoatrial node](#) several months later. These structures form the anatomical basis of the [electrocardiogram](#), whose inventor, [Willem Einthoven](#), was awarded the Nobel Prize in Medicine or Physiology in 1924.^[84]

The first successful [heart transplantation](#) was performed in 1967 by the South African surgeon [Christiaan Barnard](#) at [Groote Schuur Hospital](#) in [Cape Town](#). This marked an important milestone in [cardiac surgery](#), capturing the attention of both the medical profession and the world at large. However, long-term survival rates of patients were initially very low. [Louis Washkansky](#), the first recipient of a donated heart, died 18 days after the operation while other patients did not survive for more than a few weeks.^[85] The American surgeon [Norman Shumway](#) has been credited for his efforts to improve transplantation techniques, along with pioneers [Richard Lower](#), [Vladimir Demikhov](#) and [Adrian Kantrowitz](#). As of March 2000, more than 55,000 heart transplantations have been performed worldwide.^[86]

By the middle of the 20th century, [heart disease](#) had surpassed infectious disease as the leading cause of death in the United States, and it is currently the leading cause of deaths worldwide. Since 1948, the ongoing [Framingham Heart Study](#) has shed light on the effects of various influences on the heart, including diet, exercise, and common medications such as [aspirin](#). Although the introduction of [ACE inhibitors](#) and [beta blockers](#) has improved the management of chronic [heart failure](#), the disease continues to be an enormous medical and societal burden, with 30 to 40% of patients dying within a year of receiving the diagnosis.^[87]

Society and culture

Further information: *Sacred Heart*, *Heart symbol*, and *Blood § Cultural and religious beliefs*

Symbolism


As one of the vital organs, the heart was long identified as the center of the entire body, the seat of life, or emotion, or reason, will, intellect, purpose or the mind.^[88] The heart is an emblematic symbol in many religions, signifying "truth, conscience or moral courage in many religions - the temple or throne of God in Islamic and [Judeo-Christian](#) thought; the divine centre, or [atman](#), and the [third eye](#) of transcendent wisdom in [Hinduism](#); the diamond of purity and essence of the [Buddha](#); the [Taoist](#) centre of understanding."^[88]

In the [Hebrew Bible](#), the word for heart, *lev*, is used in these meanings, as the seat of emotion, the mind, and referring to the anatomical organ. It is also connected in function and symbolism to the stomach.^[89]


An important part of the concept of the [soul](#) in [Ancient Egyptian religion](#) was thought to be the heart, or *ib*. The *ib* or metaphysical heart was believed to be formed from one drop of blood from the child's mother's heart, taken at conception.^[90] To ancient Egyptians, the heart was the seat of [emotion](#), [thought](#), will, and [intention](#). This is evidenced by [Egyptian](#) expressions which incorporate the word *ib*, such as *Awi-ib* for "happy" (literally, "long of heart"), *Xak-ib* for "estranged" (literally, "truncated of heart").^[91] In Egyptian religion, the heart was the key to the afterlife. It was conceived as surviving death in the nether world, where it gave evidence for, or against, its possessor. It was thought that the heart was examined by [Anubis](#) and a variety of [deities](#) during the *Weighing of the Heart* ceremony. If the heart weighed more than the feather of [Maat](#), which symbolized the ideal standard of behavior. If the scales balanced, it meant the heart's possessor had lived a just life and could enter the afterlife; if the heart was heavier, it would be devoured by the monster [Ammit](#).^[92]

The [Chinese](#) character for "heart", 心, derives from a comparatively realistic depiction of a heart (indicating the heart chambers) in [seal script](#).^[93] The Chinese word *xīn* also takes the metaphorical meanings of "mind", "intention", or "core".^[94] In [Chinese medicine](#), the heart is seen as the center of [神](#) *shén* "spirit, consciousness".^[95] The heart is associated with the [small intestine](#), [tongue](#), governs the [six organs and five viscera](#), and belongs to fire in the five elements.^[96]

The Sanskrit word for heart is *hr̥d* or *hr̥daya*, found in the oldest surviving Sanskrit text, the [Rigveda](#). In Sanskrit, it may mean both the anatomical



Heart and its blood vessels, by [Leonardo da Vinci](#), 15th century



Letter [Ⴀ](#) of the [Georgian script](#) is often used as a "heart" symbol.



The [seal script](#) glyph for "heart" ([Middle Chinese](#) *sim*)

object and "mind" or "soul", representing the seat of emotion. *Hrd* may be a cognate of the word for heart in Greek, Latin, and English.^{[97][98]} Many [classical](#) philosophers and scientists, including [Aristotle](#), considered the heart the seat of thought, [reason](#), or emotion, often disregarding the brain as contributing to those functions.^[99] The identification of the heart as the seat of [emotions](#) in particular is due to the [Roman](#) physician [Galen](#), who also located the seat of the passions in the [liver](#), and the seat of reason in the brain.^[100]

The heart also played a role in the [Aztec](#) system of belief. The most common form of human sacrifice practiced by the Aztecs was heart-extraction. The Aztec believed that the heart (*tona*) was both the seat of the individual and a fragment of the Sun's heat (*istli*). To this day, the Nahua consider the Sun to be a heart-soul (*tona-tiuh*): "round, hot, pulsating".^[101]

In [Catholicism](#), there has been a long tradition of worship of the heart, stemming from worship of the wounds of [Jesus Christ](#) which gained prominence from the mid sixteenth century.^[102] This tradition influenced the development of the medieval Christian [devotion](#) to the [Sacred Heart of Jesus](#) and the parallel worship of [Immaculate Heart of Mary](#), made popular by [John Eudes](#).^[103]

The expression of a [broken heart](#) is a cross-cultural reference to [grief](#) for a lost one or to unfulfilled [romantic love](#).

The notion of "[Cupid's arrows](#)" is ancient, due to [Ovid](#), but while Ovid describes Cupid as wounding his victims with his arrows, it is not made explicit that it is the *heart* that is wounded. The familiar iconography of Cupid shooting little [heart symbols](#) is a [Renaissance](#) theme that became tied to [Valentine's day](#).^[88]

Food

Animal hearts are widely consumed as food. As they are almost entirely muscle, they are high in protein. They are often included in dishes with other [offal](#), for example in the [pan-Ottoman kokoretsi](#).

Chicken hearts are considered to be [giblets](#), and are often grilled on skewers: [Japanese hāto yakitori](#), [Brazilian churrasco de coração](#), [Indonesian chicken heart satay](#).^[104] They can also be pan-fried, as in [Jerusalem mixed grill](#). In [Egyptian cuisine](#), they can be used, finely chopped, as part of [stuffing](#) for chicken.^[105] Many recipes combined them with other giblets, such as the [Mexican pollo en menudencias](#)^[106] and the [Russian ragu iz kurinyikh potrokhov](#).^[107]

The hearts of beef, pork, and mutton can generally be interchanged in recipes. As heart is a hard-working muscle, it makes for "firm and rather dry" meat,^[108] so is generally slow-cooked. Another way of dealing with toughness is to [julienne](#) the meat, as in [Chinese](#) stir-fried heart.^[109]

[Beef](#) heart may be grilled or braised.^[108] In the [Peruvian anticuchos de corazón](#), barbecued beef hearts are grilled after being tenderized through long [marination](#) in a spice and vinegar mixture. An [Australian](#) recipe for "mock goose" is actually braised stuffed beef heart.^[110]

[Pig](#) heart is stewed, poached, braised,^[111] or made into sausage. The [Balinese oret](#) is a sort of [blood sausage](#) made with pig heart and blood. A [French](#) recipe for *cœur de porc à l'orange* is made of braised heart with an orange sauce.

Other animals

The structure of the heart varies among the different [animal groups](#). [Cephalopods](#) have two "gill hearts" also known as [branchial hearts](#) and one "systemic heart". The [vertebrate](#) heart lies in the front (ventral) part of the body cavity, dorsal to the gut. It is always surrounded by a [pericardium](#), which is usually a distinct structure, but may be continuous with the [peritoneum](#) in jawless and cartilaginous fish.

The SA node is found in all [amniotes](#) but not in more primitive vertebrates. In these animals, the muscles of the heart are relatively continuous and the sinus venosus coordinates the beat which passes in a wave through the remaining chambers. Indeed, since the sinus venosus is incorporated into the right atrium in amniotes, it is likely [homologous](#) with the SA node. In teleosts, with their vestigial sinus venosus, the main centre of coordination is, instead, in the atrium. The rate of heartbeat varies enormously between different species, ranging from around 20 beats per minute in [codfish](#) to around 600 in [hummingbirds](#)^[112] and up to 1200 bpm in the [ruby-throated hummingbird](#).^[113]

Double circulatory systems

In the heart of [lungfish](#), the septum extends part-way into the ventricle. This allows for some degree of separation between the de-oxygenated bloodstream destined for the lungs and the oxygenated stream that is delivered to the rest of the body. The absence of such a division in living amphibian species may be partly due to the amount of respiration that occurs through the skin; thus, the blood returned to the heart through the venae cavae is already partially oxygenated. As a result, there may be less need for a finer division between the two bloodstreams than in lungfish or other [tetrapods](#). Nonetheless, in at least some species of amphibian, the spongy nature of the ventricle does seem to maintain more of a separation between the bloodstreams. Also, the original valves of the [conus arteriosus](#) have been replaced by a spiral valve that divides it into two parallel parts, thereby helping to keep the two bloodstreams separate.^[112]

Adult [amphibians](#) and most [reptiles](#) have a [double circulatory system](#) but the heart is not separated into two pumps. The development of the double system is necessitated by the presence of [lungs](#) which deliver oxygenated blood directly to the heart.

In amphibians, the atrium is divided into two chambers by a muscular [septum](#) but there is only one ventricle. The sinus venosus, which remains large, connects only to the right atrium and receives blood from the [venae cavae](#), with the [pulmonary vein](#) by-passing it to enter the left atrium.

The heart of most reptiles is similar in structure to that of lungfish but the septum is generally much larger. This divides the ventricle into two halves but the septum does not reach the whole length of the heart and there is a considerable gap near the pulmonary artery and aorta openings. In most reptilian species, there appears to be little, if any, mixing between the bloodstreams, so the aorta receives, essentially, only oxygenated blood.^[112]


The fully divided heart

[Archosaurs](#) ([crocodilians](#) and [birds](#)) and [mammals](#) show complete separation of the heart into two pumps for a total of four [heart chambers](#); it is thought that the four-chambered heart of archosaurs evolved independently from that of mammals. In crocodilians, there is a small opening, the [foramen of Panizza](#), at the base of the arterial trunks and there is some degree of mixing between the blood in each side of the heart, during a dive underwater;^{[114][115]} thus, only in birds and mammals are the two streams of blood – those to the pulmonary and systemic circulations – permanently kept entirely separate by a physical barrier.^[112]

Fish

Main article: *Fish anatomy § Heart*


Primitive fish have a four-chambered heart, but the chambers are arranged sequentially so that this primitive heart is quite unlike the four-chambered



hearts of mammals and birds. The first chamber is the [sinus venosus](#), which collects deoxygenated blood, from the body, through the [hepatic](#) and [cardinal veins](#). From here, blood flows into the [atrium](#) and then to the powerful muscular [ventricle](#) where the main pumping action will take place. The fourth and final chamber is the [conus arteriosus](#) which contains several valves and sends blood to the [ventral aorta](#). The ventral aorta delivers blood to the gills where it is oxygenated and flows, through the [dorsal aorta](#), into the rest of the body. (In [tetrapods](#), the ventral aorta has divided in two; one half forms the [ascending aorta](#), while the other forms the [pulmonary artery](#)).^[112]

In the adult fish, the four chambers are not arranged in a straight row but, instead form an S-shape with the latter two chambers lying above the former two. This relatively simpler pattern is found in [cartilaginous fish](#) and in the [ray-finned fish](#). In [teleosts](#), the conus arteriosus is very small and can more accurately be described as part of the aorta rather than of the heart proper. The conus arteriosus is not present in any [amniotes](#), presumably having been absorbed into the ventricles over the course of evolution. Similarly, while the sinus venosus is present as a vestigial structure in some reptiles and birds, it is otherwise absorbed into the right atrium and is no longer distinguishable.^[112]


Invertebrates




Basic arthropod body structure – heart shown in red

Arthropods have an open circulatory system, and often some short open-ended arteries. The arthropod heart is typically a muscular tube that runs the length of the body, under the back and from the base of the head. Instead of blood the circulatory fluid is [haemolymph](#) which carries the most commonly used [respiratory pigment](#), copper-based [haemocyanin](#) as the oxygen transporter; iron-based haemoglobin is used by only a few arthropods. The heart contracts in ripples from the rear to the front of the animal transporting water and [nutrients](#). Pairs of valves run alongside the heart, allowing fluid to enter whilst preventing backflow.

In [insects](#), the circulatory system is not used to transport oxygen and so is much reduced, having no veins or arteries and consisting of a single perforated tube running dorsally which pumps [peristaltically](#). The simpler unsegmented invertebrates have no [body cavity](#), and oxygen and nutrients pass through their bodies by diffusion.




Blood flow through the fish heart: sinus venosus, atrium, ventricle, and outflow tract



The tube-like heart (green) of the mosquito *Anopheles gambiae* extends horizontally across the body, interlinked with the diamond-shaped wing muscles (also green) and surrounded by pericardial cells (red). Blue depicts cell nuclei.

Additional images



The human heart viewed from the front The human heart viewed from behind The coronary circulation An anatomical specimen of the heart Heart Illustration with circulatory system

Notes

- a. [^] From the heart to the body
- b. [^] Arteries that contain deoxygenated blood, from the heart to the lungs
- c. [^] Supplying blood to the heart itself
- d. [^] From the body to the heart
- e. [^] Veins containing oxygenated blood from the lungs to the heart
- f. [^] Veins that drain blood from the cardiac tissue itself
- g. [^] Note the muscles do **not** cause the valves to open. The pressure difference between the blood in the atria and the ventricles does this.

References

This article incorporates text from the **CC-BY** book: *OpenStax College, Anatomy & Physiology. OpenStax CNX. 30 jul 2014* ..

- [^] ^a ^b ^c ^d ^e ^f ^g ^h ⁱ ^j ^k ^l ^m ⁿ ^o ^p ^q ^r ^s ^t ^u ^v ^w ^x ^y ^z ^{aa} ^{ab} ^{ac} ^{ad} ^{ae} ^{af} ^{ag} ^{ah} ^{ai} ^{aj} ^{ak} ^{al} ^{am} ^{an} ^{ao} ^{ap} ^{aq} ^{ar} ^{as} ^{at} ^{au} ^{av} ^{aw} ^{ax} ^{ay} ^{az} ^{ba} ^{bb} ^{bc} ^{bd} ^{be} ^{bf} ^{bg} ^{bh} ^{bi} ^{bj} ^{bk} ^{bl} ^{bm} ^{bn} ^{bo} ^{bp} ^{bq} ^{br} ^{bs} ^{bt} ^{bu} ^{bv} ^{bw} ^{bx} ^{by} ^{bz} ^{ca} ^{cb} ^{cc} ^{cd} ^{ce} ^{cf} ^{cg} ^{ch} ^{ci} ^{cj} ^{ck} ^{cl} ^{cm} ^{cn} ^{co} ^{cp} ^{cq} ^{cr} ^{cs} ^{ct} ^{cu} ^{cv} ^{cw} ^{cx} ^{cy} ^{cz} ^{da} ^{db} ^{dc} ^{dd} ^{de} ^{df} ^{dg} ^{dh} ^{di} ^{dj} ^{dk} ^{dl} ^{dm} ^{dn} ^{do} ^{dp} ^{dq} ^{dr} ^{ds} ^{dt} ^{du} ^{dv} ^{dw} ^{dx} ^{dy} ^{dz} ^{ea} ^{eb} ^{ec} ^{ed} ^{ee} ^{ef} ^{eg} ^{eh} ^{ei} ^{ej} ^{ek} ^{el} ^{em} ^{en} ^{eo} ^{ep} ^{eq} ^{er} ^{es} ^{et} ^{eu} ^{ev} ^{ew} ^{ex} ^{ey} ^{ez} ^{fa} ^{fb} ^{fc} ^{fd} ^{fe} ^{ff} ^{fg} ^{fh} ^{fi} ^{fj} ^{fk} ^{fl} ^{fm} ^{fn} ^{fo} ^{fp} ^{fq} ^{fr} ^{fs} ^{ft} ^{fu} ^{fv} ^{fw} ^{fx} ^{fy} ^{fz} ^{ga} ^{gb} ^{gc} ^{gd} ^{ge} ^{gf} ^{gg} ^{gh} ^{gi} ^{gj} ^{gk} ^{gl} ^{gm} ^{gn} ^{go} ^{gp} ^{gq} ^{gr} ^{gs} ^{gt} ^{gu} ^{gv} ^{gw} ^{gx} ^{gy} ^{gz} ^{ha} ^{hb} ^{hc} ^{hd} ^{he} ^{hf} ^{hg} ^{hh} ^{hi} ^{hj} ^{hk} ^{hl} ^{hm} ^{hn} ^{ho} ^{hp} ^{hq} ^{hr} ^{hs} ^{ht} ^{hu} ^{hv} ^{hw} ^{hx} ^{hy} ^{hz} ^{ia} ^{ib} ^{ic} ^{id} ^{ie} ^{if} ^{ig} ^{ih} ⁱⁱ ^{ij} ^{ik} ^{il} ^{im} ⁱⁿ ^{io} ^{ip} ^{iq} ^{ir} ^{is} ^{it} ^{iu} ^{iv} ^{iw} ^{ix} ^{iy} ^{iz} ^{ja} ^{jb} ^{jc} ^{jd} ^{je} ^{jf} ^{jj} ^{jk} ^{jl} ^{jm} ^{jn} ^{jo} ^{jp} ^{jq} ^{jr} ^{js} ^{jt} ^{ju} ^{jv} ^{jw} ^{jx} ^{ka} ^{kb} ^{kc} ^{kd} ^{ke} ^{kf} ^{kg} ^{kh} ^{ki} ^{kj} ^{kl} ^{km} ^{kn} ^{ko} ^{kp} ^{kq} ^{kr} ^{ks} ^{kt} ^{ku} ^{kv} ^{kx} ^{ky} ^{kz} ^{la} ^{lb} ^{lc} ^{ld} ^{le} ^{lf} ^{lg} ^{lh} ^{li} ^{lj} ^{lk} ^{ll} ^{lm} ^{ln} ^{lo} ^{lp} ^{lq} ^{lr} ^{ls} ^{lt} ^{lu} ^{lv} ^{lw} ^{lx} ^{ly} ^{lz} ^{ma} ^{mb} ^{mc} ^{md} ^{me} ^{mf} ^{mg} ^{mh} ^{mi} ^{mj} ^{mk} ^{ml} ^{mm} ^{mn} ^{mo} ^{mp} ^{mq} ^{mr} ^{ms} ^{mt} ^{mu} ^{mv} ^{mw} ^{mx} ^{my} ^{mz} ^{na} ^{nb} ^{nc} nd ^{ne} ^{nf} ^{ng} ^{nh} ⁿⁱ ^{nj} ^{nk} ^{nl} ^{nm} ⁿⁿ ^{no} ^{np} ^{nq} ^{nr} ^{ns} ^{nt} ^{nu} ^{nv} ^{nw} ^{nx} ^{ny} ^{nz} ^{oa} ^{ob} ^{oc} ^{od} ^{oe} ^{of} ^{og} ^{oh} ^{oi} ^{oj} ^{ok} ^{ol} ^{om} ^{on} ^{oo} ^{op} ^{oq} ^{or} ^{os} ^{ot} ^{ou} ^{ov} ^{ow} ^{ox} ^{oy} ^{oz} ^{pa} ^{pb} ^{pc} ^{pd} ^{pe} ^{pf} ^{pg} ^{ph} ^{pi} ^{pj} ^{pk} ^{pl} ^{pm} ^{pn} ^{po} ^{pp} ^{pq} ^{pr} ^{ps} ^{pt} ^{pu} ^{pv} ^{pw} ^{px} ^{py} ^{pz} ^{qa} ^{qb} ^{qc} ^{qd} ^{qe} ^{qf} ^{qg} ^{qh} ^{qi} ^{qj} ^{qk} ^{ql} ^{qm} ^{qn} ^{qo} ^{qp} ^{qq} ^{qr} ^{qs} ^{qt} ^{qu} ^{qv} ^{qw} ^{qx} ^{qy} ^{qz} ^{ra} ^{rb} ^{rc} rd ^{re} ^{rf} ^{rg} ^{rh} ^{ri} ^{rj} ^{rk} ^{rl} ^{rm} ^{rn} ^{ro} ^{rp} ^{rq} ^{rr} ^{rs} ^{rt} ^{ru} ^{rv} ^{rw} ^{rx} ^{ry} ^{rz} ^{sa} ^{sb} ^{sc} ^{sd} ^{se} ^{sf} ^{sg} ^{sh} ^{si} ^{sj} ^{sk} ^{sl} sm ^{sn} ^{so} ^{sp} ^{sq} ^{sr} ^{ss} st ^{su} ^{sv} ^{sw} ^{sx} ^{sy} ^{sz} ^{ta} ^{tb} ^{tc} ^{td} ^{te} ^{tf} ^{tg} th ^{ti} ^{tj} ^{tk} ^{tl} tm ^{tn} ^{to} ^{tp} ^{tq} ^{tr} ^{ts} ^{tt} ^{tu} ^{tv} ^{tw} ^{tx} ^{ty} ^{tz} ^{ua} ^{ub} ^{uc} ^{ud} ^{ue} ^{uf} ^{ug} ^{uh} ^{ui} ^{uj} ^{uk} ^{ul} ^{um} ^{un} ^{uo} ^{up} ^{uq} ^{ur} ^{us} ^{ut} ^{uu} ^{uv} ^{uw} ^{ux} ^{uy} ^{uz} ^{va} ^{vb} ^{vc} ^{vd} ^{ve} ^{vf} ^{vg} ^{vh} ^{vi} ^{vj} ^{vk} ^{vl} ^{vm} ^{vn} ^{vo} ^{vp} ^{vq} ^{vr} ^{vs} ^{vt} ^{vu} ^{vv} ^{vw} ^{vx} ^{vy} ^{vz} ^{wa} ^{wb} ^{wc} ^{wd} ^{we} ^{wf} ^{wg} ^{wh} ^{wi} ^{wj} ^{wk} ^{wl} ^{wm} ^{wn} ^{wo} ^{wp} ^{wq} ^{wr} ^{ws} ^{wt} ^{wu} ^{wv} ^{ww} ^{wx} ^{wy} ^{wz} ^{xa} ^{xb} ^{xc} ^{xd} ^{xe} ^{xf} ^{xg} ^{xh} ^{xi} ^{xj} ^{xk} ^{xl} ^{xm} ^{xn} ^{xo} ^{xp} ^{xq} ^{xr} ^{xs} ^{xt} ^{xu} ^{xv} ^{xw} ^{xx} ^{xy} ^{xz} ^{ya} ^{yb} ^{yc} ^{yd} ^{ye} ^{yf} ^{yg} ^{yh} ^{yi} ^{yj} ^{yk} ^{yl} ^{ym} ^{yn} ^{yo} ^{yp} ^{yq} ^{yr} ^{ys} ^{yt} ^{yu} ^{yv} ^{yw} ^{yx} ^{yy} ^{yz} ^{za} ^{zb} ^{zc} ^{zd} ^{ze} ^{zf} ^{zg} ^{zh} ^{zi} ^{zj} ^{zk} ^{zl} ^{zm} ^{zn} ^{zo} ^{zp} ^{zq} ^{zr} ^{zs} ^{zt} ^{zu} ^{zv} ^{zw} ^{zx} ^{zy} ^{zz} ^{aa} ^{ab} ^{ac} ^{ad} ^{ae} ^{af} ^{ag} ^{ah} ^{ai} ^{aj} ^{ak} ^{al} ^{am} ^{an} ^{ao} ^{ap} ^{aq} ^{ar} ^{as} ^{at} ^{au} ^{av} ^{aw} ^{ax} ^{ay} ^{az} ^{ba} ^{bb} ^{bc} ^{bd} ^{be} ^{bf} ^{bg} ^{bh} ^{bi} ^{bj} ^{bk} ^{bl} ^{bm} ^{bn} ^{bo} ^{bp} ^{bq} ^{br} ^{bs} ^{bt} ^{bu} ^{bv} ^{bw} ^{bx} ^{by} ^{bz} ^{ca} ^{cb} ^{cc} ^{cd} ^{ce} ^{cf} ^{cg} ^{ch} ^{ci} ^{cj} ^{ck} ^{cl} ^{cm} ^{cn} ^{co} ^{cp} ^{cq} ^{cr} ^{cs} ^{ct} ^{cu} ^{cv} ^{cw} ^{cx} ^{cy} ^{cz} ^{da} ^{db} ^{dc} ^{dd} ^{de} ^{df} ^{dg} ^{dh} ^{di} ^{dj} ^{dk} ^{dl} ^{dm} ^{dn} ^{do} ^{dp} ^{dq} ^{dr} ^{ds} ^{dt} ^{du} ^{dv} ^{dw} ^{dx} ^{dy} ^{dz} ^{ea} ^{eb} ^{ec} ^{ed} ^{ee} ^{ef} ^{eg} ^{eh} ^{ei} ^{ej} ^{ek} ^{el} ^{em} ^{en} ^{eo} ^{ep} ^{eq} ^{er} ^{es} ^{et} ^{eu} ^{ev} ^{ew} ^{ex} ^{ey} ^{ez} ^{fa} ^{fb} ^{fc} ^{fd} ^{fe} ^{ff} ^{fg} ^{fh} ^{fi} ^{fj} ^{fk} ^{fl} ^{fm} ^{fn} ^{fo} ^{fp} ^{fq} ^{fr} ^{fs} ^{ft} ^{fu} ^{fv} ^{fw} ^{fx} ^{fy} ^{fz} ^{ga} ^{gb} ^{gc} ^{gd} ^{ge} ^{gf} ^{gg} ^{gh} ^{gi} ^{gj} ^{gk} ^{gl} ^{gm} ^{gn} ^{go} ^{gp} ^{gq} ^{gr} ^{gs} ^{gt} ^{gu} ^{gv} ^{gw} ^{gx} ^{gy} ^{gz} ^{ha} ^{hb} ^{hc} ^{hd} ^{he} ^{hf} ^{hg} ^{hh} ^{hi} ^{hj} ^{hk} ^{hl} ^{hm} ^{hn} ^{ho} ^{hp} ^{hq} ^{hr} ^{hs} ^{ht} ^{hu} ^{hv} ^{hw} ^{hx} ^{hy} ^{hz} ^{ia} ^{ib} ^{ic} ^{id} ^{ie} ^{if} ^{ig} ^{ih} ⁱⁱ ^{ij} ^{ik} ^{il} ^{im} ⁱⁿ ^{io} ^{ip} ^{iq} ^{ir} ^{is} ^{it} ^{iu} ^{iv} ^{iw} ^{ix} ^{iy} ^{iz} ^{ja} ^{jb} ^{jc} ^{jd} ^{je} ^{jf} ^{jj} ^{jk} ^{jl} ^{jm} ^{jn} ^{jo} ^{jp} ^{jq} ^{jr} ^{js} ^{jt} ^{ju} ^{jv} ^{jw} ^{jx} ^{ka} ^{kb} ^{kc} ^{kd} ^{ke} ^{kf} ^{kg} ^{kh} ^{ki} ^{kj} ^{kl} ^{km} ^{kn} ^{ko} ^{kp} ^{kq} ^{kr} ^{ks} ^{kt} ^{ku} ^{kv} ^{kx} ^{ky} ^{kz} ^{la} ^{lb} ^{lc} ^{ld} ^{le} ^{lf} ^{lg} ^{lh} ^{li} ^{lj} ^{lk} ^{ll} ^{lm} ^{ln} ^{lo} ^{lp} ^{lq} ^{lr} ^{ls} ^{lt} ^{lu} ^{lv} ^{lw} ^{lx} ^{ly} ^{lz} ^{ma} ^{mb} ^{mc} ^{md} ^{me} ^{mf} ^{mg} ^{mh} ^{mi} ^{mj} ^{mk} ^{ml} ^{mm} ^{mn} ^{mo} ^{mp} ^{mq} ^{mr} ^{ms} ^{mt} ^{mu} ^{mv} ^{mw} ^{mx} ^{my} ^{mz} ^{na} ^{nb} ^{nc} nd ^{ne} ^{nf} ^{ng} ^{nh} ⁿⁱ ^{nj} ^{nk} ^{nl} ^{nm} ⁿⁿ ^{no} ^{np} ^{nq} ^{nr} ^{ns} ^{nt} ^{nu} ^{nv} ^{nw} ^{nx} ^{ny} ^{nz} ^{oa} ^{ob} ^{oc} ^{od} ^{oe} ^{of} ^{og} ^{oh} ^{oi} ^{oj} ^{ok} ^{ol} ^{om} ^{on} ^{oo} ^{op} ^{oq} ^{or} ^{os} ^{ot} ^{ou} ^{ov} ^{ow} ^{ox} ^{oy} ^{oz} ^{pa} ^{pb} ^{pc} ^{pd} ^{pe} ^{pf} ^{pg} ^{ph} ^{pi} ^{pj} ^{pk} ^{pl} ^{pm} ^{pn} ^{po} ^{pp} ^{pq} ^{pr} ^{ps} ^{pt} ^{pu} ^{pv} ^{pw} ^{px} ^{py} ^{pz} ^{qa} ^{qb} ^{qc} ^{qd} ^{qe} ^{qf} ^{qg} ^{qh} ^{qi} ^{qj} ^{qk} ^{ql} ^{qm} ^{qn} ^{qo} ^{qp} ^{qq} ^{qr} ^{qs} ^{qt} ^{qu} ^{qv} ^{qw} ^{qx} ^{qy} ^{qz} ^{ra} ^{rb} ^{rc} rd ^{re} ^{rf} ^{rg} ^{rh} ^{ri} ^{rj} ^{rk} ^{rl} ^{rm} ^{rn} ^{ro} ^{rp} ^{rq} ^{rr} ^{rs} ^{rt} ^{ru} ^{rv} ^{rw} ^{rx} ^{ry} ^{rz} ^{sa} ^{sb} ^{sc} ^{sd} ^{se} ^{sf} ^{sg} ^{sh} ^{si} ^{sj} ^{sk} ^{sl} sm ^{sn} ^{so} ^{sp} ^{sq} ^{sr} ^{ss} st ^{su} ^{sv} ^{sw} ^{sx} ^{sy} ^{sz} ^{ta} ^{tb} ^{tc} ^{td} ^{te} ^{tf} ^{tg} th ^{ti} ^{tj} ^{tk} ^{tl} tm ^{tn} ^{to} ^{tp} ^{tq} ^{tr} ^{ts} ^{tt} ^{tu} ^{tv} ^{tw} ^{tx} ^{ty} ^{tz} ^{ua} ^{ub} ^{uc} ^{ud} ^{ue} ^{uf} ^{ug} ^{uh} ^{ui} ^{uj} ^{uk} ^{ul} ^{um} ^{un} ^{uo} ^{up} ^{uq} ^{ur} ^{us} ^{ut} ^{uu} ^{uv} ^{uw} ^{ux} ^{uy} ^{uz} ^{va} ^{vb} ^{vc} ^{vd} ^{ve} ^{vf} ^{vg} ^{vh} ^{vi} ^{vj} ^{vk} ^{vl} ^{vm} ^{vn} ^{vo} ^{vp} ^{vq} ^{vr} ^{vs} ^{vt} ^{vu} ^{vv} ^{vw} ^{vx} ^{vy} ^{vz} ^{wa} ^{wb} ^{wc} ^{wd} ^{we} ^{wf} ^{wg} ^{wh} ^{wi} ^{wj} ^{wk} ^{wl} ^{wm} ^{wn} ^{wo} ^{wp} ^{wq} ^{wr} ^{ws} ^{wt} ^{wu} ^{wv} ^{ww} ^{wx} ^{wy} ^{wz} ^{xa} ^{xb} ^{xc} ^{xd} ^{xe} ^{xf} ^{xg} ^{xh} ^{xi} ^{xj} ^{xk} ^{xl} ^{xm} ^{xn} ^{xo} ^{xp} ^{xq} ^{xr} ^{xs} ^{xt} ^{xu} ^{xv} ^{xw} ^{xx} ^{xy} ^{xz} ^{ya} ^{yb} ^{yc} ^{yd} ^{ye} ^{yf} ^{yg} ^{yh} ^{yi} ^{yj} ^{yk} ^{yl} ^{ym} ^{yn} ^{yo} ^{yp} ^{yq} ^{yr} ^{ys} ^{yt} ^{yu} ^{yv} ^{yw} ^{yx} ^{yy} ^{yz} ^{za} ^{zb} ^{zc} ^{zd} ^{ze} ^{zf} ^{zg} ^{zh} ^{zi} ^{zj} ^{zk} ^{zl} ^{zm} ^{zn} ^{zo} ^{zp} ^{zq} ^{zr} ^{zs} ^{zt} ^{zu} ^{zv} ^{zw} ^{zx} ^{zy} ^{zz} ^{aa} ^{ab} ^{ac} ^{ad} ^{ae} ^{af} ^{ag} ^{ah} ^{ai} ^{aj} ^{ak} ^{al} ^{am} ^{an} ^{ao} ^{ap} ^{aq} ^{ar} ^{as} ^{at} ^{au} ^{av} ^{aw} ^{ax} ^{ay} ^{az} ^{ba} ^{bb} ^{bc} ^{bd} ^{be} ^{bf} ^{bg} ^{bh} ^{bi} ^{bj} ^{bk} ^{bl} ^{bm} ^{bn} ^{bo} ^{bp} ^{bq} ^{br} ^{bs} ^{bt} ^{bu} ^{bv} ^{bw} ^{bx} ^{by} ^{bz} ^{ca} ^{cb} ^{cc} ^{cd} ^{ce} ^{cf} ^{cg} ^{ch} ^{ci} ^{cj} ^{ck} ^{cl} ^{cm} ^{cn} ^{co} ^{cp} ^{cq} ^{cr} ^{cs} ^{ct} ^{cu} ^{cv} ^{cw} ^{cx} ^{cy} ^{cz} ^{da} ^{db} ^{dc} ^{dd} ^{de} ^{df} ^{dg} ^{dh} ^{di} ^{dj} ^{dk} ^{dl} ^{dm} ^{dn} ^{do} ^{dp} ^{dq} ^{dr} ^{ds} ^{dt} ^{du} ^{dv} ^{dw} ^{dx} ^{dy} ^{dz} ^{ea} ^{eb} ^{ec} ^{ed} ^{ee} ^{ef} ^{eg} ^{eh} ^{ei} ^{ej} ^{ek} ^{el} ^{em} ^{en} ^{eo} ^{ep} ^{eq} ^{er} ^{es} ^{et} ^{eu} ^{ev} ^{ew} ^{ex} ^{ey} ^{ez} ^{fa} ^{fb} ^{fc} ^{fd} ^{fe} ^{ff} ^{fg} ^{fh} ^{fi}

- Other Societies on Cardiovascular Disease Prevention in Clinical Practice (Constituted by representatives of nine societies and by invited experts)". *European Heart Journal*. **28** (19): 2375–414. doi:10.1093/eurheartj/ehm316 . PMID 17726041 .
14. ^ "Gray's Anatomy of the Human Body – 6. Surface Markings of the Thorax" . Bartleby.com. Retrieved 2010-10-18.
 15. ^ Dorland's (2012). *Dorland's Illustrated Medical Dictionary* (32nd ed.). Elsevier. p. 1461. ISBN 978-1-4160-6257-8
 16. ^ <http://health.howstuffworks.com/human-body/systems/circulatory/heart1.htm>
 17. ^ ^a ^b Gray's Anatomy 2008, p. 960–962.
 18. ^ Gray's Anatomy 2008, p. 964–967.
 19. ^ Pocock, Gillian (2006). *Human Physiology*. Oxford University Press. p. 264. ISBN 978-0-19-856878-0.
 20. ^ ^a ^b ^c Gray's Anatomy 2008, p. 966–967.
 21. ^ Gray's Anatomy 2008, p. 970.
 22. ^ University of Minnesota. "Papillary Muscles" . *Atlas of Human Cardiac Anatomy* . Retrieved 7 March 2016.
 23. ^ "pectinate muscle" . Retrieved 2016-07-31.
 24. ^ Gray's Anatomy 2008, p. 959.
 25. ^ Davidson's 2010, p. 525.
 26. ^ Gray's Anatomy 2008, p. 981.
 27. ^ ^a ^b Gray's Anatomy 2008, p. 982.
 28. ^ Davidson's 2010, p. 526.
 29. ^ Gray's Anatomy 2008, p. 945.
 30. ^ "Main Frame Heart Development" . Meddean.luc.edu. Retrieved 2010-10-17.
 31. ^ DuBose, T. J.; Cunyus, J. A.; Johnson, L. (1990). "Embryonic Heart Rate and Age". *J Diagn Med Sonography* . **6** (3): 151–157. doi:10.1177/875647939000600306 .
 32. ^ DuBose, TJ (1996) *Fetal Sonography*, pp. 263–274; Philadelphia: WB Saunders ISBN 0-7216-5432-0
 33. ^ Terry J. DuBose *Sex, Heart Rate and Age*
 34. ^ Guyton & Hall 2011, pp. 110–113.
 35. ^ Pocock, Gillian (2006). *Human Physiology* (Third ed.). Oxford University Press. p. 266. ISBN 978-0-19-856878-0.
 36. ^ Antz, Matthias; et al. (1998). "Electrical Conduction Between the Right Atrium and the Left Atrium via the Musculature of the Coronary Sinus". *Circulation*. **98** (17): 1790–1795. doi:10.1161/01.CIR.98.17.1790 . PMID 9788835 .
 37. ^ De Ponti, Roberto; et al. (2002). "Electroanatomic Analysis of Sinus Impulse Propagation in Normal Human Atria". *Journal of Cardiovascular Electrophysiology* . **13** (1): 1–10. doi:10.1046/j.1540-8167.2002.00001.x . PMID 11843475 .
 38. ^ "SA node definition – Medical Dictionary definitions of popular medical terms easily defined on MedTerms" . Medterms.com. 27 April 2011 . Retrieved 7 June 2011
 39. ^ "Purkinje Fibers" . Biology.about.com. 9 April 2012 . Retrieved 7 June 2012.
 40. ^ Guyton & Hall 2011, pp. 115–120.
 41. ^ Davis, J. P.; Tikunova, S. B. (2008). "^{Ca}2+ exchange with troponin C and cardiac muscle dynamics" . *Cardiovascular Research*. **77** (4): 619–626. doi:10.1093/cvr/cvm098 . PMID 18079104 .
 42. ^ "Resting pulse rate reference data for children, adolescents and adults, United States 1999–2008" (PDF). Retrieved 30 December 2015.
 43. ^ Hall, Arthur C. Guyton, John E. (2005). *Textbook of medical physiology* (11th ed.). Philadelphia: W.B. Saunders. pp. 116–122. ISBN 978-0-7216-0240-0
 44. ^ Guyton & Hall 2011, p. 208.
 45. ^ Guyton & Hall 2011, p. 212.
 46. ^ ^a ^b Talley, Nicholas J.; O'Connor, Simon. *Clinical Examination*. Churchill Livingstone. pp. 76–82. ISBN 978-0-7295-4198-5
 47. ^ Dorland's (2012). *Dorland's Illustrated Medical Dictionary* (32nd ed.). Elsevier Saunders. p. 1189. ISBN 978-1-4160-6257-8
 48. ^ Newburger, Jane (2006). *Nadas' Pediatric Cardiology 2nd Edition*. Philadelphia: Elsevier. p. 358. ISBN 978-1-4160-2390-6
 49. ^ Cantarini L, Lopalco G; et al. (Oct 2014). "Autoimmunity and autoinflammation as the yin and yang of idiopathic recurrent acute pericarditis". *Autoimmun Rev*. **14** (2): 90–7. doi:10.1016/j.autrev.2014.10.005 . PMID 25308531 .
 50. ^ ^a ^b ^c "Cardiovascular diseases (CVDs)" . World Health Organization . Retrieved 2016-03-09.
 51. ^ "Your Heart Failure Healthcare Team" . www.heart.org. Retrieved 2016-03-09.
 52. ^ ^a ^b ^c "Different heart diseases | World Heart Federation" . www.world-heart-federation.org. Retrieved 2016-03-09.
 53. ^ Harrison's 2011, p. 1501.
 54. ^ Davidson's 2010, p. 554.
 55. ^ Davidson's 2010, p. 544.
 56. ^ Davidson's 2010, pp. 543–545.
 57. ^ Davidson's 2010, pp. 635–637.
 58. ^ ^a ^b ^c ^d Davidson's 2010, pp. 556–559.
 59. ^ Davidson's 2010, pp. 624–625.
 60. ^ Davidson's 2010, pp. 612–613.
 61. ^ ^a ^b ^c Davidson's 2010, pp. 560–570.
 62. ^ Davidson's 2010, p. 542.
 63. ^ Davidson's 2010, pp. 638–639.
 64. ^ "Patent Foramen Ovale (PFO)" . www.heart.org. Retrieved 2016-03-09.
 65. ^ ^a ^b Davidson's 2010, pp. 628–630.
 66. ^ Harrison's 2011, pp. 1458–1465.
 67. ^ Davidson's 2010, pp. 527–534.
 68. ^ ^a ^b Britton, the editors Nicki R. Colledge, Brian R. Walker, Stuart H. Ralston ; illustrated by Robert (2010). *Davidson's principles and practice of medicine*. (21st ed.). Edinburgh: Churchill Livingstone/Elsevier. pp. 522–536. ISBN 978-0-7020-3084-0
 69. ^ ^a ^b ^c ^d ^e ^f Davidson's 2010, pp. 522–536.
 70. ^ William F. Armstrong; Thomas Ryan; Harvey Feigenbaum (2010). *Feigenbaum's Echocardiography* . Lippincott Williams & Wilkins. ISBN 978-0-7817-9557-9
 71. ^ Gheorghade, M. (22 June 2004). "Digoxin in the Management of Cardiovascular Disorders". *Circulation*. **109** (24): 2959–2964. doi:10.1161/01.CIR.0000132482.95686.87 .
 72. ^ ^a ^b ^c ^d ^e Davidson's 2010, pp. 521–640.
 73. ^ ^a ^b Davidson's 2010, pp. 585–588, 614–623.
 74. ^ Davidson's 2010, pp. 585–588, 614–623.
 75. ^ ^a ^b ^c "Anatomy of the Heart" . University of Sydney Online Museum . Retrieved 2 August 2016.
 76. ^ ^a ^b ^c ^d Meletis, John; Konstantopoulos, Kostas (2010). "The Beliefs, Myths, and Reality Surrounding the Word Hema (Blood) from Homer to the Present". *Anemia*. **2010**: 1–6. doi:10.1155/2010/857657 .
 77. ^ Katz, A. M. (1 May 2008). "The "Modern" View of Heart Failure: How Did We Get Here?". *Circulation: Heart Failure* . **1** (1): 63–71. doi:10.1161/CIRCHEARTFAILURE.108.772756 .
 78. ^ ^a ^b ^c ^d ^e ^f AIRD, W. C. (July 2011). "Discovery of the cardiovascular system: from Galen to William Harvey". *Journal of Thrombosis and Haemostasis* . **9**: 118–129. doi:10.1111/j.1538-7836.2011.04312.x .
 79. ^ Michelakis, E. D. (19 June 2014). "Pulmonary Arterial Hypertension: Yesterday, Today, Tomorrow" . *Circulation Research*. **115** (1): 109–114. doi:10.1161/CIRCRESAHA.115.301132 .
 80. ^ West, John (2008). "Ibn al-Nafis, the pulmonary circulation, and the Islamic Golden Age" . *Journal of Applied Physiology* . **105** (6): 1877–1880. doi:10.1152/jappphysiol.91171.2008 . PMC 2612469 . PMID 18845773 . Retrieved 28 May 2014.
 81. ^ Bondke Persson, A.; Persson, P. B. (2014). "Form and function in the vascular system" . *Acta Physiologica*. **211** (3): 468–470. doi:10.1111/apha.12309 .
 82. ^ ^a ^b West, J. B. (30 May 2014). "Galen and the beginnings of Western physiology" . *AJP: Lung Cellular and Molecular Physiology* . **307** (2): L121–L128. doi:10.1152/ajplung.00123.2014 .
 83. ^ AIRD, W. C. (2011). "Discovery of the cardiovascular system: from Galen to William Harvey" . *Journal of Thrombosis and Haemostasis* . **9**: 118–129. doi:10.1111/j.1538-7836.2011.04312.x .
 84. ^ Silverman, M. E. (13 June 2006). "Why Does the Heart Beat?: The Discovery of the Electrical System of the Heart" . *Circulation*. **113** (23): 2775–2781. doi:10.1161/CIRCULATIONAHA.106.616771 . PMID 16769927 .
 85. ^ Cooley, Denton A. (2011). "Recollections of the Early Years of Heart Transplantation and the Total Artificial Heart" . *Artificial Organs*. **35** (4): 353–357.

- doi:10.1111/j.1525-1594.2011.01235.x . PMID 21501184 .
86. [^] Miniati, Douglas N.; Robbins, Robert C. (2002). "Heart transplantation: a thirty-year perspective: A Thirty-Year Perspective" . *Annual Review of Medicine* . **53** (1): 189–205. doi:10.1146/annurev.med.53.082901.104050 .
 87. [^] Neubauer, Stefan (15 March 2007). "The Failing Heart — An Engine Out of Fuel" . *New England Journal of Medicine* . **356** (11): 1140–1151. doi:10.1056/NEJMra063052 . PMID 17360992 .
 88. [^] ^a ^b ^c *The Watkins Dictionary of Symbols*. pp. "Heart". ISBN 978-1-78028-357-9.
 89. [^] Rosner, Fred (1995). *Medicine in the Bible and the Talmud : selections from classical Jewish sources* (Augm. ed.). Hoboken, NJ: KTAV Pub. House. pp. 87–96. ISBN 978-0-88125-506-5.
 90. [^] *Britannica*, *lb* ; *Slider*, *Ab*, *Egyptian heart and soul conception* . The word was also transcribed by *Wallis Budge* as *Ab*.
 91. [^] Allen, James P. (2000). *Middle Egyptian: An Introduction to the Language and Culture of Hieroglyphs* . Cambridge University Press. pp. 453, 465
 92. [^] Taylor, John H. (2001). *Death and the Afterlife in Ancient Egypt* . University of Chicago Press. pp. 35–38
 93. [^] Xigui, Qiu; Mattos, Gilbert L (2000). *Chinese writing = Wenzhi-xue-gaiyao* . Berkeley: Society for the Study of Early China [u.a.] p. 176. ISBN 1-55729-071-7.
 94. [^] MDBG online dictionary. "心". From <http://www.mdbg.net/chindict/chindict.php?page=worddict&wdrst=0&wdqb=%E5%BF%83> , accessed 7 February 2016.
 95. [^] Rogers, Bob Flaws, Michael Johnston & Timothy (2007). *Statements of fact in traditional Chinese medicine* (3rd ed.). Boulder, Colo.: Blue Poppy Press. p. 47. ISBN 978-0-936185-52-1.
 96. [^] Ye, Nigel Wiseman, Feng (1998). *A practical dictionary of Chinese medicine* (1st ed.). Brookline, Mass.: Paradigm Publications. p. 260. ISBN 978-0-912111-54-9.
 97. [^] Sven Sellmer (2004), "The Heart in the *Rg veda*", in Piotr Balcerowicz; Marek Mejer, *Essays in Indian Philosophy, Religion and Literature* , Delhi: Motilal Banarsidass Publishers, pp. 71–83, ISBN 978-81-208-1978-8
 98. [^] Lanman, Charles Rockwell (1996). *A Sanskrit reader : text and vocabulary and notes* (repr ed.). Delhi: Motilal Banarsidass. p. 287. ISBN 978-81-208-1363-2.
 99. [^] Aristotle. *On the Parts of Animals* . book 3, ch. 4 (*De partibus animalium*)
 100. [^] Galen, *De usu partium corporis humani* ("The Use of the Parts of the Human Body"), book 6.
 101. [^] Alan Sandstrom, *Corn is Our Life*, 1991, 239–240
 102. [^] *Nelson's Dictionary of Christianity: The Authoritative Resource on the Christian World* . Thomas Nelson Inc. 2001. p. Sacred Heart of Jesus. ISBN 978-1-4185-398-1 [first1= missing |last1= in Authors list (help)]
 103. [^] Murray, Tom Devonshire Jones; Linda Murray; Peter (2013). *The Oxford dictionary of christian art and architecture*. (Second ed.). Corby: Oxford University Press. p. Heart. ISBN 978-0-19-968027-6.
 104. [^] *Indonesia Magazine*, **25** (1994), p. 67
 105. [^] Samia Abdenmour, "Firakh mahshiya wi mihammara" recipe 117, *Egyptian Cooking: And Other Middle Eastern Recipes* , American University in Cairo Press, 2010.
 106. [^] *Diana Kennedy* , *My Mexico: A Culinary Odyssey with Recipes* , University of Texas Press, updated edition, 2013, ISBN 0-292-74840-X, p. 100
 107. [^] Alla Sacharow, *Classic Russian Cuisine: A Magnificent Selection of More Than 400 Traditional Recipes* , 1993, ISBN 1-55970-174-9, page unknown
 108. [^] ^a ^b *Irma S. Rombauer* , Marion Rombauer Becker, *The Joy of Cooking*, 1975, p. 508
 109. [^] Calvin W. Schwabe, *Unmentionable Cuisine*, University of Virginia Press, 1979 (reprint), ISBN 0-8139-1162-1, p. 96
 110. [^] John Torode, *Beef: And Other Bovine Matters* , Taunton Press, 2009, ISBN 1-60085-126-6, p. 230
 111. [^] Jennie Milsom, *The Connoisseur's Guide to Meat*, 2009m ISBN 1-4027-7050-2, p. 171
 112. [^] ^a ^b ^c ^d ^e ^f Romer, Alfred Sherwood; Parsons, Thomas S. (1977). *The Vertebrate Body*. Philadelphia, PA: Holt-Saunders International. pp. 437–442. ISBN 0-03-910284-X.
 113. [^] June Osborne (1998). *The Ruby-Throated Hummingbird*. University of Texas Press. p. 14. ISBN 0-292-76047-7.
 114. [^] Crigg, Gordon; Johansen, Kjell (1987). "Cardiovascular Dynamics In *Crocodylus Porosus* Breathing Air And During Voluntary Aerobic Dives" (PDF). *Journal of Comparative Physiology B*. Springer-Verlag. **157** (3): 381–392. doi:10.1007/BF00693365 . Archived from the original on 3 July 2012 . Retrieved 3 July 2012.
 115. [^] Axelsson, Michael; Craig, Franklin; Löfman, Carl; Nilsson, Stefan; Crigg, Gordon (1996). "Dynamic Anatomical Study Of Cardiac Shunting In Crocodiles Using High-Resolution Angioscopy" (PDF). *The Journal of Experimental Biology* . The Company of Biologists Limited. **199** (2): 359–365. PMID 9317958 . Retrieved 3 July 2012.

Bibliography

Hall, John (2011). *Guyton and Hall textbook of medical physiology* (12th ed.). Philadelphia, Pa.: Saunders/Elsevier. [ISBN 978-1-4160-4574-8](#).

Longo, Dan; Fauci, Anthony; Kasper, Dennis; Hauser, Stephen; Jameson, J.; Loscalzo, Joseph (August 11, 2011). *Harrison's Principles of Internal Medicine* (18 ed.). McGraw-Hill Professional. [ISBN 978-0-07-174889-6](#).

Susan Standing; Neil R. Borley; et al., eds. (2008). *Gray's anatomy : the anatomical basis of clinical practice* (40th ed.). London: Churchill Livingstone. [ISBN 978-0-8089-2371-8](#).

Nicki R. Colledge; Brian R. Walker; Stuart H. Ralston, eds. (2010). *Davidson's principles and practice of medicine* (21st ed.). Edinburgh: Churchill Livingstone/Elsevier. [ISBN 978-0-7020-3085-7](#).

External links

- What Is the Heart? – NIH
- The Gross Physiology of the Cardiovascular System (2nd Ed., 2012) – Robert M. Anderson, M.D. (CC-BY-NC)
- Atlas of Human Cardiac Anatomy
- Dissection review of the anatomy of the Human Heart including vessels, internal and external features
- Prenatal human heart development
- Anatomy of the Human Heart – Texas Heart Institute
- Animal hearts: fish, squid

Find more about **Heart** at Wikipedia's sister projects

Definitions from Wiktionary

Media from Commons

Quotations from Wikiquote

Texts from Wikisource

Textbooks from Wikibooks

Learning resources from Wikiversity

V • T • EAnatomy of the heart		
General	Surface	base • apex • sulci (coronary • interatrial • anterior interventricular • posterior interventricular) • borders (right • left)
	Internal	atria (interatrial septum • pectinate muscles • terminal sulcus) • ventricles (interventricular septum • trabeculae carneae • chordae tendineae • papillary muscle) • valves • cusps • atrioventricular septum
		cardiac skeleton • intervenous tubercle
Chambers	Right heart	(<i>venae cavae</i> , <i>coronary sinus</i>) → right atrium (atrial appendage, fossa ovalis, limbus of fossa ovalis, crista terminalis, valve of inferior vena cava , valve of coronary sinus) → tricuspid valve → right ventricle (infundibulum, moderator band/septomarginal trabecula) → pulmonary valve → (<i>pulmonary artery and pulmonary circulation</i>)
	Left heart	(<i>pulmonary veins</i>) → left atrium (atrial appendage) → mitral valve → left ventricle → aortic valve (aortic sinus) → (<i>aorta and systemic circulation</i>)
Layers	Endocardium	heart valves
	Myocardium	Conduction system (cardiac pacemaker • SA node • AV node • bundle of His • bundle branches • Purkinje fibers)
	Pericardial cavity	pericardial sinus
	Pericardium	fibrous pericardium (sternopericardial ligaments) • serous pericardium (epicardium/visceral layer) • fold of left vena cava
Blood supply	Coronary circulation	

V • T • EPhysiology of the cardiovascular system		
Heart	Cardiac output	Cardiac cycle • Cardiac output (Heart rate • Stroke volume) • Stroke volume (End-diastolic volume • End-systolic volume) • Afterload • Preload • Frank–Starling law of the heart • Cardiac function curve • Venous return curve • Wiggers diagram • Pressure volume diagram
	Ultrasound	Fractional shortening = (End-diastolic dimension • End-systolic dimension) / End-diastolic dimension • Aortic valve area calculation • Ejection fraction • Cardiac index
	Heart rate	Cardiac pacemaker • Chronotropic (Heart rate) • Dromotropic (Conduction velocity) • Inotropic (Contractility) • Bathmotropic (Excitability) • Lusitropic (Relaxation)
	Conduction	Conduction system • Cardiac electrophysiology • Action potential (cardiac • atrial • ventricular) • Effective refractory period • Pacemaker potential • Electrocardiography (P wave • PR interval • QRS complex • QT interval • ST segment • T wave • U wave) • Hexaxial reference system
	Chamber pressure	Central venous • Right (atrial • ventricular) • pulmonary artery (wedge) • Left (atrial • ventricular) • Aortic
	Other	Ventricular remodeling
Vascular system / Hemodynamics	Blood flow	Compliance • Vascular resistance • Pulse • Perfusion
	Blood pressure	Pulse pressure (Systolic • Diastolic) • Mean arterial pressure • Jugular venous pressure • Portal venous pressure
	Regulation of BP	Baroreflex • Kinin–kallikrein system • Renin–angiotensin system • Vasoconstrictors • Vasodilators • Autoregulation (Myogenic mechanism • Tubuloglomerular feedback • Cerebral autoregulation) • Paraganglia (Aortic body • Carotid body • Glomus cell)
Authority control	GND: 4024632-2 • NDL: 00571194	

Categories: Heart | Cardiac anatomy | Organs (anatomy)