

Article
Talk

Read
View source
View history

Search Wikipedia

# Heart

From Wikipedia, the free encyclopedia

 *This article is about the internal organ. For other uses, see *Heart (disambiguation)*.*

The **heart** is a **muscular organ** in most animals, which pumps **blood** through the **blood vessels** of the **circulatory system**.<sup>[1]</sup> Blood provides the body with **oxygen** and **nutrients**, as well as assists in the removal of **metabolic wastes**.<sup>[2]</sup> In humans, the heart is located between the lungs, in the **middle compartment** of the **chest**.<sup>[3]</sup>

In humans, other mammals, and birds, the heart is divided into four chambers: upper left and right **atria**; and lower left and right **ventricles**.<sup>[4][5]</sup> Commonly the right atrium and ventricle are referred together as the *right heart* and their left counterparts as the *left heart*.<sup>[6]</sup> Fish, in contrast, have two chambers, an atrium and a ventricle, while reptiles have three chambers.<sup>[5]</sup> In a healthy heart blood flows one way through the heart due to **heart valves**, which prevent **backflow**.<sup>[3]</sup> The heart is enclosed in a protective sac, the **pericardium**, which also contains a small amount of **fluid**. The wall of the heart is made up of three layers: **epicardium**, **myocardium**, and **endocardium**.<sup>[7]</sup>

The heart pumps blood with a rhythm determined by a group of pacemaking cells in the **sinoatrial node**. These generate a current that causes contraction of the heart, traveling through the **atrioventricular node** and along the **conduction system of the heart**. The heart receives blood low in oxygen from the **systemic circulation**, which enters the right atrium from the **superior** and **inferior venae cavae** and passes to the right ventricle. From here it is pumped into the **pulmonary circulation**, through the **lungs** where it receives oxygen and gives off carbon dioxide. Oxygenated blood then returns to the left atrium, passes through the left ventricle and is pumped out through the **aorta** to the systemic circulation—where the oxygen is used and **metabolized** to **carbon dioxide**.<sup>[8]</sup> The heart beats at a resting **rate** close to 72 beats per minute.<sup>[9]</sup> **Exercise** temporarily increases the rate, but lowers **resting heart rate** in the long term, and is good for heart health.<sup>[10]</sup>

**Cardiovascular diseases** (CVD) are the most common cause of death globally as of 2008, accounting for 30% of deaths.<sup>[11][12]</sup> Of these more than three quarters are a result of **coronary artery disease** and **stroke**.<sup>[11]</sup> Risk factors include: **smoking**, being **overweight**, little exercise, **high cholesterol**, **high blood pressure**, and poorly controlled **diabetes**, among others.<sup>[13]</sup> Cardiovascular diseases frequently do not have symptoms or may cause **chest pain** or **shortness of breath**. Diagnosis of heart disease is often done by the taking of a **medical history**, **listening** to the **heart-sounds** with a **stethoscope**, **ECG**, and **ultrasound**.<sup>[3]</sup> Specialists who focus on diseases of the heart are called **cardiologists**, although many specialties of medicine may be involved in treatment.<sup>[12]</sup>

## Contents

- 1

Structure
- 1.1

Location and shape

1.2

Chambers

1.3

Heart wall

1.4

Pericardium

1.5

Coronary circulation

1.6

Nerve supply
- 2

Development
- 3

Physiology
- 3.1

Blood flow

3.2

Electrical conduction

3.3

Heart rate

4

Clinical significance

4.1

Diseases

4.2

Diagnosis

4.3

Treatment

5

History

5.1

Ancient

5.2

Pre-modern

5.3

Modern

6

Society and culture

6.1

Symbolism

6.2

Food

7

Other animals

7.1

Other vertebrates

7.2

Double circulatory systems

7.3

The fully divided heart

7.4

Fish

7.5

Invertebrates

8

Additional images

9

Notes

10

References

11

Bibliography

## Heart

The human heart

### Details

<b>System</b>	Circulatory
<b>Artery</b>	Aorta, <sup>[a]</sup> <b>pulmonary trunk</b> and right and left <b>pulmonary arteries</b> <sup>[b]</sup> <b>Right coronary artery</b> , <b>left main coronary artery</b> <sup>[c]</sup>
<b>Vein</b>	<b>Superior vena cava</b> , <b>inferior vena cava</b> , <sup>[d]</sup> <b>right</b> and <b>left pulmonary veins</b> , <sup>[e]</sup> <b>great cardiac vein</b> , <b>middle cardiac vein</b> , <b>small cardiac vein</b> , <b>anterior cardiac veins</b> . <sup>[f]</sup>
<b>Nerve</b>	<b>Accelerans nerve</b> , <b>vagus nerve</b>

### Identifiers

<b>Latin</b>	<i>cor</i>
<b>Greek</b>	<i>kardía</i> ( <i>καρδιά</i> )
<b>MeSH</b>	<i>A07.541</i>
<b>TA</b>	A12.1.00.001
<b>FMA</b>	7088

### Anatomical terminology

[edit on Wikidata]

### Normal heart sounds

Normal heart sounds as heard with a stethoscope

*Problems playing this file? See media help.*

Main page
Contents
Featured content
Current events
Random article
Donate to Wikipedia
Wikipedia store

Interaction

Help
About Wikipedia
Community portal
Recent changes
Contact page

Tools

What links here
Related changes
Upload file
Special pages
Permanent link
Page information
Wikidata item
Cite this page

Print/export

Create a book
Download as PDF
Printable version

In other projects

Wikimedia Commons
Wikiquote

Languages

Afrikaans
Alemannisch
አማርኛ
Ænglisc
العربية
Aragonés
ܐܪܡܝܐ
Armãneashti
অসমীয়া
Asturianu
Atikamekw
Avañe'ẽ
Авар
Aymar aru
Azərbaycanca
Башҡортса
Беларуская
Беларуская (тарашкевіца)
Bikol Central
Български
Bosanski
Brezhoneg
Буряад
Català
ЧӀавашла
Cebuano
Čeština
Chavacano de Zamboanga
ChiShona
Corsu
Cymraeg
Dansk
Deutsch

Deutsch • Deutsch • Deutsch • Deutsch

Eesti
Ελληνικά
Emiliàn e rumagnòl
Español
Esperanto
Euskara
فارسی
Fiji Hindi
Føroyskt
Français
Frysk
Gaeilge
Galego
गोंयची कोंकणी / Göychi

- Konknni

客家語/Hak-kâ-ngî

Халъмр

한국어

Հայերեն

हिन्दी

Hrvatski

Ido

Ilokano

Bahasa Indonesia

Interlingua

Interlingue

IsiXhosa

Íslenska

Italiano

עברית

Basa Jawa

ಕನ್ನಡ

Kapampangan

ქართული

Қазақша

Kiswahili

Кomi

Kreyòl ayisyen

Кыргызча

Кырык мары

Лакку

Latina

Latviešu

Lëtzebuergesch

Lietuvių

Limburgs

Lingála

La .lojban.

Lumbaart

Magyar

Македонски

മലയാളം

Malti

मराठी

მარგალური

مصرى

Bahasa Melayu

Ming-dêng-ngû

Монгол

မၤနଁဟံသာဝါ

Nāhuatl

Nederlands

नेपाली

नेपाल भाषा

日本語

Napulitano

Нохчийн

Norsk

Norsk nynorsk

Nouormand

Occitan

Олык мари

ଓଡ଼ିଆ

O'zbekcha/Ўзбекча

ਪੰਜਾਬੀ

Pangasinan

پنجابی

بيننوو

Plattdüütsch

Polski

Português

Română

Runa Simi

Русиньскый

Русский

Саха тыла

संस्कृतम्

Sardu

Scots

Shqip

Sicilianu

සිංහල

Simple English

Slovenčina

Slovenščina

Ślůnski

Soomaaliga

کوردی

Српски / srpski

Srpskohrvatski / српскохрватски

Basa Sunda

Suomi

Svenska

Tagalog

தமிழ்

Татарча/tatarça

తెలుగు

ไทย

ትግርኛ

Тоҷикӣ

Türkçe

Українська

اردو

ئۇيغۇرچە / Uyghurche

Vahcuengh

- 12 External links

## Structure

### Location and shape

The human heart is situated in the **middle mediastinum** , at the level of **thoracic vertebrae T5-T8**. A double-membraned sac called the **pericardium** surrounds the heart and attaches to the mediastinum.<sup>[15]</sup> The back surface of the heart lies near the **vertebral column**, and the front surface sits behind the **sternum** and **rib cartilages**.<sup>[7]</sup> The upper part of the heart is the attachment point for several large blood vessels – the **venae cavae**, **aorta** and **pulmonary trunk**. The upper part of the heart is located at the level of the third costal cartilage.<sup>[7]</sup> The lower tip of the heart, the apex, lies to the left of the sternum (8 to 9 cm from the **midsternal line**) between the junction of the fourth and fifth ribs near their **articulation** with the costal cartilages.<sup>[7]</sup>

The largest part of the heart is usually slightly offset to the left side of the chest (though occasionally it may be **offset to the right** ) and is felt to be on the left because the **left heart** is stronger and larger, since it pumps to all body parts. Because the heart is between the **lungs**, the left lung is smaller than the right lung and has a cardiac notch in its border to accommodate the heart.<sup>[7]</sup> The heart is cone-shaped, with its base positioned upwards and tapering down to the apex.<sup>[7]</sup> An adult heart has a mass of 250–350 grams (9–12 oz).<sup>[16]</sup> The heart is typically the size of a fist: 12 cm (5 in) in length, 8 cm (3.5 in) wide, and 6 cm (2.5 in) in thickness.<sup>[7]</sup> Well-trained **athletes** can have much larger hearts due to the effects of exercise on the heart muscle, similar to the response of skeletal muscle.<sup>[7]</sup>

### Chambers

The heart has four chambers, two upper **atria**, the receiving chambers, and two lower **ventricles**, the discharging chambers. The atria open into the ventricles via the atrioventricular valves, present in the **atrioventricular septum**. This distinction is visible also on the surface of the heart as the **coronary sulcus**.<sup>[17]</sup> There is an ear-shaped structure in the upper right atrium called the right atrial appendage, or auricle, and another in the upper left atrium, the left atrial appendage.<sup>[18]</sup> The right atrium and the right ventricle together are sometimes referred to as the *right heart*. Similarly, the left atrium and the left ventricle together are sometimes referred to as the *left heart*.<sup>[6]</sup> The ventricles are separated from each other by the **interventricular septum**, visible on the surface of the heart as the **anterior longitudinal sulcus** and the **posterior interventricular sulcus**.<sup>[17]</sup>

The **cardiac skeleton** is made of **dense connective tissue** and this gives structure to the heart. It forms the **atrioventricular septum** which separates the atria from the ventricles, and the fibrous rings which serve as bases for the four **heart valves**.<sup>[19]</sup> The cardiac skeleton also provides an important boundary in the heart's electrical conduction system since collagen cannot conduct electricity. The interatrial septum separates the atria and the interventricular septum separates the ventricles.<sup>[7]</sup> The interventricular septum is much thicker than the interatrial septum, since the ventricles need to generate greater pressure when they contract.<sup>[7]</sup>

#### Valves

*Main article: **Heart valves***

The heart has four valves, which separate its chambers. One valve lies between each atrium and ventricle, and one valve rests at the exit of each ventricle.<sup>[7]</sup>







The valves between the atria and ventricles are called the atrioventricular valves. Between the right atrium and the right ventricle is the **tricuspid valve**. The tricuspid valve has three cusps,<sup>[20]</sup> which connect to chordae tendinae and three papillary muscles named the anterior, posterior, and septal muscles, after their relative positions.<sup>[20]</sup> The **mitral valve** lies between the left atrium and left ventricle. It is also known as the bicuspid valve due to its having two cusps, an anterior and a posterior cusp. These cusps are also attached via chordae tendinae to two papillary muscles projecting from the ventricular wall.<sup>[21]</sup>

The **papillary muscles** extend from the walls of the heart to valves by cartilaginous connections called **chordae tendinae**. These muscles prevent the valves from falling too far back when they close.<sup>[22]</sup> During the relaxation phase of the cardiac cycle, the papillary muscles are also relaxed and the tension on the chordae tendineae is slight. As the heart chambers contract, so do the papillary muscles. This creates tension on the chordae tendineae, helping to hold the cusps of the atrioventricular valves in place and preventing them from being blown back into the atria.<sup>[7]</sup> <sup>[g]</sup><sup>[20]</sup>

Two additional semilunar valves sit at the exit of each of the ventricles. The **pulmonary valve** is located at the base of the **pulmonary artery**. This has three cusps which are not attached to any papillary muscles. When the ventricle relaxes blood flows back into the ventricle from the artery and this flow of blood fills the pocket-like valve, pressing against the cusps which close to seal the valve. The semilunar **aortic valve** is at the base of the **aorta** and also is not attached to papillary muscles. This too has three cusps which close with the pressure of the blood flowing back from the aorta.<sup>[7]</sup>

#### Right heart

The right heart consists of two chambers, the right atrium and the right ventricle, separated by a valve, the **tricuspid valve**.<sup>[7]</sup>



The heart, showing valves, arteries and veins. The white

The right atrium receives blood almost continuously from the body's two major veins, the [superior](#) and [inferior venae cavae](#). A small amount of blood from the coronary circulation also drains into the right atrium via the [coronary](#)

[sinus](#), which is immediately above and to the middle of the opening of the inferior vena cava.<sup>[7]</sup> In the wall of the right atrium is an oval-shaped depression known as the [fossa ovalis](#), which is a remnant of an opening in the fetal heart known as the [foramen ovale](#).<sup>[7]</sup> Most of the internal surface of the right atrium is smooth, the depression of the fossa ovalis is medial, and the anterior surface has prominent ridges of [pectinate muscles](#), which are also present in the [right atrial appendage](#).<sup>[7]</sup>

The right atrium is connected to the right ventricle by the [tricuspid valve](#).<sup>[7]</sup> The walls of the right ventricle are lined with [trabeculae carneae](#), ridges of cardiac muscle covered by endocardium. In addition to these muscular ridges, a band of cardiac muscle, also covered by endocardium, known as the [moderator band](#) reinforces the thin walls of the right ventricle and plays a crucial role in cardiac conduction. It arises from the lower part of the interventricular septum and crosses the interior space of the right ventricle to connect with the inferior papillary muscle.<sup>[7]</sup> The right ventricle tapers into the [pulmonary trunk](#), into which it ejects blood when contracting. The pulmonary trunk branches into the left and right pulmonary arteries that carry the blood to each lung. The pulmonary valve lies between the right heart and the pulmonary trunk.<sup>[7]</sup>

#### Left heart

The left heart has two chambers: the left atrium, and the left ventricle, separated by the [mitral valve](#).<sup>[7]</sup>

The left atrium receives oxygenated blood back from the lungs via one of the four [pulmonary veins](#). The left atrium has an outpouching called the [left atrial appendage](#). Like the right atrium, the left atrium is lined by [pectinate muscles](#).<sup>[23]</sup> The left atrium is connected to the left ventricle by the [mitral valve](#).<sup>[7]</sup>

The left ventricle is much thicker as compared with the right, due to the greater force needed to pump blood to the entire body. Like the right ventricle, the left also has [trabeculae carneae](#), but there is no [moderator band](#). The left ventricle pumps blood to the body through the [aortic valve](#) and into the [aorta](#). Two small openings above the aortic valve carry blood to the heart itself, the [left main coronary artery](#) and the [right coronary artery](#).<sup>[7]</sup>

#### Heart wall

*Further information:* [Cardiac muscle](#)

The heart wall is made up of three layers: the inner [endocardium](#), middle [myocardium](#) and outer [epicardium](#). These are surrounded by a double-membraned sac called the [pericardium](#).

The innermost layer of the heart is called the endocardium. It is made up of [simple squamous epithelium](#), and covers heart chambers and valves. It is continuous with the [endothelium](#) of the veins and arteries of the heart, and is joined to the myocardium with a thin layer of connective tissue.<sup>[7]</sup> The endocardium, by secreting [endothelins](#), may also play a role in regulating the contraction of the myocardium.<sup>[7]</sup>

The middle layer of the heart wall is the myocardium, which is the [cardiac muscle](#) – a layer of involuntary [striated muscle tissue](#) surrounded by a framework of [collagen](#). The cardiac muscle pattern is elegant and complex, as the muscle cells swirl and spiral around the chambers of the heart, with the outer muscles forming a figure 8 pattern around the atria and around the bases of the great vessels, and inner muscles forming a figure 8 around the two ventricles and proceed toward the apex. This complex swirling pattern allows the heart to pump blood more effectively.<sup>[7]</sup>

There are two types of cells in cardiac muscle: [muscle cells](#) which have the ability to contract easily, and [pacemaker cells](#) of the conducting system. The muscle cells make up the bulk (99%) of cells in the atria and ventricles. These contractile cells are connected by [intercalated discs](#) which allow a rapid response to impulses of [action potential](#) from the pacemaker cells. The intercalated discs allow the cells to act as a [syncytium](#) and enable the contractions that pump blood through the heart and into the [major arteries](#).<sup>[7]</sup> The pacemaker cells make up 1% of cells and form the conduction system of the heart. They are generally much smaller than the contractile cells and have few [myofibrils](#) which gives them limited contractibility. Their function is similar in many respects to [neurons](#).<sup>[7]</sup> Cardiac muscle tissue has [autorhythmicity](#), the unique ability to initiate a [cardiac action potential](#) at a fixed rate – spreading the impulse rapidly from cell to cell to trigger the contraction of the entire heart.<sup>[7]</sup>

There are specific [proteins expressed](#) in cardiac muscle cells.<sup>[24][25]</sup> These are mostly associated with muscle contraction, and bind with [actin](#), [myosin](#), [tropomyosin](#), and [troponin](#). They include [MYH6](#), [ACTC1](#), [TNNI3](#), [CDH2](#) and [PKP2](#). Other proteins expressed are [MYH7](#) and [LDB3](#) that are also expressed in skeletal muscle.<sup>[26]</sup>

#### Pericardium

The [pericardium](#) is the sack that surrounds the heart. The tough outer surface of the pericardium is called the fibrous membrane. This is lined by a double inner membrane called the serous membrane that produces [pericardial fluid](#) to lubricate the surface of the heart.<sup>[27]</sup> The part of the serous membrane attached to the fibrous membrane is called the parietal pericardium, while the part of the serous membrane attached to the heart is known as the visceral pericardium. The pericardium is present in order to lubricate its movement against other structures within the chest, to keep the heart's position stabilised within the chest, and to protect the heart from infection.<sup>[28]</sup>

#### Coronary circulation

*Main article:* [Coronary circulation](#)

Heart tissue, like all cells in the body, needs to be supplied with [oxygen](#), [nutrients](#) and a way of removing [metabolic wastes](#). This is achieved by the [coronary circulation](#), which includes [arteries](#), [veins](#), and [lymphatic vessels](#). Blood flow through the coronary vessels occurs in peaks and troughs relating to the heart muscle's relaxation or contraction.<sup>[7]</sup>

Heart tissue receives blood from two arteries which arise just above the aortic valve. These are the [left main coronary artery](#) and the [right coronary artery](#). The left main coronary artery splits shortly after leaving the [aorta](#) into two vessels, the [left anterior descending](#) and the [left circumflex artery](#). The left anterior descending artery supplies heart tissue and the front, outer side, and the septum of the left ventricle. It does this by

arrows show the normal direction of blood flow.

Frontal section showing papillary muscles attached to the tricuspid valve on the right and to the mitral valve on the left via chordae tendineae.<sup>[7]</sup>

Layers of the heart wall, including visceral and parietal pericardium.

The swirling pattern of myocardium helps the heart pump effectively

Arterial supply to the heart (red), with other areas labelled (blue).

branching into smaller arteries – diagonal and septal branches. The left circumflex supplies the back and underneath of the left ventricle. The right coronary artery supplies the right atrium, right ventricle, and lower posterior sections of the left ventricle. The right coronary artery also supplies blood to the atrioventricular node (in about 90% of people) and the sinoatrial node (in about 60% of people). The right coronary artery runs in a groove at the back of the heart and the left anterior descending artery runs in a groove at the front. There is significant variation between people in the anatomy of the arteries that supply the heart <sup>[29]</sup> The arteries divide at their furthest reaches into smaller branches that join together at the edges of each arterial distribution.<sup>[7]</sup>

The **coronary sinus** is a large vein that drains into the right atrium, and receives most of the venous drainage of the heart. It receives blood from the **great cardiac vein** (receiving the left atrium and both ventricles), the **posterior cardiac vein** (draining the back of the left ventricle), the **middle cardiac vein** (draining the bottom of the left and right ventricles), and **small cardiac veins**.<sup>[30]</sup> The **anterior cardiac veins** drain the front of the right ventricle and drain directly into the right atrium.<sup>[7]</sup>

Small lymphatic networks called **plexuses** exist beneath each of the three layers of the heart. These networks collect into a main left and a main right trunk, which travel up the groove between the ventricles that exists on the heart's surface, receiving smaller vessels as they travel up. These vessels then travel into the atrioventricular groove, and receive a third vessel which drains the section of the left ventricle sitting on the diaphragm. The left vessel joins with this third vessel, and travels along the pulmonary artery and left atrium, ending in the **inferior tracheobronchial node**. The right vessel travels along the right atrium and the part of the right ventricle sitting on the diaphragm. It usually then travels in front of the ascending aorta and then ends in a brachiocephalic node.<sup>[31]</sup>

## Nerve supply

The heart receives nerve signals from the **vagus nerve** and from nerves arising from the **sympathetic trunk**. These nerves act to influence, but not control, the heart rate. **Sympathetic nerves** also influence the force of heart contraction.<sup>[32]</sup> Signals that travel along these nerves arise from two paired **cardiovascular centres** in the **medulla oblongata**. The **vagus nerve** of the parasympathetic nervous system acts to decrease the heart rate, and nerves from the **sympathetic trunk** act to increase the heart rate.<sup>[7]</sup> These nerves form a network of nerves that lies over the heart called the **cardiac plexus**.<sup>[7][31]</sup>

The vagus nerve is a long, wandering nerve that emerges from the **brainstem** and provides parasympathetic stimulation to a large number of organs in the thorax and abdomen, including the heart.<sup>[33]</sup> The nerves from the sympathetic trunk emerge through the T1-T4 **thoracic ganglia** and travel to both the sinoatrial and atrioventricular nodes, as well as to the atria and ventricles. The ventricles are more richly innervated by sympathetic fibers than parasympathetic fibers. Sympathetic stimulation causes the release of the neurotransmitter **norepinephrine** (also known as **noradrenaline**) at the **neuromuscular junction** of the cardiac nerves. This shortens the repolarization period, thus speeding the rate of depolarization and contraction, which results in an increased heart rate. It opens chemical or ligand-gated sodium and calcium ion channels, allowing an influx of **positively charged ions**.<sup>[7]</sup> Norepinephrine binds to the **beta-1 receptor**.<sup>[7]</sup>

## Development

*Main articles: **Heart development** and **Human embryogenesis***

The heart is the first functional organ to develop and starts to beat and pump blood at about three weeks into **embryogenesis**. This early start is crucial for subsequent embryonic and **prenatal development**.

The heart derives from **splanchnopleuric mesenchyme** in the neural plate which forms the cardiogenic region.

Two **endocardial tubes** form here that fuse to form a primitive heart tube known as the **tubular heart**.<sup>[34]</sup> Between the third and fourth week, the heart tube lengthens, and begins to fold to form an S-shape within the pericardium. This places the chambers and major vessels into the correct alignment for the developed heart. Further development will include the septa and valves formation and remodelling of the heart chambers. By the end of the fifth week the septa are complete and the heart valves are completed by the ninth week.<sup>[7]</sup>

Before the fifth week, there is an opening in the fetal heart known as the **foramen ovale**. The foramen ovale allowed blood in the fetal heart to pass directly from the right atrium to the left atrium, allowing some blood to bypass the lungs. Within seconds after birth, a flap of tissue known as the **septum primum** that previously acted as a valve closes the foramen ovale and establishes the typical cardiac circulation pattern. A depression in the surface of the right atrium remains where the foramen ovale once walls, called the fossa ovalis.<sup>[7]</sup>

The **embryonic** heart begins beating at around 22 days after conception (5 weeks after the last normal menstrual period, LMP). It starts to beat at a rate near to the mother's which is about 75–80 **beats per minute** (bpm). The embryonic heart rate then accelerates and reaches a peak rate of 165–185 bpm early in the early 7th week (early 9th week after the LMP).<sup>[35][36]</sup> After 9 weeks (start of the **fetal** stage) it starts to decelerate, slowing to around 145 (±25) bpm at birth. There is no difference in female and male heart rates before birth.<sup>[37]</sup>

## Physiology

*Main article: **Cardiac physiology***

### Blood flow

The heart functions as a pump in the **circulatory system** to provide a continuous **flow of blood** throughout the body. This circulation consists of the **systemic circulation** to and from the body and the **pulmonary circulation** to and from the lungs. Blood in the pulmonary circulation exchanges carbon dioxide for oxygen in the lungs through the process of **respiration**. The systemic circulation then transports oxygen to the body and returns carbon

Autonomic innervation of the heart

Development of the human heart during the first eight weeks (top), and the formation of the heart chambers (bottom). In this figure, the blue and red colors represent blood inflow and outflow (not venous and arterial blood). Initially, all venous blood flows from the tail/atria to the ventricles/head, a very different pattern from that of an adult.<sup>[7]</sup>

Blood flow through the valves

dioxide and relatively deoxygenated blood to the heart for transfer to the lungs.<sup>[7]</sup>

The **right heart** collects deoxygenated blood from two large veins, the **superior** and **inferior venae cavae**. Blood collects in the right and left atrium continuously.<sup>[7]</sup> The superior vena cava drains blood from above the **diaphragm** and empties into the upper back part of the right atrium. The inferior vena cava drains the blood from below the diaphragm and empties into the back part of the atrium below the opening for the superior vena cava. Immediately above and to the middle of the opening of the inferior vena cava is the opening of the thin-walled coronary sinus.<sup>[7]</sup> Additionally, the **coronary sinus** returns deoxygenated blood from the myocardium to the right atrium. The blood collects in the right atrium. When the right atrium contracts, the blood is pumped through the **tricuspid**

Play media

Blood flow through the heart from the Khan academy

**valve** into the right ventricle. As the right ventricle contracts, the tricuspid valve closes and the blood is pumped into the pulmonary trunk through the **pulmonary valve**. The pulmonary trunk divides into pulmonary arteries and progressively smaller arteries throughout the lungs, until it reaches **capillaries**. As these pass by **alveoli** carbon dioxide is **exchanged** for oxygen. This happens through the passive process of **diffusion**.

In the **left heart**, oxygenated blood is returned to the left atrium via the **pulmonary veins**. It is then pumped into the left ventricle through the **mitral valve** and into the aorta through the **aortic valve** for systemic circulation. The aorta is a large artery that branches into many smaller arteries, **arterioles**, and ultimately **capillaries**. In the capillaries, oxygen and nutrients from blood are supplied to body cells for metabolism, and exchanged for carbon dioxide and waste products.<sup>[7]</sup> Capillary blood, now deoxygenated, travels into **venules** and **veins** that ultimately collect in the superior and inferior vena cavae, and into the right heart.

### Cardiac cycle

*Main articles: **Cardiac cycle**, **Systole**, and **Diastole***

The **cardiac cycle** refers to the sequence of events in which the heart contracts and relaxes with every heartbeat.<sup>[9]</sup> The period of time during which the ventricles contract, forcing blood out into the aorta and main pulmonary artery, is known as **systole**, while the period during which the ventricles relax and refill with blood is known as **diastole**. The atria and ventricles work in concert, so in systole when the ventricles are contracting, the atria are relaxed and collecting blood. When the ventricles are relaxed in diastole, the atria contract to pump blood to the ventricles. This coordination ensures blood is pumped efficiently to the body.<sup>[7]</sup>

At the beginning of the cardiac cycle, the ventricles are relaxing. As they do so, they are filled by blood passing through the open **mitral** and **tricuspid** valves. After the ventricles have completed most of their filling, the atria contract, forcing further blood into the ventricles and priming the pump. Next, the ventricles start to contract. As the pressure rises within the cavities of the ventricles, the mitral and tricuspid valves are forced shut. As the pressure within the ventricles rises further, exceeding the pressure with the **aorta** and **pulmonary** arteries, the **aortic** and **pulmonary** valves open. Blood is ejected from the heart, causing the pressure within the ventricles to fall. Simultaneously, the atria refill as blood flows into the **right atrium** through the superior and **inferior vena cavae**, and into the **left atrium** through the **pulmonary veins**. Finally, when the pressure within the ventricles falls below the pressure within the aorta and pulmonary arteries, the aortic and pulmonary valves close. The ventricles start to relax, the mitral and tricuspid valves open, and the cycle begins again.<sup>[9]</sup>

The cardiac cycle as correlated to the ECG

### Cardiac output

*Main article: **Cardiac output***

**Cardiac output** (CO) is a measurement of the amount of blood pumped by each ventricle (stroke volume) in one minute. This is calculated by multiplying the stroke volume (SV) by the beats per minute of the heart rate (HR). So that:  $CO = SV \times HR$ .<sup>[7]</sup> The cardiac output is normalized to body size through **body surface area** and is called the **cardiac index**.

The average cardiac output, using an average stroke volume of about 70mL, is 5.25 L/min, with a normal range of 4.0–8.0 L/min.<sup>[7]</sup> The stroke volume is normally measured using an **echocardiogram** and can be influenced by the size of the heart, physical and mental condition of the individual, **sex**, **contractility**, duration of contraction, **preload** and **afterload**.<sup>[7]</sup>

**Preload** refers to the filling pressure of the atria at the end of diastole, when they are at their fullest. A main factor is how long it takes the ventricles to fill—if the ventricles contract faster, then there is less time to fill and the preload will be less.<sup>[7]</sup> Preload can also be affected by a person's blood volume. The force of each contraction of the heart muscle is proportional to the preload, described as the **Frank-Starling mechanism**. This states that the force of contraction is directly proportional to the initial length of muscle fiber, meaning a ventricle will contract more forcefully, the more it is stretched.<sup>[7][38]</sup>

The x-axis reflects time with a recording of the heart sounds. The y-axis represents pressure.<sup>[7]</sup>

**Afterload**, or how much pressure the heart must generate to eject blood at systole, is influenced by **vascular resistance**. It can be influenced by narrowing of the heart valves (**stenosis**) or contraction or relaxation of the peripheral blood vessels.<sup>[7]</sup>

The strength of heart muscle contractions controls the stroke volume. This can be influenced positively or negatively by agents termed **inotropes**.<sup>[39]</sup> These agents can be a result of changes within the body, or be given as drugs as part of treatment for a medical disorder, or as a form of **life support**, particularly in **intensive care units**. Inotropes that increase the force of contraction are "positive" inotropes, and include **sympathetic** agents such as **adrenaline**, **noradrenaline** and **dopamine**.<sup>[40]</sup> "Negative" inotropes decrease the force of contraction and include **calcium channel blockers**.<sup>[39]</sup>

### Electrical conduction

*Main articles: **Electrical conduction system of the heart** and **Heart rate***

The normal rhythmical heart beat, called **sinus rhythm**, is established by the **sinoatrial node**, the heart's **pacemaker**. Here an electrical signal is created that travels through the heart, causing the heart muscle to contract.

The sinoatrial node is found in the upper part of the **right atrium** near to the junction with the superior vena cava.<sup>[41]</sup> The electrical signal generated by the sinoatrial node travels through the right atrium in a radial way that is not completely understood. It travels to the left atrium via **Bachmann's bundle**, such that the muscles of the left and right atria contract together.<sup>[42][43][44]</sup> The signal then travels to the **atrioventricular node**. This is found at the bottom of the right atrium in the **atrioventricular septum**—the boundary between the right atrium and the left ventricle. The septum is part of the **cardiac skeleton**, tissue within the heart that the electrical signal cannot pass through, which forces the signal to pass through the atrioventricular



node only.<sup>[7]</sup> The signal then travels along the [bundle of His](#) to left and right [bundle branches](#) through to the ventricles of the heart. In the ventricles the signal is carried by specialized tissue called the [Purkinje fibers](#) which then transmit the electric charge to the heart muscle.<sup>[45]</sup>

## Heart rate

*Main article: [Heart rate](#)*

The normal resting heart rate is called the [sinus rhythm](#), created and sustained by the [sinoatrial node](#), a group of pacemaking cells found in the wall of the right atrium. Cells in the sinoatrial node do this by creating an [action potential](#). The [cardiac action potential](#) is created by the movement of specific [electrolytes](#) into and out of the pacemaker cells. The action potential then spreads to nearby cells.<sup>[46]</sup>

When the sinoatrial cells are resting, they have a negative charge on their membranes. However a rapid influx of [sodium](#) ions causes the membrane's charge to become positive. This is called [depolarisation](#) and occurs spontaneously.<sup>[7]</sup> Once the cell has a sufficiently high charge, the sodium channels close and [calcium](#) ions then begin to enter the cell, shortly after which [potassium](#) begins to leave it. All the ions travel through [ion channels](#) in the membrane of the sinoatrial cells. The potassium and calcium only start to move out of and into the cell once it has a sufficiently high charge, and so are called [voltage-gated](#). Shortly after this, the calcium channels close and [potassium channels](#) open, allowing potassium to leave the cell. This causes the cell to have a negative resting charge and is called [repolarization](#). When the membrane potential reaches approximately −60 mV, the potassium channels close and the process may begin again.<sup>[7]</sup>

The ions move from areas where they are concentrated to where they are not. For this reason sodium moves into the cell from outside, and potassium moves from within the cell to outside the cell. Calcium also plays a critical role. Their influx through slow channels means that the sinoatrial cells have a prolonged "plateau" phase when they have a positive charge. A part of this is called the [absolute refractory period](#). Calcium ions also combine with the regulatory protein [troponin C](#) in the [troponin complex](#) to enable [contraction](#) of the cardiac muscle, and separate from the protein to allow relaxation.<sup>[47]</sup>

The adult resting heart rate ranges from 60 to 100 bpm. The resting heart rate of a [newborn](#) can be 129 beats per minute (bpm) and this gradually decreases until maturity.<sup>[48]</sup> An athlete's heart rate can be lower than 60 bpm. During exercise the rate can be 150 bpm with maximum rates reaching from 200 to 220 bpm.<sup>[7]</sup>

## Influences

The normal [sinus rhythm](#) of the heart, giving the resting [heart rate](#), is influenced a number of factors. The cardiovascular centres in the brainstem that control the sympathetic and parasympathetic influences to the heart through the vagus nerve and sympathetic trunk.<sup>[49]</sup> These cardiovascular centres receive input from a series of receptors including [baroreceptors](#), sensing stretch the stretching of blood vessels and [chemoreceptors](#), sensing the amount of oxygen and carbon dioxide in the blood and its pH. Through a series of reflexes these help regulate and sustain blood flow.<sup>[7]</sup>

Baroreceptors are stretch receptors located in the [aortic sinus](#), [carotid bodies](#), the venae cavae, and other locations, including pulmonary vessels and the right side of the heart itself. Baroreceptors fire at a rate determined by how much they are stretched,<sup>[50]</sup> which is influenced by blood pressure, level of physical activity, and the relative distribution of blood. With increased pressure and stretch, the rate of baroreceptor firing increases, and the cardiac centers decrease sympathetic stimulation and increase parasympathetic stimulation. As pressure and stretch decrease, the rate of baroreceptor firing decreases, and the cardiac centers increase sympathetic stimulation and decrease parasympathetic stimulation.<sup>[7]</sup> There is a similar reflex, called the atrial reflex or [Bainbridge reflex](#), associated with varying rates of blood flow to the atria. Increased venous return stretches the walls of the atria where specialized baroreceptors are located. However, as the atrial baroreceptors increase their rate of firing and as they stretch due to the increased blood pressure, the cardiac center responds by increasing sympathetic stimulation and inhibiting parasympathetic stimulation to increase heart rate. The opposite is also true.<sup>[7]</sup> Chemoreceptors present in the carotid body or adjacent to the aorta in an aortic body respond to the blood's oxygen, carbon dioxide levels. Low oxygen or high carbon dioxide will stimulate firing of the receptors.<sup>[51]</sup>

Exercise and fitness levels, age, body temperature, [basal metabolic rate](#), and even a person's emotional state can all affect the heart rate. High levels of the hormones [epinephrine](#), norepinephrine, and [thyroid hormones](#) can increase the heart rate. The levels of electrolytes including calcium, potassium, and sodium can also influence the speed and regularity of the heart rate; [low blood oxygen](#), low [blood pressure](#) and [dehydration](#) may increase it.<sup>[7]</sup>

## Clinical significance

### Diseases

[Cardiovascular diseases](#), which include diseases of the heart, are the leading cause of death worldwide.<sup>[52]</sup> The majority of cardiovascular disease is noncommunicable and related to lifestyle and other factors, becoming more prevalent with ageing.<sup>[52]</sup> Heart disease is a major cause of death, accounting for an average of 30% of all deaths in 2008, globally.<sup>[11]</sup> This rate varies from a lower 28% to a high 40% in [high-income countries](#).<sup>[12]</sup> Doctors that specialise in the heart are called [cardiologists](#). Many other medical professionals are involved in treating diseases of the heart, including [doctors](#) such as [general practitioners](#), [cardiothoracic surgeons](#) and [intensivists](#), and [allied health](#) practitioners including [physiotherapists](#) and [dieticians](#).<sup>[53]</sup>

### Ischaemic heart disease

*Main article: [Coronary artery disease](#)*

[Coronary artery disease](#), also known as ischemic heart disease, is caused by [atherosclerosis](#) – a build-up of fatty material along the inner walls of the arteries. These fatty deposits known as atherosclerotic plaques [narrow](#) the coronary arteries, and if severe may reduce blood flow to the heart.<sup>[54]</sup> If a narrowing (or stenosis) is relatively minor then the patient may not experience any symptoms. Severe narrowings may

Transmission of a [cardiac action potential](#) through the heart's conduction system

Conduction system of the heart

The prepotential is due to a slow influx of sodium ions until the threshold is reached followed by a rapid depolarization and repolarization. The prepotential accounts for the membrane reaching threshold and initiates the spontaneous depolarization and contraction of the cell; there is no resting potential.<sup>[7]</sup>

The stethoscope is used for [auscultation](#) of the heart, and is one of the most iconic symbols for [medicine](#). A number of diseases can be detected primarily by listening for [heart murmurs](#).

cause chest pain ([angina](#)) or breathlessness during exercise or even at rest. The thin covering of an atherosclerotic plaque can rupture, exposing the fatty centre to the circulating blood. In this case a clot or thrombus can form, blocking the artery, and restricting blood flow to an area of heart muscle causing a myocardial infarction (a heart attack) or [unstable angina](#).<sup>[55]</sup> In the worst case this may cause cardiac arrest, a sudden and utter loss of output from the heart.<sup>[56]</sup> [Obesity](#), [high blood pressure](#), uncontrolled [diabetes](#), smoking and high [cholesterol](#) can all increase the risk of developing atherosclerosis and coronary artery disease.<sup>[52][54]</sup>

#### Heart failure

*Main article: [heart failure](#)*

[Heart failure](#) is defined as a condition in which the heart is unable to pump enough blood to meet the demands of the body.<sup>[57]</sup> Patients with heart failure may experience breathlessness especially when lying flat, as well as ankle swelling, known as [peripheral oedema](#). Heart failure is the end result of many diseases affecting the heart, but is most commonly associated with [ischaemic heart disease](#), [valvular heart disease](#), or [high blood pressure](#). Less common causes include various [cardiomyopathies](#). Heart failure is frequently associated with weakness of the heart muscle in the [ventricles](#) ([systolic](#) heart failure), but can also be seen in patients with heart muscle that is strong but stiff ([diastolic](#) heart failure). The condition may affect the [left ventricle](#) (causing predominantly breathlessness), the [right ventricle](#) (causing predominantly swelling of the legs and an elevated [jugular venous pressure](#)), or both ventricles. Patients with heart failure are at higher risk of developing dangerous heart rhythm disturbances or [arrhythmias](#).<sup>[57]</sup>

#### Cardiomyopathies

*Main article: [Cardiomyopathy](#)*

Cardiomyopathies are diseases affecting the muscle of the heart. Some cause abnormal thickening of the heart muscle ([hypertrophic cardiomyopathy](#)), some cause the heart to abnormally expand and weaken ([dilated cardiomyopathy](#)), some cause the heart muscle to become stiff and unable to fully relax between contractions ([restrictive cardiomyopathy](#)) and some make the heart prone to abnormal heart rhythms ([arrhythmogenic cardiomyopathy](#)). These conditions are often genetic and can be [inherited](#), but some such as dilated cardiomyopathy may be caused by damage from toxins such as alcohol. Some cardiomyopathies such as hypertrophic cardiomyopathy are linked to a higher risk of sudden cardiac death, particularly in athletes.<sup>[7]</sup> Many cardiomyopathies can lead to [heart failure](#) in the later stages of the disease.<sup>[57]</sup>

#### Valvular heart disease

*Main article: [Valvular heart disease](#)*

Healthy heart valves allow blood to flow easily in one direction, but prevent it from flowing in the other direction. Diseased heart valves may have a narrow opening and therefore restrict the flow of blood in the forward direction (referred to as a [stenotic valve](#)), or may allow blood to leak in the reverse direction (referred to as [valvular regurgitation](#)). Valvular heart disease may cause breathlessness, blackouts, or chest pain, but may be asymptomatic and only detected on a routine examination by hearing abnormal heart sounds or a [heart murmur](#). In the developed world, valvular heart disease is most commonly caused by degeneration secondary to old age, but may also be caused by infection of the heart valves ([endocarditis](#)). In some parts of the world [rheumatic heart disease](#) is a major cause of valvular heart disease, typically leading to mitral or aortic stenosis and caused by the body's immune system reacting to a [streptococcal](#) throat infection.<sup>[58][59]</sup>

#### Cardiac arrhythmias

*Main article: [Heart arrhythmia](#)*

While in the healthy heart, waves of electrical impulses originate in the [sinus node](#) before spreading to the rest of the atria, the [atrioventricular node](#), and finally the ventricles (referred to as a [normal sinus rhythm](#)), this normal rhythm can be disrupted. Abnormal heart rhythms or [arrhythmias](#) may be asymptomatic or may cause palpitations, blackouts, or breathlessness. Some types of arrhythmia such as [atrial fibrillation](#) increase the long term risk of [stroke](#).<sup>[60]</sup>

Some arrhythmias cause the heart to beat abnormally slowly, referred to as a [bradycardia](#) or bradyarrhythmia. This may be caused by an [abnormally slow sinus node](#) or damage within the cardiac conduction system ([heart block](#)).<sup>[61]</sup> In other arrhythmias the heart may beat abnormally rapidly, referred to as a [tachycardia](#) or tachyarrhythmia. These arrhythmias can take many forms and can originate from different structures within the heart – some arise from the atria (e.g. [atrial flutter](#)), some from the atrioventricular node (e.g. [AV nodal re-entrant tachycardia](#)) whilst others arise from the ventricles (e.g. [ventricular tachycardia](#)). Some tachyarrhythmias are caused by scarring within the heart (e.g. some forms of [ventricular tachycardia](#)), others by an irritable focus (e.g. focal [atrial tachycardia](#)), while others are caused by additional abnormal conduction tissue that has been present since birth (e.g. [Wolff-Parkinson-White syndrome](#)). The most dangerous form of heart racing is [ventricular fibrillation](#), in which the ventricles quiver rather than contract, and which if untreated is rapidly fatal.<sup>[62]</sup>

#### Pericardial disease

The sack which surrounds the heart, called the pericardium, can become inflamed in a condition known as [pericarditis](#). This condition typically causes chest pain that may spread to the back, and is often caused by a viral infection ([glandular fever](#), [cytomegalovirus](#), or [coxsackievirus](#)). Fluid can build up within the pericardial sack, referred to as a [pericardial effusion](#). Pericardial effusions often occur secondary to pericarditis, kidney failure, or tumours, and frequently do not cause any symptoms. However, large effusions or effusions which accumulate rapidly can compress the heart in a condition known as [cardiac tamponade](#), causing breathlessness and potentially fatal low blood pressure. Fluid can be removed from the pericardial space for diagnosis or to relieve tamponade using a syringe in a procedure called [pericardiocentesis](#).<sup>[63]</sup>

#### Congenital heart disease

*Main article: [Congenital heart defect](#)*

Some people are born with hearts that are abnormal and these abnormalities are known as [congenital heart defects](#). They may range from the relatively minor (e.g. [patent foramen ovale](#), arguably a variant of normal) to serious life-threatening abnormalities (e.g. [hypoplastic left heart syndrome](#)). Common abnormalities include those that affect the heart muscle that separates the two sides of the heart (a 'hole in the heart' e.g. [ventricular septal defect](#)). Other defects include those affecting the heart valves (e.g. [congenital aortic stenosis](#)), or the main blood vessels that lead from the heart (e.g. [coarctation of the aorta](#)). More complex syndromes are seen that affect more than one part of the heart (e.g. [Tetralogy of Fallot](#)).

Some congenital heart defects allow blood that is low in oxygen that would normally be returned to the lungs to instead be pumped back to the rest of the body. These are known as [cyanotic congenital heart defects](#) and are often more serious. Major congenital heart defects are often picked up in childhood, shortly after birth, or even before a child is born (e.g. [transposition of the great arteries](#)), causing breathlessness and a lower rate of growth. More minor forms of congenital heart disease may remain undetected for many years and only reveal themselves in adult life (e.g. [atrial septal defect](#)).<sup>[64][65]</sup>

#### Diagnosis

Atherosclerosis is a condition affecting the circulatory system. If the coronary arteries are affected, [angina pectoris](#) may result or at worse a [heart attack](#).

Heart disease is diagnosed by the taking of a [medical history](#), a [cardiac examination](#), and further investigations, including [blood tests](#), [echocardiograms](#), [ECGs](#) and [imaging](#). Other invasive procedures such as [cardiac catheterisation](#) can also play a role. <sup>[66]</sup>

## Examination

*Main articles: [Cardiac examination](#) and [Heart sounds](#)*

The cardiac examination includes inspection, feeling the chest with the hands ([palpation](#)) and listening with a stethoscope ([auscultation](#)).<sup>[67][68]</sup> It involves assessment of [signs](#) that may be visible on a person's hands (such as [splinter haemorrhages](#)), joints and other areas. A person's pulse is taken, usually at the [radial artery](#) near the wrist, in order to assess for the rhythm and strength of the pulse. The [blood pressure](#) is taken, using either a manual or automatic [sphygmomanometer](#) or using a [more invasive measurement](#) from within the artery. Any elevation of the [jugular venous pulse](#) is noted. A person's [chest](#) is felt for any transmitted vibrations from the heart, and then listened to with a stethoscope.

## Heart sounds

Typically, healthy hearts have only two audible [heart sounds](#), called S1 and S2. The first heart sound S1, is the sound created by the closing of the atrioventricular valves during ventricular contraction and is normally described as "lub". The second heart sound, S2, is the sound of the semilunar valves closing during ventricular diastole and is described as "dub".<sup>[7]</sup> Each sound consists of two components, reflecting the slight difference in time as the two valves close.<sup>[69]</sup> S2 may [split](#) into two distinct sounds, either as a result of inspiration or different valvular or cardiac problems.<sup>[69]</sup> Additional heart sounds may also be present and these give rise to [gallop rhythms](#). A [third heart sound](#), S3 usually indicates an increase in ventricular blood volume. A [fourth heart sound](#) S4 is referred to as an atrial gallop and is produced by the sound of blood being forced into a stiff ventricle. The combined presence of S3 and S4 give a quadruple gallop.<sup>[7]</sup>

[Heart murmurs](#) are abnormal heart sounds which can be either related to disease or benign, and there are several kinds.<sup>[70]</sup> There are normally two heart sounds, and abnormal heart sounds can either be extra sounds, or "murmurs" related to the flow of blood between the sounds. Murmurs are graded by volume, from 1 (the quietest), to 6 (the loudest), and evaluated by their relationship to the heart sounds, position in the cardiac cycle, and additional features such as their radiation to other sites, changes with a person's position, the frequency of the sound as determined by the side of the [stethoscope](#) by which they are heard, and site at which they are heard loudest.<sup>[70]</sup> Murmurs may be caused by [damaged heart valves](#), congenital heart disease such as [ventricular septal defects](#), or may be heard in normal hearts. A different type of sound, a [pericardial friction rub](#) can be heard in cases of pericarditis where the inflamed membranes can rub together.

## Blood tests

Blood tests play an important role in the diagnosis and treatment of many cardiovascular conditions.

[Troponin](#) is a sensitive [biomarker](#) for a heart with insufficient blood supply. It is released 4–6 hours after injury, and usually peaks at about 12–24 hours.<sup>[40]</sup> Two tests of troponin are often taken – one at the time of initial presentation, and another within 3–6 hours, <sup>[71]</sup> with either a high level or a significant rise being diagnostic. A test for [brain natriuretic peptide](#) (BNP) can be used to evaluate for the presence of heart failure, and rises when there is increased demand on the left ventricle. These tests are considered [biomarkers](#) because they are highly specific for cardiac disease.<sup>[72]</sup> Testing for the [MB form of creatine kinase](#) provides information about the heart's blood supply, but is used less frequently because it is less specific and sensitive.<sup>[73]</sup>

Other blood tests are often taken to help understand a person's general health and risk factors that may contribute to heart disease. These often include a [full blood count](#) investigating for [anaemia](#), and [basic metabolic panel](#) that may reveal any disturbances in electrolytes. A [coagulation screen](#) is often required to ensure that the right level of anticoagulation is given. [Fasting lipids](#) and [fasting blood glucose](#) (or an [HbA1c](#) level) are often ordered to evaluate a person's [cholesterol](#) and [diabetes](#) status, respectively. <sup>[74]</sup>

## Electrocardiogram

*Main article: [Electrocardiography](#)*

Using surface electrodes on the body, it is possible to record the electrical activity of the heart. This tracing of the electrical signal is the electrocardiogram (ECG) or (EKG). An ECG is a [bedside test](#) and involves the placement of ten leads on the body. This produces a "12 lead" ECG (three extra leads are calculated mathematically, and one lead is a [ground](#)).<sup>[75]</sup>

There are five prominent features on the ECG: the P wave (atrial depolarisation), the QRS complex (ventricular depolarisation<sup>[h]</sup>) and the T wave (ventricular repolarisation).<sup>[7]</sup> As the heart cells contract, they create a current that travels through the heart. A downward deflection on the ECG implies cells are becoming more positive in charge ("depolarising") in the direction of that lead, whereas an upward inflection implies cells are becoming more negative ("repolarising") in the direction of the lead. This depends on the position of the lead, so if a wave of depolarising moved from left to right, a lead on the left would show a negative deflection, and a lead on the right would show a positive deflection. The ECG is a useful tool in detecting rhythm disturbances and in detecting insufficient blood supply to the heart.<sup>[75]</sup> Sometimes abnormalities are suspected, but not immediately visible on the ECG. [Testing when exercising](#) can be used to provoke an abnormality, or an ECG can be worn for a longer period such as a 24-hour [Holter monitor](#) if a suspected rhythm abnormality is not present at the time of assessment.<sup>[75]</sup>

## Imaging

*Main article: [Cardiac imaging](#)*

Several [imaging methods](#) can be used to assess the anatomy and function of the heart, including [ultrasound \(echocardiography\)](#), [angiography](#), [CT scans](#), [MRI](#) and [PET](#). An echocardiogram is an ultrasound of the heart used to measure the heart's function, assess for valve disease, and look for any abnormalities. Echocardiography can be conducted by a probe on the chest ("transthoracic") or by a probe in the [esophagus](#) ("transoesophageal"). A typical echocardiography report will include information about the width of the valves noting any [stenosis](#), whether there is any backflow of blood ([regurgitation](#)) and information about the blood volumes at the end of systole and diastole, including an [ejection fraction](#), which describes how much blood is ejected from the left and right ventricles after systole. Ejection fraction can then be obtained by dividing the volume ejected by the heart (stroke volume) by the volume of the filled heart (end-diastolic volume).<sup>[76]</sup> Echocardiograms can also be conducted under circumstances when the body is more stressed, in order to examine for signs of lack of blood supply. This [cardiac stress test](#) involves either direct exercise, or where this is not possible, injection of a drug such as [dobutamine](#).<sup>[68]</sup>

CT scans, [chest X-rays](#) and other forms of [imaging](#) can help evaluate the heart's size, evaluate for signs of [pulmonary oedema](#), and indicate whether there is [fluid around the heart](#). They are also useful for evaluating the aorta, the major blood vessel which leaves the heart. <sup>[68]</sup>

## Treatment

3D echocardiogram showing the mitral valve (right), tricuspid and mitral valves (top left) and aortic valve (top right). The closure of the heart valves causes the [heart sounds](#).

Cardiac cycle shown against ECG



Diseases affecting the heart can be treated by a variety of methods including lifestyle modification, drug treatment, and surgery.

### Ischaemic heart disease

*Main articles: [Coronary artery disease](#), [Coronary artery bypass surgery](#), and [Coronary stent](#)*

Narrowings of the coronary arteries ([ischaemic heart disease](#)) are treated to relieve symptoms of chest pain caused by a partially narrowed artery ([angina pectoris](#)), to minimise heart muscle damage when an artery is completely occluded ([myocardial infarction](#)), or to prevent a myocardial infarction from occurring. Medications to improve angina symptoms include [nitroglycerin](#), [beta blockers](#), and calcium channel blockers, while preventative treatments include [antiplatelets](#) such as [aspirin](#) and [statins](#), lifestyle measures such as stopping smoking and weight loss, and treatment of risk factors such as [high blood pressure](#) and [diabetes](#).<sup>[77]</sup>

In addition to using medications, narrowed heart arteries can be treated by expanding the narrowings or redirecting the flow of blood to bypass an obstruction. This may be performed using a [percutaneous coronary intervention](#), during which narrowings can be expanded by passing small balloon-tipped wires into the coronary arteries, inflating the balloon to expand the narrowing, and sometimes leaving behind a metal scaffold known as a stent to keep the artery open.<sup>[78]</sup>

If the narrowings in coronary arteries are unsuitable for treatment with a percutaneous coronary intervention, open surgery may be required. A [coronary artery bypass graft](#) can be performed, whereby a blood vessel from another part of the body (the [saphenous vein](#), [radial artery](#), or [internal mammary artery](#)) is used to redirect blood from a point before the narrowing (typically the [aorta](#)) to a point beyond the obstruction.<sup>[78][79]</sup>

### Valvular heart disease

*Main article: [Artificial heart valve](#)*

Diseased [heart valves](#) that have become abnormally narrow or abnormally leaky may require surgery. This is traditionally performed as an open surgical procedure to replace the damaged heart valve with a tissue or metallic [prosthetic valve](#). In some circumstances, the [tricuspid](#) or [mitral](#) valves can be repaired [surgically](#), avoiding the need for a valve replacement. Heart valves can also be treated percutaneously, using techniques that share many similarities with percutaneous coronary intervention. [Transcatheter aortic valve replacement](#) is increasingly used for patients consider very high risk for open valve replacement.<sup>[58]</sup>

### Cardiac arrhythmias

*Main articles: [Heart arrhythmia](#), [Radiofrequency ablation](#), and [Artificial cardiac pacemaker](#)*

Abnormal heart rhythms ([arrhythmias](#)) can be treated using antiarrhythmic drugs. These may work by manipulating the flow of electrolytes across the cell membrane (such as [calcium channel blockers](#), [sodium channel blockers](#), [amiodarone](#), or [digoxin](#)), or modify the autonomic nervous system's effect on the heart ([beta blockers](#) and [atropine](#)). In some arrhythmias such as atrial fibrillation which increase the risk of stroke, this risk can be reduced using anticoagulants such as [warfarin](#) or [novel oral anticoagulants](#).<sup>[60]</sup>

If medications fail to control an arrhythmia, another treatment option may be [catheter ablation](#). In these procedures, wires are passed from a [vein](#) or [artery](#) in the leg to the heart to find the abnormal area of tissue that is causing the arrhythmia. The abnormal tissue can be intentionally damaged, or ablated, by [heating](#) or [freezing](#) to prevent further heart rhythm disturbances. Whilst the majority of arrhythmias can be treated using minimally invasive catheter techniques, some arrhythmias (particularly [atrial fibrillation](#)) can also be treated using open or [thoracoscopic](#) surgery, either at the time of other cardiac surgery or as a standalone procedure. A [cardioversion](#), whereby an electric shock is used to stun the heart out of an abnormal rhythm, may also be used.

Cardiac devices in the form of [pacemakers](#) or [implantable defibrillators](#) may also be required to treat arrhythmias. Pacemakers, comprising a small battery powered generator implanted under the skin and one or more leads that extend to the heart, are most commonly used to treat abnormally [slow heart rhythms](#).<sup>[61]</sup> Implantable defibrillators are used to treat serious life-threatening rapid heart rhythms. These devices monitor the heart, and if dangerous heart racing is detected can automatically deliver a shock to restore the heart to a normal rhythm. Implantable defibrillators are most commonly used in patients with [heart failure](#), [cardiomyopathies](#), or inherited arrhythmia syndromes.

### Heart failure

*Main article: [Heart failure](#)*

As well as addressing the underlying cause for a patient's heart failure (most commonly [ischaemic heart disease](#) or [hypertension](#)), the mainstay of heart failure treatment is with medication. These include drugs to prevent fluid from accumulating in the lungs by increasing the amount of urine a patient produces ([diuretics](#)), and drugs that attempt to preserve the pumping function of the heart ([beta blockers](#), [ACE inhibitors](#) and [mineralocorticoid receptor antagonists](#)).<sup>[57]</sup>

In some patients with heart failure, a specialised [pacemaker](#) known as [cardiac resynchronisation therapy](#) can be used to improve the heart's pumping efficiency.<sup>[61]</sup> These devices are frequently combined with a defibrillator. In very severe cases of heart failure, a small pump called a [ventricular assist device](#) may be implanted which supplements the heart's own pumping ability. In the most severe cases, a [cardiac transplant](#) may be considered.<sup>[57]</sup>

## History

### Ancient

Humans have known about the heart since ancient times, although its precise function and anatomy were not clearly understood.<sup>[80]</sup> From the primarily religious views of earlier societies towards the heart, [ancient Greeks](#) are considered to have been the primary seat of scientific understanding of the heart in the ancient world.<sup>[81][82][83]</sup> [Aristotle](#) considered the heart to be organ responsible for creating blood; [Plato](#) considered the heart as the source of circulating blood and [Hippocrates](#) noted blood circulating cyclically from the body through the heart to the lungs.<sup>[81][83]</sup> [Erasistratos](#) (304–250 BCE) noted the heart as a pump, causing dilation of blood vessels, and noted that arteries and veins both radiate from the heart, becoming progressively smaller with distance, although he believed they were filled with air and not blood. He also discovered the heart valves.<sup>[81]</sup>

The Greek physician [Galen](#) (2nd century CE) knew blood vessels carried blood and identified venous (dark red) and arterial (brighter and thinner) blood, each with distinct and separate functions.<sup>[81]</sup> Galen, noting the heart as the hottest organ in the body, concluded that it provided heat to the body.<sup>[83]</sup> The heart did not pump blood around, the heart's motion sucked blood in during diastole and the blood moved by the pulsation of the arteries themselves.<sup>[83]</sup> Galen believed the arterial blood was created by venous blood passing from the left ventricle to the right through 'pores' between the ventricles.<sup>[80]</sup> Air from the lungs passed from the lungs via the pulmonary artery to the left side of the heart and created arterial blood.<sup>[83]</sup>

These ideas went unchallenged for almost a thousand years.<sup>[80][83]</sup>

Heart and its blood vessels, by [Leonardo da Vinci](#), 15th century

Pre-modern

The earliest descriptions of the [coronary](#) and [pulmonary circulation](#) systems can be found in the *Commentary on Anatomy in Avicenna's Canon* , published in 1242 by [Ibn al-Nafis](#).<sup>[84]</sup> In his manuscript, al-Nafis wrote that blood passes through the pulmonary circulation instead of moving from the right to the left ventricle as previously believed by Galen.<sup>[85]</sup> His work was later translated into [Latin](#) by [Andrea Alpago](#).<sup>[86]</sup>

In Europe, the teachings of Galen continued to dominate the academic community and his doctrines were adopted as the official canon of the Church. [Andreas Vesalius](#) questioned some of Galen's beliefs of the heart in *De humani corporis fabrica* (1543), but his *magnum opus* was interpreted as a challenge to the authorities and he was subjected to a number of attacks.<sup>[87]</sup> [Michael Servetus](#) wrote in *Christianismi Restitutio* (1553) that blood flows from one side of the heart to the other via the lungs.<sup>[87]</sup>

Modern

A breakthrough in understanding the flow of blood through the heart and body came with the publication of *De Motu Cordis* (1628) by the English physician [William Harvey](#). Harvey's book completely describes the [systemic circulation](#) and the mechanical force of the heart, leading to an overhaul of the Galenic doctrines. <sup>[83]</sup> [Otto Frank](#) (1865–1944) was a German physiologist; among his many published works are detailed studies of this important heart relationship. [Ernest Starling](#) (1866–1927) was an important English physiologist who also studied the heart. Although they worked largely independently, their combined efforts and similar conclusions have been recognized in the name "[Frank–Starling mechanism](#)".<sup>[7]</sup>

Although [Purkinje fibers](#) and the [bundle of His](#) were discovered as early as the 19th century, their specific role in the [electrical conduction system of the heart](#) remained unknown until [Sunao Tawara](#) published his monograph, titled *Das Reizleitungssystem des Säugetierherzens*, in 1906. Tawara's discovery of the [atrioventricular node](#) prompted [Arthur Keith](#) and [Martin Flack](#) to look for similar structures in the heart, leading to their discovery of the [sinoatrial node](#) several months later. These structures form the anatomical basis of the [electrocardiogram](#), whose inventor, [Willem Einthoven](#), was awarded the Nobel Prize in Medicine or Physiology in 1924.<sup>[88]</sup>

The first successful [heart transplantation](#) was performed in 1967 by the South African surgeon [Christiaan Barnard](#) at [Groote Schuur Hospital](#) in [Cape Town](#). This marked an important milestone in [cardiac surgery](#), capturing the attention of both the medical profession and the world at large. However, long-term survival rates of patients were initially very low. [Louis Washkansky](#), the first recipient of a donated heart, died 18 days after the operation while other patients did not survive for more than a few weeks.<sup>[89]</sup> The American surgeon [Norman Shumway](#) has been credited for his efforts to improve transplantation techniques, along with pioneers [Richard Lower](#), [Vladimir Demikhov](#) and [Adrian Kantrowitz](#). As of March 2000, more than 55,000 heart transplantations have been performed worldwide.<sup>[90]</sup>

By the middle of the 20th century, [heart disease](#) had surpassed infectious disease as the leading cause of death in the United States, and it is currently the leading cause of deaths worldwide. Since 1948, the ongoing [Framingham Heart Study](#) has shed light on the effects of various influences on the heart, including diet, exercise, and common medications such as [aspirin](#). Although the introduction of [ACE inhibitors](#) and [beta blockers](#) has improved the management of chronic [heart failure](#), the disease continues to be an enormous medical and societal burden, with 30 to 40% of patients dying within a year of receiving the diagnosis.<sup>[91]</sup>

Society and culture

Further information: *[Sacred Heart](#), [Heart symbol](#), and [Blood § Cultural and religious beliefs](#)*

Symbolism

As one of the vital organs, the heart was long identified as the center of the entire body, the seat of life, or emotion, or reason, will, intellect, purpose or the mind.<sup>[92]</sup> The heart is an emblematic symbol in many religions, signifying "truth, conscience or moral courage in many religions – the temple or throne of God in Islamic and [Judeo-Christian](#) thought; the divine centre, or [atman](#), and the [third eye](#) of transcendent wisdom in [Hinduism](#); the diamond of purity and essence of the [Buddha](#); the [Taoist](#) centre of understanding."<sup>[92]</sup>

In the [Hebrew Bible](#), the word for heart, *lev*, is used in these meanings, as the seat of emotion, the mind, and referring to the anatomical organ. It is also connected in function and symbolism to the stomach.<sup>[93]</sup>



An important part of the concept of the [soul](#) in [Ancient Egyptian religion](#) was thought to be the heart, or *ib*. The *ib* or metaphysical heart was believed to be formed from one drop of blood from the child's mother's heart, taken at conception.<sup>[94]</sup> To ancient Egyptians, the heart was the seat of [emotion](#), [thought](#), will, and [intention](#). This is evidenced by [Egyptian](#) expressions which incorporate the word *ib*, such as *Awī-ib* for "happy" (literally, "long of heart"), *Xak-ib* for "estranged" (literally, "truncated of heart").<sup>[95]</sup> In Egyptian religion, the heart was the key to the afterlife. It was conceived as surviving death in the nether world, where it gave evidence for, or against, its possessor. It was thought that the heart was examined by [Anubis](#) and a variety of [deities](#) during the *Weighing of the Heart* ceremony. If the heart weighed more than the feather of [Maat](#), which symbolized the ideal standard of behavior. If the scales balanced, it meant the heart's possessor had lived a just life and could enter the afterlife; if the heart was heavier, it would be devoured by the monster [Ammit](#).<sup>[96]</sup>

The [Chinese](#) character for "heart", 心, derives from a comparatively realistic depiction of a heart (indicating the heart chambers) in [seal script](#).<sup>[97]</sup> The Chinese word *xīn* also takes the metaphorical meanings of "mind", "intention", or "core".<sup>[98]</sup> In [Chinese medicine](#), the heart is seen as the center of [神 \*shén\*](#) "spirit, consciousness".<sup>[99]</sup> The heart is associated with the [small intestine](#), [tongue](#), governs the [six organs and five viscera](#), and belongs to fire in the five elements.<sup>[100]</sup>

The Sanskrit word for heart is *hr̥d* or *hr̥daya*, found in the oldest surviving Sanskrit text, the [Rigveda](#). In Sanskrit, it may mean both the anatomical object and "mind" or "soul", representing the seat of emotion. *Hrd* may be a cognate of the word for heart in Greek, Latin, and English.<sup>[101][102]</sup>

Many [classical](#) philosophers and scientists, including [Aristotle](#), considered the heart the seat of thought, [reason](#), or emotion, often disregarding the brain as contributing to those functions.<sup>[103]</sup> The identification of the heart as the seat of [emotions](#) in particular is due to the [Roman](#) physician [Galen](#), who also located the seat of the passions in the [liver](#), and the seat of reason in the brain.<sup>[104]</sup>

The heart also played a role in the [Aztec](#) system of belief. The most common form of human sacrifice practiced by the Aztecs was heart-extraction. The Aztec believed that the heart (*tona*) was both the seat of the individual and a fragment of the Sun's heat ( *istli*). To this day, the Nahua consider



the Sun to be a heart-soul (*tona-tiuh*): "round, hot, pulsating".<sup>[105]</sup>

In **Catholicism**, there has been a long tradition of veneration of the heart, stemming from worship of the wounds of **Jesus Christ** which gained prominence from the mid sixteenth century.<sup>[106]</sup> This tradition influenced the development of the medieval Christian **devotion** to the **Sacred Heart of Jesus** and the parallel worship of **Immaculate Heart of Mary**, made popular by **John Eudes**.<sup>[107]</sup>

The expression of a **broken heart** is a cross-cultural reference to **grief** for a lost one or to unfulfilled **romantic love**.

The notion of "**Cupid's arrows**" is ancient, due to **Ovid**, but while Ovid describes Cupid as wounding his victims with his arrows, it is not made explicit that it is the *heart* that is wounded. The familiar iconography of Cupid shooting little **heart symbols** is a **Renaissance** theme that became tied to **Valentine's day**.<sup>[92]</sup>

## Food

Animal hearts are widely consumed as food. As they are almost entirely muscle, they are high in protein. They are often included in dishes with other offal, for example in the **pan-Ottoman kokoretsi**.

Chicken hearts are considered to be **giblets**, and are often grilled on skewers: **Japanese hāto yakitori**, **Brazilian churrasco de coração**, **Indonesian chicken heart satay**.<sup>[108]</sup> They can also be pan-fried, as in **Jerusalem mixed grill**. In **Egyptian cuisine**, they can be used, finely chopped, as part of **stuffing** for chicken.<sup>[109]</sup> Many recipes combined them with other giblets, such as the **Mexican pollo en menudencias**<sup>[110]</sup> and the **Russian ragu iz kurinyikh potrokhov**.<sup>[111]</sup>

The hearts of beef, pork, and mutton can generally be interchanged in recipes. As heart is a hard-working muscle, it makes for "firm and rather dry" meat,<sup>[112]</sup> so is generally slow-cooked. Another way of dealing with toughness is to **julienne** the meat, as in **Chinese** stir-fried heart.<sup>[113]</sup>

**Beef** heart may be grilled or braised.<sup>[114]</sup> In the **Peruvian anticuchos de corazón**, barbecued beef hearts are grilled after being tenderized through long **marination** in a spice and vinegar mixture. An **Australian** recipe for "mock goose" is actually braised stuffed beef heart.<sup>[115]</sup>

**Pig** heart is stewed, poached, braised,<sup>[116]</sup> or made into sausage. The **Balinese oret** is a sort of **blood sausage** made with pig heart and blood. A **French** recipe for *cœur de porc à l'orange* is made of braised heart with an orange sauce.

## Other animals

*See also: **Circulatory system***

### Other vertebrates

The size of the heart varies among the different **animal groups**, with hearts in **vertebrates** ranging from those of the smallest mice (12 mg) to the blue whale (600 kg).<sup>[117]</sup> In **vertebrates**, the heart lies in the middle of the ventral part of the body, surrounded by a **pericardium**.<sup>[118]</sup> which in some fish may be connected to the **peritoneum**.<sup>[119]</sup>

The SA node is found in all **amniotes** but not in more primitive vertebrates. In these animals, the muscles of the heart are relatively continuous and the sinus venosus coordinates the beat which passes in a wave through the remaining chambers. Indeed, since the sinus venosus is incorporated into the right atrium in amniotes, it is likely **homologous** with the SA node. In teleosts, with their vestigial sinus venosus, the main centre of coordination is, instead, in the atrium. The rate of heartbeat varies enormously between different species, ranging from around 20 beats per minute in **codfish** to around 600 in **hummingbirds**<sup>[120]</sup> and up to 1200 bpm in the **ruby-throated hummingbird**.<sup>[121]</sup>

### Double circulatory systems

Adult **amphibians** and most **reptiles** have a **double circulatory system**, meaning a circulatory system divided into arterial and venous parts. However, the heart itself is not completely separated into two sides. Instead, it is separated into three chambers – two atria and one ventricle. Blood returning from both the systemic circulation and the lungs is returned, and blood is pumped simultaneously into the systemic circulation and the lungs. The double system allows blood to circulate to and from **lungs** which deliver oxygenated blood directly to the heart.<sup>[122]</sup>

In reptiles, the heart is usually situated around the middle of the thorax, and in snakes, usually between the junction of the upper first and second third. There is a heart with three chambers: two atria and one ventricle. The ventricle is incompletely separated into two halves by a wall (**septum**), with a considerable gap near the pulmonary artery and aortic openings. In most reptilian species, there appears to be little, if any, mixing between the bloodstreams, so the aorta receives, essentially, only oxygenated blood.<sup>[120][122]</sup> The exception to this rule is **crocodiles**, which have a four-chambered heart.<sup>[123]</sup>

In the heart of **lungfish**, the septum extends part-way into the ventricle. This allows for some degree of separation between the de-oxygenated bloodstream destined for the lungs and the oxygenated stream that is delivered to the rest of the body. The absence of such a division in living amphibian species may be partly due to the amount of respiration that occurs through the skin; thus, the blood returned to the heart through the venae cavae is already partially oxygenated. As a result, there may be less need for a finer division between the two bloodstreams than in lungfish or other **tetrapods**. Nonetheless, in at least some species of amphibian, the spongy nature of the ventricle does seem to maintain more of a separation between the bloodstreams. Also, the original valves of the **conus arteriosus** have been replaced by a spiral valve that divides it into two parallel parts, thereby helping to keep the two bloodstreams separate.<sup>[120]</sup>

### The fully divided heart

**Archosaurs** (**crocodilians** and **birds**) and **mammals** show complete separation of the heart into two pumps for a total of four **heart chambers**; it is thought that the four-chambered heart of archosaurs evolved independently from that of mammals. In crocodilians, there is a small opening, the **foramen of Panizza**, at the base of the arterial trunks and there is some degree of mixing between the blood in each side of the heart, during a dive underwater;<sup>[124][125]</sup> thus, only in birds and mammals are the two streams of blood – those to the pulmonary and systemic circulations – permanently kept entirely separate by a physical barrier.<sup>[120]</sup>

### Fish

*Main article: **Fish anatomy § Heart***

Fish have what is often described as a two-chambered heart,<sup>[126]</sup> consisting of one atrium to receive blood and one ventricle to pump it,<sup>[127]</sup> However, the fish heart has entry and exit compartments that may be called chambers, so it is also sometimes described as three-chambered<sup>[127]</sup> or four-chambered,<sup>[128]</sup> depending on what is counted as a chamber. The atrium and ventricle are sometimes considered "true chambers", while the others are considered "accessory chambers".<sup>[129]</sup>


A cross section of a three chambered adult amphibian heart, note the single ventricle. The purple regions represent areas where mixing of oxygenated and de-oxygenated blood occurs.

1. Pulmonary vein
2. Left atrium
3. Right atrium
4. Ventricle
5. Conus arteriosus
6. Sinus venosus

Primitive fish have a four-chambered heart, but the chambers are arranged sequentially so that this primitive heart is quite unlike the four-chambered hearts of mammals and birds. The first chamber is the [sinus venosus](#), which collects deoxygenated blood, from the body, through the [hepatic](#) and [cardinal veins](#). From here, blood flows into the [atrium](#) and then to the powerful muscular [ventricle](#) where the main pumping action will take place. The fourth and final chamber is the [conus arteriosus](#) which contains several valves and sends blood to the [ventral aorta](#). The ventral aorta delivers blood to the gills where it is oxygenated and flows, through the [dorsal aorta](#), into the rest of the body. (In [tetrapods](#), the ventral aorta has divided in two; one half forms the [ascending aorta](#), while the other forms the [pulmonary artery](#)).<sup>[120]</sup>

In the adult fish, the four chambers are not arranged in a straight row but, instead form an S-shape with the latter two chambers lying above the former two. This relatively simpler pattern is found in [cartilaginous fish](#) and in the [ray-finned fish](#). In [teleosts](#), the conus arteriosus is very small and can more accurately be described as part of the aorta rather than of the heart proper. The conus arteriosus is not present in any [amniotes](#), presumably having been absorbed into the ventricles over the course of evolution. Similarly, while the sinus venosus is present as a vestigial structure in some reptiles and birds, it is otherwise absorbed into the right atrium and is no longer distinguishable.<sup>[120]</sup>

Invertebrates



Basic arthropod body structure – heart shown in red

[Arthropods](#) and most [mollusks](#) have an open circulatory system. In this system, deoxygenated blood collects around the heart in cavities ([sinuses](#)). This blood slowly permeates the heart through many small one-way channels. The heart then pumps the blood into the [hemocoel](#), a cavity between the organs. The heart in arthropods is typically a muscular tube that runs the length of the body, under the back and from the base of the head. Instead of blood the circulatory fluid is [haemolymph](#) which carries the most commonly used [respiratory pigment](#), copper-based [haemocyanin](#) as the oxygen transporter; iron-based haemoglobin is used by only a few arthropods.<sup>[130]</sup>

In some other invertebrates such as [earthworms](#), the circulatory system is not used to transport oxygen and so is much reduced, having no veins or arteries and consisting of two connected tubes. Oxygen travels by diffusion and there are five small muscular vessels that connect these vessels that contract at the front of the animals that can be thought of as "hearts".<sup>[130]</sup>

[Squids and other cephalopods](#) have two "gill hearts" also known as [branchial hearts](#), and one "systemic heart". The brachial hearts have two atria and one ventricle each, and pump to the [gills](#), whereas the systemic heart pumps to the body.<sup>[131][132]</sup>

The tube-like heart (green) of the mosquito [Anopheles gambiae](#) extends horizontally across the body, interlinked with the diamond-shaped [wing muscles](#) (also green) and surrounded by [pericardial cells](#) (red). Blue depicts [cell nuclei](#).

Additional images



The human heart viewed from the front      The human heart viewed from behind      The [coronary circulation](#)      The human heart viewed from the front and from behind      Frontal section of the human heart



An anatomical specimen of the heart      Heart illustration with circulatory system

Notes

- a. ^ From the heart to the body
- b. ^ Arteries that contain deoxygenated blood, from the heart to the lungs
- c. ^ Supplying blood to the heart itself
- d. ^ From the body to the heart
- e. ^ Veins containing oxygenated blood from the lungs to the heart
- f. ^ Veins that drain blood from the cardiac tissue itself
- g. ^ Note the muscles do **not** cause the valves to open. The pressure difference between the blood in the atria and the ventricles does this.
- h. ^ Depolarisation of the ventricles occurs concurrently, but is not significant enough to be detected on an ECG. <sup>[75]</sup>

## References

This article incorporates text from the **CC-BY** book: *OpenStax College, Anatomy & Physiology. OpenStax CNX. 30 jul 2014* ..

- <sup>1</sup> <sup>^</sup> Taber, Clarence Wilbur; Venes, Donald (2009). *Taber's cyclopedic medical dictionary* . F. A. Davis Co. pp. 1018–23. ISBN 0-8036-1559-0
- <sup>2</sup> <sup>^</sup> <sup>a</sup> <sup>b</sup> <sup>c</sup> Guyton & Hall 2011, p. 157.
- <sup>3</sup> <sup>^</sup> <sup>a</sup> <sup>b</sup> <sup>c</sup> Moore, Keith L.; Dalley, Arthur F.; Agur, Anne M. R. "1". *Clinically Oriented Anatomy* . Wolters Kluwer Health/Lippincott Williams & Wilkins. pp. 127–73. ISBN 978-1-60547-652-0.
- <sup>4</sup> <sup>^</sup> Starr, Cecie; Evers, Christine; Starr, Lisa (2 January 2009). *Biology: Today and Tomorrow With Physiology* . Cengage Learning. p. 422. ISBN 978-0-495-56157-6 Archived from the original on 2 May 2016.
- <sup>5</sup> <sup>^</sup> <sup>a</sup> <sup>b</sup> <sup>c</sup> Reed, C. Roebuck; Brainerd, Lee Wherry; Lee., Rodney; Inc, the staff of Kaplan, (2008). *CSET : California Subject Examinations for Teachers* (3rd ed.). New York, NY: Kaplan Pub. p. 154. ISBN 978-1-4195-5281-6 Archived from the original on 4 May 2016.
- <sup>6</sup> <sup>^</sup> <sup>a</sup> <sup>b</sup> <sup>c</sup> Gray's Anatomy 2008, p. 960.
- <sup>7</sup> <sup>^</sup> <sup>a</sup> <sup>b</sup> <sup>c</sup> <sup>d</sup> <sup>e</sup> <sup>f</sup> <sup>g</sup> <sup>h</sup> <sup>i</sup> <sup>j</sup> <sup>k</sup> <sup>l</sup> <sup>m</sup> <sup>n</sup> <sup>o</sup> <sup>p</sup> <sup>q</sup> <sup>r</sup> <sup>s</sup> <sup>t</sup> <sup>u</sup> <sup>v</sup> <sup>w</sup> <sup>x</sup> <sup>y</sup> <sup>z</sup> <sup>aa</sup> <sup>ab</sup> <sup>ac</sup> <sup>ad</sup> <sup>ae</sup> <sup>af</sup> <sup>ag</sup> <sup>ah</sup> <sup>ai</sup> <sup>aj</sup> <sup>ak</sup> <sup>al</sup> <sup>am</sup> <sup>an</sup> <sup>ao</sup> <sup>ap</sup> <sup>aq</sup> <sup>ar</sup> <sup>as</sup> <sup>at</sup> <sup>au</sup> <sup>av</sup> <sup>aw</sup> <sup>ax</sup> <sup>ay</sup> <sup>az</sup> <sup>ba</sup> <sup>bb</sup> <sup>bc</sup> <sup>bd</sup> <sup>be</sup> <sup>bf</sup> <sup>bg</sup> <sup>bh</sup> <sup>bi</sup> <sup>bj</sup> <sup>bk</sup> <sup>bl</sup> <sup>bm</sup> <sup>bn</sup> <sup>bo</sup> Betts, J. Gordon (2013). *Anatomy & physiology* . pp. 787–846. ISBN 1-938168-13-5. Retrieved 11 August 2014.
- <sup>8</sup> <sup>^</sup> Guyton & Hall 2011, pp. 101, 157.
- <sup>9</sup> <sup>^</sup> <sup>a</sup> <sup>b</sup> <sup>c</sup> Guyton & Hall 2011, pp. 105–07.
- <sup>10</sup> <sup>^</sup> Guyton & Hall 2011, pp. 1039–41.
- <sup>11</sup> <sup>^</sup> <sup>a</sup> <sup>b</sup> <sup>c</sup> "Cardiovascular diseases (CVDs) Fact sheet N°317 March 2013" . WHO. World Health Organization. Archived from the original on 19 September 2014 . Retrieved 20 September 2014.
- <sup>12</sup> <sup>^</sup> <sup>a</sup> <sup>b</sup> <sup>c</sup> Longo, Dan; Fauci, Anthony; Kasper, Dennis; Hauser, Stephen; Jameson, J.; Loscalzo, Joseph (August 11, 2011). *Harrison's Principles of Internal Medicine* (18 ed.). McGraw-Hill Professional. p. 1811. ISBN 978-0-07-174889-6.
- <sup>13</sup> <sup>^</sup> Graham, I; Atar, D; Borch-Johnsen, K; Boysen, G; Burell, G; Cifkova, R; Dallongeville, J; De Backer, G; Ebrahim, S; Gjelsvik, B; Herrmann-Lingen, C; Hoes, A; Humphries, S; Knapton, M; Perk, J; Priori, SG; Pyörälä, K; Reiner, Z; Ruilope, L; Sans-Menéndez, S; Scholte op Reimer, W; Weissberg, P; Wood, D; Yarnell, J; Zamorano, JL; Walma, E; Fitzgerald, T; Cooney, MT; Dudina, A; European Society of Cardiology (ESC) Committee for Practice Guidelines, (CPG) (Oct 2007). "European guidelines on cardiovascular disease prevention in clinical practice: executive summary: Fourth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (Constituted by representatives of nine societies and by invited experts)". *European Heart Journal*. **28** (19): 2375–414. doi:10.1093/eurheartj/ehm316 . PMID 17726041 .
- <sup>14</sup> <sup>^</sup> "Gray's Anatomy of the Human Body – 6. Surface Markings of the Thorax" . Bartleby.com. Archived from the original on 20 November 2010 . Retrieved 18 October 2010.
- <sup>15</sup> <sup>^</sup> Dorland's (2012). *Dorland's Illustrated Medical Dictionary* (32nd ed.). Elsevier. p. 1461. ISBN 978-1-4160-6257-8
- <sup>16</sup> <sup>^</sup> Bianco, Carl. "How Your Heart Works" . HowStuffWorks. Archived from the original on 29 July 2016 . Retrieved 14 August 2016.
- <sup>17</sup> <sup>^</sup> <sup>a</sup> <sup>b</sup> <sup>c</sup> Gray's Anatomy 2008, pp. 960–62.
- <sup>18</sup> <sup>^</sup> Gray's Anatomy 2008, pp. 964–67.
- <sup>19</sup> <sup>^</sup> Pocock, Gillian (2006). *Human Physiology*. Oxford University Press. p. 264. ISBN 978-0-19-856878-0.
- <sup>20</sup> <sup>^</sup> <sup>a</sup> <sup>b</sup> <sup>c</sup> <sup>d</sup> Gray's Anatomy 2008, pp. 966–67.
- <sup>21</sup> <sup>^</sup> Gray's Anatomy 2008, p. 970.
- <sup>22</sup> <sup>^</sup> University of Minnesota. "Papillary Muscles" . Atlas of Human Cardiac Anatomy. Archived from the original on 17 March 2016 . Retrieved 7 March 2016.
- <sup>23</sup> <sup>^</sup> "pectinate muscle" . The Free Dictionary . Retrieved 2016-07-31.
- <sup>24</sup> <sup>^</sup> "The human proteome in heart - The Human Protein Atlas" . www.proteinatlas.org. Retrieved 2017-09-29.
- <sup>25</sup> <sup>^</sup> Uhlen, Matthias; Fagerberg, Linn; Hallström, Björn M.; Lindskog, Cecilia; Oksvold, Per; Mardinoglu, Adil; Sivertsson, Åsa; Kampf, Caroline; Sjöstedt, Evelina (2015-01-23). "Tissue-based map of the human proteome" . *Science*. **347** (6220): 1260419. doi:10.1126/science.1260419 . ISSN 0036-8075 . PMID 25613900 .
- <sup>26</sup> <sup>^</sup> Lindskog, Cecilia; Linné, Jerker; Fagerberg, Linn; Hallström, Björn M.; Sundberg, Carl Johan; Lindholm, Malene; Huss, Mikael; Kampf, Caroline; Choi, Howard (2015-06-25). "The human cardiac and skeletal muscle proteomes defined by transcriptomics and antibody-based profiling" . *BMC Genomics*. **16**: 475. doi:10.1186/s12864-015-1686-y . ISSN 1471-2164 .
- <sup>27</sup> <sup>^</sup> Gray's Anatomy 2008, p. 959.
- <sup>28</sup> <sup>^</sup> J., Tortora, Gerard (2009). *Principles of human anatomy* . Nielsen, Mark T. (Mark Thomas) (11th ed.). Hoboken, NJ: J. Wiley. ISBN 9780471789314. OCLC 213300667 .
- <sup>29</sup> <sup>^</sup> Davidson's 2010, p. 525.
- <sup>30</sup> <sup>^</sup> Gray's Anatomy 2008, p. 981.
- <sup>31</sup> <sup>^</sup> <sup>a</sup> <sup>b</sup> <sup>c</sup> Gray's Anatomy 2008, p. 982.
- <sup>32</sup> <sup>^</sup> Davidson's 2010, p. 526.
- <sup>33</sup> <sup>^</sup> Gray's Anatomy 2008, p. 945.
- <sup>34</sup> <sup>^</sup> "Main Frame Heart Development" . Meddean.luc.edu. Archived from the original on 16 November 2001 . Retrieved 17 October 2010.
- <sup>35</sup> <sup>^</sup> DuBose, T. J.; Cunyus, J. A.; Johnson, L. (1990). "Embryonic Heart Rate and Age". *J Diagn Med Sonography* . **6** (3): 151–57. doi:10.1177/875647939000600306
- <sup>36</sup> <sup>^</sup> DuBose, TJ (1996) *Fetal Sonography*, pp. 263–74; Philadelphia: WB Saunders ISBN 0-7216-5432-0
- <sup>37</sup> <sup>^</sup> DuBose, Terry J. (2011-07-26) *Sex, Heart Rate and Age* Archived 2 May 2014 at the Wayback Machine .. obgyn.net
- <sup>38</sup> <sup>^</sup> Guyton & Hall 2011, pp. 110–13.
- <sup>39</sup> <sup>^</sup> <sup>a</sup> <sup>b</sup> Berry, William; McKenzie, Catherine (1 January 2010). "Use of inotropes in critical care" . *Clinical Pharmacist*. **2**: 395. Archived from the original on 28 November 2016.
- <sup>40</sup> <sup>^</sup> <sup>a</sup> <sup>b</sup> Bersten, Andrew (2013). *Oh's Intensive Care Manual* (7. ed.). London: Elsevier Health Sciences. pp. 912–22. ISBN 978-0-7020-4762-6
- <sup>41</sup> <sup>^</sup> Pocock, Gillian (2006). *Human Physiology* (Third ed.). Oxford University Press. p. 266. ISBN 978-0-19-856878-0.
- <sup>42</sup> <sup>^</sup> Antz, Matthias; et al. (1998). "Electrical Conduction Between the Right Atrium and the Left Atrium via the Musculature of the Coronary Sinus". *Circulation*. **98** (17): 1790–95. doi:10.1161/01.CIR.98.17.1790 . PMID 9788835 .
- <sup>43</sup> <sup>^</sup> De Ponti, Roberto; et al. (2002). "Electroanatomic Analysis of Sinus Impulse Propagation in Normal Human Atria". *Journal of Cardiovascular Electrophysiology* . **13** (1): 1–10. doi:10.1046/j.1540-8167.2002.00001.x . PMID 11843475 .
- <sup>44</sup> <sup>^</sup> "Definition of SA node" . MedicineNet.com. 27 April 2011. Archived from the original on 1 August 2012 . Retrieved 7 June 2012.
- <sup>45</sup> <sup>^</sup> "Purkinje Fibers" . About.com. 9 April 2012. Archived from the original on 14 April 2012 . Retrieved 7 June 2012.
- <sup>46</sup> <sup>^</sup> Guyton & Hall 2011, pp. 115–20.
- <sup>47</sup> <sup>^</sup> Davis, J. P.; Tikunova, S. B. (2008). "Ca<sup>2+</sup> exchange with troponin C and cardiac muscle dynamics". *Cardiovascular Research*. **77** (4): 619–26. doi:10.1093/cvr/cvm098 . PMID 18079104 .
- <sup>48</sup> <sup>^</sup> Ostchega, Y; Porter, K. S.; Hughes, J; Dillon, C. F.; Nwankwo, T (2011). "Resting pulse rate reference data for children, adolescents and adults, United States 1999-2008" (PDF). *National health statistics reports* (41): 1–16. PMID 21905522 . Archived (PDF) from the original on 23 June 2017.
- <sup>49</sup> <sup>^</sup> Hall, Arthur C. Guyton, John E. (2005). *Textbook of medical physiology* (11th ed.). Philadelphia: W.B. Saunders. pp. 116–22. ISBN 978-0-7216-0240-0
- <sup>50</sup> <sup>^</sup> Guyton & Hall 2011, p. 208.
- <sup>51</sup> <sup>^</sup> Guyton & Hall 2011, p. 212.
- <sup>52</sup> <sup>^</sup> <sup>a</sup> <sup>b</sup> <sup>c</sup> "Cardiovascular diseases (CVDs)" . World Health Organization. Archived from the original on 10 March 2016 . Retrieved 9 March 2016.
- <sup>53</sup> <sup>^</sup> "Your Heart Failure Healthcare Team" . www.heart.org. Archived from the original on 10 March 2016 . Retrieved 9 March 2016.
- <sup>54</sup> <sup>^</sup> <sup>a</sup> <sup>b</sup> "Different heart diseases" . World Heart Federation . Archived from the original on 12 March 2016 . Retrieved 9 March 2016.
- <sup>55</sup> <sup>^</sup> Harrison's 2011, p. 1501.
- <sup>56</sup> <sup>^</sup> Davidson's 2010, p. 554.
- <sup>57</sup> <sup>^</sup> <sup>a</sup> <sup>b</sup> <sup>c</sup> <sup>d</sup> <sup>e</sup> Ponikowski, Piotr; Voors, Adriaan A.; Anker, Stefan D.; Bueno, Héctor; Cleland, John G. F.; Coats, Andrew J. S.; Falk, Volkmar; González-Juanatey, José Ramón; Harjola, Veli-Pekka (August 2016). "2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: The Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC). Developed with the special contribution of the Heart Failure Association



- (HFA) of the ESC". *European Journal of Heart Failure*. **18** (8): 891–975. doi:10.1002/ehf.592 . ISSN 1879-0844 . PMID 27207191 .
58. <sup>a</sup> <sup>b</sup> Vahanian, Alec; Alfieri, Ottavio; Andreotti, Felicità; Antunes, Manuel J.; Barón-Esquivias, Gonzalo; Baumgartner, Helmut; Borger, Michael Andrew; Carrel, Thierry P.; De Bonis, Michele (October 2012). "Guidelines on the management of valvular heart disease (version 2012): the Joint Task Force on the Management of Valvular Heart Disease of the European Society of Cardiology (ESC) and the European Association for Cardio-Thoracic Surgery (EACTS)". *European Journal of Cardio-Thoracic Surgery: Official Journal of the European Association for Cardio-Thoracic Surgery*. **42** (4): S1–44. doi:10.1093/ejcts/ezs455 . ISSN 1873-734X . PMID 22922698 .
  59. <sup>a</sup> Davidson's 2010, pp. 612–13.
  60. <sup>a</sup> <sup>b</sup> Kirchhof, Paulus; Benussi, Stefano; Kotecha, Dipak; Ahlsson, Anders; Atar, Dan; Casadei, Barbara; Castella, Manuel; Diener, Hans-Christoph; Heidbuchel, Hein (November 2016). "2016 ESC Guidelines for the management of atrial fibrillation developed in collaboration with EACTS". *Europace: European Pacing, Arrhythmias, and Cardiac Electrophysiology: Journal of the Working Groups on Cardiac Pacing, Arrhythmias, and Cardiac Cellular Electrophysiology of the European Society of Cardiology*. **18** (11): 1609–1678. doi:10.1093/europace/euw295 . ISSN 1532-2092 . PMID 27567465 .
  61. <sup>a</sup> <sup>b</sup> <sup>c</sup> European Society of Cardiology (ESC); European Heart Rhythm Association (EHRA); Brignole, Michele; Auricchio, Angelo; Baron-Esquivias, Gonzalo; Bordachar, Pierre; Boriani, Giuseppe; Breithardt, Ole-A.; Cleland, John (August 2013). "2013 ESC guidelines on cardiac pacing and cardiac resynchronization therapy: the task force on cardiac pacing and resynchronization therapy of the European Society of Cardiology (ESC). Developed in collaboration with the European Heart Rhythm Association (EHRA)". *Europace: European Pacing, Arrhythmias, and Cardiac Electrophysiology: Journal of the Working Groups on Cardiac Pacing, Arrhythmias, and Cardiac Cellular Electrophysiology of the European Society of Cardiology*. **15** (8): 1070–1118. doi:10.1093/europace/eut206 . ISSN 1532-2092 . PMID 23801827 .
  62. <sup>a</sup> Blomström-Lundqvist, Carina; Scheinman, Melvin M.; Aliot, Etienne M.; Alpert, Joseph S.; Calkins, Hugh; Camm, A. John; Campbell, W. Barton; Haines, David E.; Kuck, Karl H. (2003-10-14). "ACC/AHA/ESC guidelines for the management of patients with supraventricular arrhythmias—executive summary: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and the European Society of Cardiology Committee for Practice Guidelines (Writing Committee to Develop Guidelines for the Management of Patients With Supraventricular Arrhythmias)". *Circulation*. **108** (15): 1871–1909. doi:10.1161/01.CIR.0000091380.04100.84 . ISSN 1524-4539 . PMID 14557344 .
  63. <sup>a</sup> Davidson's 2010, pp. 638–639.
  64. <sup>a</sup> Baumgartner, Helmut; Bonhoeffer, Philipp; De Groot, Natasja M. S.; de Haan, Fokko; Deanfield, John Erik; Galie, Nazzareno; Gatzoulis, Michael A.; Gohlke-Baerwolf, Christa; Kaemmerer, Harald (December 2010). "ESC Guidelines for the management of grown-up congenital heart disease (new version 2010)". *European Heart Journal*. **31** (23): 2915–2957. doi:10.1093/eurheartj/ehq249 . ISSN 1522-9645 . PMID 20801927 .
  65. <sup>a</sup> Harrison's 2011, p. 1458–65.
  66. <sup>a</sup> Davidson's 2010, pp. 527–34.
  67. <sup>a</sup> Britton, the editors Nicki R. Colledge, Brian R. Walker, Stuart H. Ralston; illustrated by Robert (2010). *Davidson's principles and practice of medicine* (21st ed.). Edinburgh: Churchill Livingstone/Elsevier. pp. 522–36. ISBN 978-0-7020-3084-0
  68. <sup>a</sup> <sup>b</sup> <sup>c</sup> Davidson's 2010, pp. 522–36.
  69. <sup>a</sup> <sup>b</sup> Talley, Nicholas J.; O'Connor, Simon. *Clinical Examination*. Churchill Livingstone. pp. 76–82. ISBN 978-0-7295-4198-5
  70. <sup>a</sup> <sup>b</sup> Davidson's 2010, pp. 556–59.
  71. <sup>a</sup> Coven, David; Yang, Eric. "Acute Coronary Syndrome Workup". *Medscape*. Archived from the original on 6 August 2016. Retrieved 14 August 2016.
  72. <sup>a</sup> Davidson's 2010, pp. 531.
  73. <sup>a</sup> Harrison's 2011, p. 1534.
  74. <sup>a</sup> Davidson's 2010, pp. 521–640.
  75. <sup>a</sup> <sup>b</sup> <sup>c</sup> <sup>d</sup> Davidson's 2010, pp. 528–30.
  76. <sup>a</sup> Armstrong, William F.; Ryan, Thomas; Feigenbaum, Harvey (2010). *Feigenbaum's Echocardiography*. Lippincott Williams & Wilkins. ISBN 978-0-7817-9557-9 Archived from the original on 23 April 2016.
  77. <sup>a</sup> Authors/Task Force Members;; Piepoli, Massimo F.; Hoes, Arno W.; Agewall, Stefan; Albus, Christian; Brotons, Carlos; Catapano, Alberico L.; Cooney, Marie-Therese; Corrà, Ugo (September 2016). "2016 European Guidelines on cardiovascular disease prevention in clinical practice: The Sixth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (constituted by representatives of 10 societies and by invited experts) Developed with the special contribution of the European Association for Cardiovascular Prevention & Rehabilitation (EACPR)". *Atherosclerosis*. **252**: 207–274. doi:10.1016/j.atherosclerosis.2016.05.037 . ISSN 1879-1484 . PMID 27664503 .
  78. <sup>a</sup> <sup>b</sup> Koli, Philippe; Windecker, Stephan; Alfonso, Fernando; Collet, Jean-Philippe; Cremer, Jochen; Falk, Volkmar; Filippatos, Gerasimos; Hamm, Christian; Head, Stuart J. (October 2014). "2014 ESC/EACTS Guidelines on myocardial revascularization: the Task Force on Myocardial Revascularization of the European Society of Cardiology (ESC) and the European Association for Cardio-Thoracic Surgery (EACTS). Developed with the special contribution of the European Association of Percutaneous Cardiovascular Interventions (EAPCI)". *European Journal of Cardio-Thoracic Surgery: Official Journal of the European Association for Cardio-Thoracic Surgery*. **46** (4): 517–592. doi:10.1093/ejcts/ezu366 . ISSN 1873-734X . PMID 25173601 .
  79. <sup>a</sup> Davidson's 2010, pp. 585–88, 614–23.
  80. <sup>a</sup> <sup>b</sup> <sup>c</sup> "Anatomy of the Heart". *University of Sydney Online Museum*. Archived from the original on 18 August 2016. Retrieved 2 August 2016.
  81. <sup>a</sup> <sup>b</sup> <sup>c</sup> <sup>d</sup> Meletis, John; Konstantopoulos, Kostas (2010). "The Beliefs, Myths, and Reality Surrounding the Word Hema (Blood) from Homer to the Present". *Anemia*. **2010**: 1–6. doi:10.1155/2010/857657 .
  82. <sup>a</sup> Katz, A. M. (1 May 2008). "The "Modern" View of Heart Failure: How Did We Get Here?". *Circulation: Heart Failure*. **1** (1): 63–71. doi:10.1161/CIRCHEARTFAILURE.108.772756 .
  83. <sup>a</sup> <sup>b</sup> <sup>c</sup> <sup>d</sup> <sup>e</sup> <sup>f</sup> <sup>g</sup> Aird, W. C. (July 2011). "Discovery of the cardiovascular system: from Galen to William Harvey". *Journal of Thrombosis and Haemostasis*. **9**: 118–29. doi:10.1111/j.1538-7836.2011.04312.x . PMID 21781247 .
  84. <sup>a</sup> Michelakis, E. D. (19 June 2014). "Pulmonary Arterial Hypertension: Yesterday, Today, Tomorrow". *Circulation Research*. **115** (1): 109–14. doi:10.1161/CIRCRESAHA.115.301132 .
  85. <sup>a</sup> West, John (2008). "Ibn al-Nafis, the pulmonary circulation, and the Islamic Golden Age". *Journal of Applied Physiology*. **105** (6): 1877–80. doi:10.1152/jappphysiol.91171.2008 . PMC 2612469 . PMID 18845773 .
  86. <sup>a</sup> Bondke Persson, A.; Persson, P. B. (2014). "Form and function in the vascular system". *Acta Physiologica*. **211** (3): 468–70. doi:10.1111/apha.12309 .
  87. <sup>a</sup> <sup>b</sup> West, J. B. (30 May 2014). "Galen and the beginnings of Western physiology". *AJP: Lung Cellular and Molecular Physiology*. **307** (2): L121–L128. doi:10.1152/ajplung.00123.2014 .
  88. <sup>a</sup> Silverman, M. E. (13 June 2006). "Why Does the Heart Beat?: The Discovery of the Electrical System of the Heart". *Circulation*. **113** (23): 2775–81. doi:10.1161/CIRCULATIONAHA.106.616771 . PMID 16769927 .
  89. <sup>a</sup> Cooley, Denton A. (2011). "Recollections of the Early Years of Heart Transplantation and the Total Artificial Heart". *Artificial Organs*. **35** (4): 353–57. doi:10.1111/j.1525-1594.2011.01235.x . PMID 21501184 .
  90. <sup>a</sup> Miniati, Douglas N.; Robbins, Robert C. (2002). "Heart transplantation: a thirty-year perspective: A Thirty-Year Perspective". *Annual Review of Medicine*. **53** (1): 189–205. doi:10.1146/annurev.med.53.082901.104050 .
  91. <sup>a</sup> Neubauer, Stefan (15 March 2007). "The Failing Heart – An Engine Out of Fuel". *New England Journal of Medicine*. **356** (11): 1140–51. doi:10.1056/NEJMr063052 . PMID 17360992 .
  92. <sup>a</sup> <sup>b</sup> <sup>c</sup> "Heart". *The Watkins Dictionary of Symbols*. ISBN 978-1-78028-357-9.
  93. <sup>a</sup> Rosner, Fred (1995). *Medicine in the Bible and the Talmud : selections from classical Jewish sources* (Augm. ed.). Hoboken, NJ: KTAV Pub. House. pp. 87–96. ISBN 978-0-88125-506-5.
  94. <sup>a</sup> *Britannica, lb* Archived 7 January 2009 at the Wayback Machine.; *Slider, Ab, Egyptian heart and soul conception* [permanent dead link]. The word was also transcribed by Wallis Budge as Ab.
  95. <sup>a</sup> Allen, James P. (2014). *Middle Egyptian : an introduction to the language and culture of hieroglyphs* (3rd ed.). pp. 453, 465. ISBN 978-1-107-66328-2.
  96. <sup>a</sup> Taylor, John H. (2001). *Death and the afterlife in ancient Egypt*. Chicago: The University of Chicago Press. pp. 35–38. ISBN 978-0-226-79164-7.
  97. <sup>a</sup> Xigui, Qiu; Mattos, Gilbert L. (2000). *Chinese writing = Wenzi-xue-gaiyao*. Berkeley: Society for the Study of Early China [u.a.]. p. 176. ISBN 1-55729-071-7.
  98. <sup>a</sup> MDBG online dictionary. "心" Archived 4 October 2016 at the Wayback Machine..
  99. <sup>a</sup> Rogers, Flaws, Bob (2007). *Statements of fact in traditional Chinese medicine* (3rd ed.). Boulder, Colo.: Blue Poppy Press. p. 47. ISBN 978-0-936185-52-1.
  100. <sup>a</sup> Wiseman, Nigel and Ye, Feng (1998). *A practical dictionary of Chinese medicine* (1st ed.). Brookline, Mass.: Paradigm Publications. p. 260. ISBN 978-0-912111-54.
  101. <sup>a</sup> Sellmer, Sven (2004), "The Heart in the *Rg veda*", in Piotr Balcerowicz; Marek Meior, *Essays in Indian Philosophy, Religion and Literature*, Delhi: Motilal Banarsidass Publishers, pp. 71–83, ISBN 978-81-208-1978-8, archived from the original on 6 December 2016
  102. <sup>a</sup> Lanman, Charles Rockwell (1996). *A Sanskrit reader : text and vocabulary and notes* (repr ed.). Delhi: Motilal Banarsidass. p. 287. ISBN 978-81-208-1363-2.
  103. <sup>a</sup> Aristotle. *On the Parts of Animals*. book 3, ch. 4. Archived from the original on 14 August 2016 (*De partibus animalium*)

104. <sup>^</sup> [Galen](#), *De usu partium corporis humani* ("The Use of the Parts of the Human Body"), book 6.
105. <sup>^</sup> Sandstrom, Alan (1991) *Corn is Our Blood*. University of Oklahoma Press. pp. 239–40. [ISBN 0-8061-2403-2](#)
106. <sup>^</sup> Kurian G (2001). "Sacred Heart of Jesus". *Nelson's Dictionary of Christianity: The Authoritative Resource on the Christian World* . Thomas Nelson Inc. [ISBN 978-1-4185-3981-8](#).
107. <sup>^</sup> Murray, Tom Devonshire Jones; Linda Murray; Peter (2013). "Heart". *The Oxford dictionary of christian art and architecture* (Second ed.). Corby: Oxford University Press. [ISBN 978-0-19-968027-6](#).
108. <sup>^</sup> *Indonesia Magazine*, **25** (1994), p. 67
109. <sup>^</sup> Abdenmour, Samia (2010) "Firakh mahshiya wi mihammara", recipe 117, *Egyptian Cooking: And Other Middle Eastern Recipes* , American University in Cairo Press [ISBN 977-424-926-7](#).
110. <sup>^</sup> Kennedy, Diana (2013) *My Mexico: A Culinary Odyssey with Recipes* , University of Texas Press. p. 100. [ISBN 0-292-74840-X](#).
111. <sup>^</sup> Sacharow, Alla (1993) *Classic Russian Cuisine: A Magnificent Selection of More Than 400 Traditional Recipes* . [ISBN 1-55970-174-9](#)
112. <sup>^</sup> Rombauer, Irma S. ; Becker, Marion Rombauer; Becker, Ethan (1975). *The Joy of Cooking*. The Bobbs-Merrill Company. p. 508. [ISBN 978-0-02-604570-4](#).
113. <sup>^</sup> Schwabe, Calvin W. (1979) *Unmentionable Cuisine*, University of Virginia Press, [ISBN 0-8139-1162-1](#), p. 96
114. <sup>^</sup> Rombauer, Irma S. and Rombauer Becker, Marion (1975) *The Joy of Cooking*, p. 508
115. <sup>^</sup> Torode, John (2009) *Beef: And Other Bovine Matters* , Taunton Press, [ISBN 1-60085-126-6](#), p. 230
116. <sup>^</sup> Milsom, Jennie (2009) *The Connoisseur's Guide to Meat*. Sterling Publishing Company. p. 171. [ISBN 1-4027-7050-2](#)
117. <sup>^</sup> Dobson, Geoffrey P (August 2003). "On Being the Right Size: Heart Design, Mitochondrial Efficiency and Lifespan Potential". *Clinical and Experimental Pharmacology and Physiology*. **30** (8): 590–97. doi:10.1046/j.1440-1681.2003.03876.x .
118. <sup>^</sup> Hyman, L. Henrietta (15 September 1992). *Hyman's Comparative Vertebrate Anatomy* . University of Chicago Press. pp. 448–. [ISBN 978-0-226-87013-7](#). Archived from the original on 6 December 2016.
119. <sup>^</sup> Shuttleworth, Trevor J., ed. (1988). *Physiology of Elasmobranch Fishes* . Berlin, Heidelberg: Springer Berlin Heidelberg. p. 3. [ISBN 978-3-642-73336-9](#).
120. <sup>^</sup> <sup>a</sup> <sup>b</sup> <sup>c</sup> <sup>d</sup> <sup>e</sup> <sup>f</sup> Romer, Alfred Sherwood; Parsons, Thomas S. (1977). *The Vertebrate Body*. Philadelphia, PA: Holt-Saunders International. pp. 437–42. [ISBN 0-03-910284-X](#).
121. <sup>^</sup> Osborne, June (1998). *The Ruby-Throated Hummingbird*. University of Texas Press. p. 14. [ISBN 0-292-76047-7](#).
122. <sup>^</sup> <sup>a</sup> <sup>b</sup> Grimm, Kurt A.; Lamont, Leigh A.; Tranquilli, William J.; Greene, Stephen A.; Robertson, Sheila A. (16 March 2015). *Veterinary Anesthesia and Analgesia* . John Wiley & Sons. p. 418. [ISBN 978-1-118-52620-0](#). Archived from the original on 6 December 2016.
123. <sup>^</sup> Colville, Thomas P.; Bassert, Joanna M. (10 March 2015). *Clinical Anatomy and Physiology for Veterinary Technicians* . Elsevier Health Sciences. p. 547. [ISBN 978-0-323-35620-6](#). Archived from the original on 6 December 2016.
124. <sup>^</sup> Crigg, Gordon; Johansen, Kjell (1987). "Cardiovascular Dynamics In Crocodylus Porosus Breathing Air And During Voluntary Aerobic Dives" (PDF). *Journal of Comparative Physiology B*. Springer-Verlag. **157** (3): 381–92. doi:10.1007/BF00693365 . Archived from the original on 3 July 2012 . Retrieved 3 July 2012.
125. <sup>^</sup> Axelsson, Michael; Craig, Franklin; Löfman, Carl; Nilsson, Stefan; Crigg, Gordon (1996). "Dynamic Anatomical Study Of Cardiac Shunting In Crocodiles Using High-Resolution Angioscopy" (PDF). *The Journal of Experimental Biology* . The Company of Biologists Limited. **199** (2): 359–65. PMID 9317958 . Archived (PDF) from the original on 3 March 2015. Retrieved 3 July 2012.
126. <sup>^</sup> Jurd, Richard David (January 2004). *Instant Notes Animal Biology* . Garland Science. p. 134. [ISBN 978-1-85996-325-8](#). Archived from the original on 6 December 2016.
127. <sup>^</sup> <sup>a</sup> <sup>b</sup> Ostrander, Gary Kent (2000). *The Laboratory Fish* . Elsevier. pp. 154–55. [ISBN 978-0-12-529650-2](#). Archived from the original on 6 December 2016.
128. <sup>^</sup> Farrell, Anthony P, ed. (1 June 2011). *Encyclopedia of Fish Physiology: From Genome to Environment* . Stevens, E Don; Cech, Jr., Joseph J; Richards, Jeffrey G Academic Press. p. 2315. [ISBN 978-0-08-092323-9](#). Archived from the original on 6 December 2016.
129. <sup>^</sup> Shukla, J.P. *Fish & Fisheries* . Rastogi Publications. pp. 154–55. [ISBN 978-81-7133-800-9](#). Archived from the original on 6 December 2016.
130. <sup>^</sup> <sup>a</sup> <sup>b</sup> Solomon, Eldra; Berg, Linda; Martin, Diana W. (15 September 2010). *Biology* . Cengage Learning. p. 939. [ISBN 1-133-17032-3](#). Archived from the original on 6 December 2016.
131. <sup>^</sup> "Meet our animals" . *Smithsonian National Zoological Park*. Archived from the original on 29 July 2016 . Retrieved 14 August 2016.
132. <sup>^</sup> Ladd, Prosser C (18 March 1991). *Comparative Animal Physiology, Environmental and Metabolic Animal Physiology* . John Wiley & Sons. pp. 537–. [ISBN 978-0-471-85767-9](#). Archived from the original on 6 December 2016.













Bibliography

Hall, John (2011). *Guyton and Hall textbook of medical physiology* (12th ed.). Philadelphia, Pa.: Saunders/Elsevier. ISBN 978-1-4160-4574-8.

Longo, Dan; Fauci, Anthony; Kasper, Dennis; Hauser, Stephen; Jameson, J.; Loscalzo, Joseph (August 11, 2011). *Harrison's Principles of Internal Medicine* (18 ed.). McGraw-Hill Professional. ISBN 978-0-07-174889-6.

Susan Standing; Neil R. Borley; et al., eds. (2008). *Gray's anatomy : the anatomical basis of clinical practice* (40th ed.). London: Churchill Livingstone. ISBN 978-0-8089-2371-8.

Nicki R. Colledge; Brian R. Walker; Stuart H. Ralston, eds. (2010). *Davidson's principles and practice of medicine* (21st ed.). Edinburgh: Churchill Livingstone/Elsevier. ISBN 978-0-7020-3085-7.

External links

- What Is the Heart? – NIH
- Atlas of Human Cardiac Anatomy
- Dissection review of the anatomy of the Human Heart including vessels, internal and external features
- Prenatal human heart development
- Animal hearts: fish, squid

Find more about  
**Heart**  
at Wikipedia's sister projects

Definitions from Wiktionary

Media from Wikimedia Commons

Quotations from Wikiquote

Texts from Wikisource

Textbooks from Wikibooks

Learning resources from Wikiversity

v · t · eAnatomy of the heart		
General	Surface	base · apex · sulci (coronary · interatrial · anterior interventricular · posterior interventricular) · borders (right · left)
	Internal	atria (interatrial septum · pectinate muscles · terminal sulcus) · ventricles (interventricular septum · trabeculae carneae · chordae tendineae · papillary muscle) · valves · cusps · atrioventricular septum
Chambers	Right heart	( <i>venae cavae</i> , <i>coronary sinus</i> ) → right atrium (atrial appendage, fossa ovalis, limbus of fossa ovalis, crista terminalis, valve of inferior vena cava , valve of coronary sinus) → tricuspid valve → right ventricle (infundibulum, moderator band/septomarginal trabecula) → pulmonary valve → ( <i>pulmonary artery and pulmonary circulation</i> )
	Left heart	( <i>pulmonary veins</i> ) → left atrium (atrial appendage) → mitral valve → left ventricle → aortic valve (aortic sinus) → ( <i>aorta and systemic circulation</i> )
Layers	Endocardium	heart valves
	Myocardium	Conduction system (cardiac pacemaker · SA node · Bachmann's bundle · AV node · bundle of His · bundle branches · Purkinje fibers)
	Pericardial cavity	pericardial sinus
	Pericardium	fibrous pericardium (sternopericardial ligaments) · serous pericardium (epicardium/visceral layer) · fold of left vena cava
Blood supply	Circulatory system · Coronary circulation · Coronary arteries	

v · t · ePhysiology of the cardiovascular system		
Heart	Cardiac output	Cardiac cycle · Cardiac output (Heart rate · Stroke volume) · Stroke volume (End-diastolic volume · End-systolic volume) · Afterload · Preload · Frank–Starling law · Cardiac function curve · Venous return curve · Wiggers diagram · Pressure volume diagram
	Ultrasound	Fractional shortening = ( End-diastolic dimension · End-systolic dimension ) / End-diastolic dimension · Aortic valve area calculation · Ejection fraction · Cardiac index
	Heart rate	Cardiac pacemaker · Chronotropic (Heart rate) · Dromotropic (Conduction velocity) · Inotropic (Contractility) · Bathmotropic (Excitability) · Lusitropic (Relaxation)
	Conduction	Conduction system · Cardiac electrophysiology · Action potential (cardiac · atrial · ventricular) · Effective refractory period · Pacemaker potential · Electrocardiography (P wave · PR interval · QRS complex · QT interval · ST segment · T wave ·

		U wave) · Hexaxial reference system
	Chamber pressure	Central venous · Right (atrial · ventricular) · pulmonary artery (wedge) · Left (atrial · ventricular) · Aortic
	Other	Ventricular remodeling
Vascular system/ Hemodynamics	Blood flow	Compliance · Vascular resistance · Pulse · Perfusion
	Blood pressure	Pulse pressure (Systolic · Diastolic) · Mean arterial pressure · Jugular venous pressure · Portal venous pressure
	Regulation of BP	Baroreflex · Kinin–kallikrein system · Renin–angiotensin system · Vasoconstrictors · Vasodilators · Autoregulation (Myogenic mechanism · Tubuloglomerular feedback · Cerebral autoregulation) · Paraganglia (Aortic body · Carotid body · Glomus cell)
Authority control	GND: 4024632-2 · NDL: 00571194	
Categories: Heart   Cardiac anatomy   Organs (anatomy)		