

Monte Carlo Methods

Introduction

Generating Samples

Markov Chain Monte Carlo + Discrepancy

Improving Efficiency

Selected Topics

Git website and repository
Canvas

Fred Hickernell, Fall 2025

Updated 2025 October 14




Improving Efficiency

Owen, Chapters 11– 12

Test 2 Oct 29

MATH 565 Monte Carlo Methods, Fred Hickernell, Fall 2025





Variance Reduction

$$\mu = \mathbb{E}[f(X)] = \mathbb{E}[g(X)], \quad X \sim \mathcal{U}[0,1]^d$$

Is it better to use

$$\hat{\mu}_{f,n} = \frac{1}{n} \sum_{i=0}^{n-1} f(X_i) \quad \text{or} \quad \hat{\mu}_{g,n} = \frac{1}{n} \sum_{i=0}^{n-1} g(X_i), \quad X_0, X_1, \dots \stackrel{\text{iid}}{\sim} \mathcal{U}[0,1]^d$$

to approximate μ ? Since both sample means are unbiased, we can compare their variances (mean squared errors):

$$\text{var}(\hat{\mu}_{f,n}) = \frac{\text{var}(f(X))}{n} \quad \text{var}(\hat{\mu}_{g,n}) = \frac{\text{var}(g(X))}{n}$$

so it depends on the variances of the two functions



Keister example

$$\mu = \int_{\mathbb{R}^d} \cos(\|t\|) \exp(-\|t\|^2) dt$$

$$\mathbf{x} = (\Phi(at_1), \dots, \Phi(at_d)), \quad \Phi \text{ is the CDF of the standard normal}$$

$$d\mathbf{x} = a^d \phi(at_1) \cdots \phi(at_d) dt = \frac{a^d \exp(-a^2 \|t\|^2/2)}{(2\pi)^{d/2}} dt$$

$$t = (\Phi^{-1}(x_1), \dots, \Phi^{-1}(x_d)) / a$$

$$\mu = \int_{[0,1]^d} \frac{(2\pi)^{d/2}}{a^d} \cos\left(\left\|(\Phi^{-1}(x_1), \dots, \Phi^{-1}(x_d))\right\| / a\right)$$

$$\times \exp\left(-\frac{2-a^2}{2a^2} \left\|(\Phi^{-1}(x_1), \dots, \Phi^{-1}(x_d))\right\|^2\right) d\mathbf{x}, \quad (a^2 \leq 2)$$

Whole family $f(\mathbf{x}; a)$ for which $\mu = \mathbb{E}[f(X; a)]$ for $X \sim \mathcal{U}[0,1]^d$



Variable transformations

Suppose that you want to approximate

$$\mu = \int_{\mathcal{T}} g(\mathbf{t}) \, d\mathbf{t}$$

by Monte Carlo. You need to write it as an expectation. Let $\mathbf{t} = \boldsymbol{\psi}(\mathbf{x})$, for some **invertible transformation**, $\boldsymbol{\psi} : \mathcal{X} \rightarrow \mathcal{T}$. Define the **Jacobian determinant** as

$$J(\mathbf{x}; \boldsymbol{\psi}) := \det \left(\left(\frac{\partial \psi_i}{\partial x_j} \right)_{i,j=1}^d \right), \quad J(\mathbf{t}; \boldsymbol{\psi}^{-1}) := \det \left(\left(\frac{\partial \psi_i^{-1}}{\partial t_j} \right)_{i,j=1}^d \right), \quad J(\boldsymbol{\psi}(\mathbf{x}); \boldsymbol{\psi}^{-1}) = \frac{1}{J(\mathbf{x}; \boldsymbol{\psi})}$$

Then

$$\mu = \int_{\mathcal{X}} g(\boldsymbol{\psi}(\mathbf{x})) |J(\mathbf{x}; \boldsymbol{\psi})| \, d\mathbf{x} = \int_{\mathcal{X}} \underbrace{\frac{g(\boldsymbol{\psi}(\mathbf{x}))}{\varrho_{\mathbf{X}}(\mathbf{x})} |J(\mathbf{x}; \boldsymbol{\psi})|}_{f(\mathbf{x})} \varrho_{\mathbf{X}}(\mathbf{x}) \, d\mathbf{x} = \mathbb{E}_{\mathbf{X} \sim \varrho_{\mathbf{X}}} [f(\mathbf{X})]$$



Variable transformations

$$\mu = \int_{\mathcal{T}} g(t) \, dt = ?$$

Let $t = \psi(x)$, for some **invertible transformation**, $\psi : \mathcal{X} \rightarrow \mathcal{T}$. Then

$$\mu = \int_{\mathcal{X}} g(\psi(x)) |J(x; \psi)| \, dx = \int_{\mathcal{X}} \underbrace{\frac{g(\psi(x))}{\varrho(x)} |J(x; \psi)|}_{f(x)} \varrho(x) \, dx = \mathbb{E}_{\mathbf{X} \sim \varrho}[f(\mathbf{X})]$$

If $\mathbf{X} \sim \mathcal{U}[0,1]^d$ (because we know how generate **uniform** samples), then

$$\mu = \int_{[0,1]^d} \underbrace{g(\psi(x)) |J(x; \psi)|}_{f(x)} \, dx = \mathbb{E}_{\mathbf{X} \sim \mathcal{U}[0,1]^d}[f(\mathbf{X})]$$



Random variable transformations

Let

- T be a random variable with sample space $\mathcal{T}$ and density ϱ_T
- X be a random variable with sample space $\mathcal{X}$ and density ϱ_X
- $T = \psi(X)$, for some **invertible transformation**, $\psi : \mathcal{X} \rightarrow \mathcal{T}$. Then

$$\varrho_T(t) = \varrho_X(\psi^{-1}(t)) |J(t; \psi^{-1})|, \quad \varrho_T(\psi(x)) = \varrho_X(x) |J(\psi(x)); \psi^{-1})| = \frac{\varrho_X(x)}{|J(x; \psi)|}$$

$$\mu = \int_{\mathcal{T}} \underbrace{h(t) \varrho_T(t)}_{g(t)} dt = \mathbb{E}_{T \sim \varrho_T}[h(T)] = \mathbb{E}_{X \sim \varrho_X}[h(\psi(X))] = \int_{\mathcal{X}} h(\psi(x)) \varrho_X(x) dx$$

$$\text{or } = \int_{\mathcal{X}} \frac{h(\psi(x)) \varrho_T(\psi(x))}{\varrho_X(x)} |J(x; \psi)| \varrho_X(x) dx = \int_{\mathcal{X}} h(\psi(x)) \varrho_X(x) dx$$



Importance sampling

Suppose that you want to approximate

$$\mu = \mathbb{E}_{\mathbf{T} \sim \varrho_{\mathbf{T}}} [h(\mathbf{X})] = \int_{\mathcal{T}} h(\mathbf{t}) \varrho_{\mathbf{T}}(\mathbf{t}) \, d\mathbf{t} = \int_{\mathcal{T}} \underbrace{h(\mathbf{t}) \frac{\varrho_{\mathbf{T}}(\mathbf{t})}{\varrho_{\mathbf{X}}(\mathbf{t})}}_{f(\mathbf{t})} \varrho_{\mathbf{X}}(\mathbf{t}) \, d\mathbf{t} = \mathbb{E}_{\mathbf{X} \sim \varrho_{\mathbf{X}}} [f(\mathbf{X})]$$

likelihood ratio

We try to choose $\varrho_{\mathbf{X}}$ to make $\text{var}[f(\mathbf{X})]$ small when doing IID sampling. If $\mathbf{T} = \boldsymbol{\psi}(\mathbf{X})$ for $\mathbf{X} \sim \mathcal{X}[0,1]^d$ —since often find generate non-uniform samples from uniform ones—then

$$\mu = \int_{\mathcal{T}} h(\mathbf{t}) \varrho_{\mathbf{T}}(\mathbf{t}) \, d\mathbf{t} = \mathbb{E}_{\mathbf{T} \sim \varrho_{\mathbf{T}}} [h(\mathbf{T})] = \mathbb{E}_{\mathbf{X} \sim \mathcal{U}[0,1]^d} [h(\boldsymbol{\psi}(\mathbf{X}))] = \int_{[0,1]^d} h(\boldsymbol{\psi}(\mathbf{x})) \, d\mathbf{x}$$



Variable transformation / importance sampling cautions

- The choice of sampling density or variable transformation affects the variance of the sample mean
- You must choose the variable transformation so that
 - The Jacobian determinant, $J(\mathbf{x}; \boldsymbol{\psi})$, is **finite**
 - The probability density, $q_{\mathbf{X}}(\mathbf{x})$, does not **vanish**