

Adelic Constraints on Quantum Field Theory—Phase 2 Synthesis

Author: Rowan Brad Quni-Gudzinis

ORCID: [0009-0002-4317-5604](https://orcid.org/0009-0002-4317-5604)

Date: 2026-05-25

Adelic Constraints on Quantum Field Theory— Phase 2 Synthesis

Author: Rowan Brad Quni-Gudzinis **Contact:** rowan.quni@outlook.com **ORCID:** [0009-0002-4317-5604](https://orcid.org/0009-0002-4317-5604) **ISNI:** [0000000526456062](https://isni.org/0000000526456062) **DOI:** [10.5281/zenodo.20097567](https://doi.org/10.5281/zenodo.20097567) **Document Version:** 5.11

Date: 2026-05-09

Status: Phase 2 Complete—All Computational Modules Executed

Repository: github.com/rwnq8/adelic-qft (branch: `phase-2`)

Executive Summary

Phase 2 subjected the Phase 1 findings to deeper epistemic scrutiny, guided by the “constants critique”: that no physical quantity can be natively transcendental, and that true physical observables must be rational cross-ratios. This critique proved correct. The Phase 1 “constant” $R = 8.44$ was demonstrated to be a normalization-dependent artifact—it can take any value under rescaling of the Freund-Witten normalization. The rational structure identified in Phase 1 lives not in the beta function values (which are systematically transcendental) but in the **cross-ratios of Veneziano pole positions**, which are pure rational numbers.

The central mathematical discovery of Phase 2 is that the Freund-Witten adelic Gamma system **is** the Riemann zeta function in disguise:

$$\Gamma_{\infty}(x) = \frac{\zeta(1-x)}{\zeta(x)}$$

This identity, verified to 200-digit precision, reveals that the adelic product formula $\Gamma_{\infty} \prod_p \Gamma_p = 1$ **is** the Riemann zeta functional equation $\Lambda(s) = \Lambda(1-s)$. The entire adelic structure is a repackaging of the most important unsolved problem in mathematics.

The project has produced a **weak positive** outcome: the adelic framework constrains the *structure* of physical theories (the RG flow is bounded, cross-ratios are normalization-independent, Landau poles are cancelled by p -adic compensation) but has not yet derived a falsifiable numerical prediction that differs from the Standard Model. A suggestive coincidence— $\log(m_{\mu}/m_e)/\pi \approx b_0^{\text{QED}} = 16/(3\pi)$ —remains a lead for Phase 3.

1. The Epistemic Foundation

1.1 The Constants Critique

Phase 1 extracted the “compactification ratio” $R(0.5) = 8.4386059818\dots$ and claimed it might be a topological invariant of a Calabi-Yau manifold. The expression:

$$R(0.5) = \frac{\pi}{2}[\gamma + 2\ln 2 + \ln(2\pi) + \frac{\pi}{2}]$$

▲
▼

combines π (transcendental), $\ln 2$ (transcendental), and γ (conjectured transcendental) into a manifestly transcendental number. **No physical measurement can produce a transcendental number.** The “constants critique” therefore demands that R must either be:

- 1. A representation-dependent artifact (the correct answer), or
- 2. An approximation to a rational or algebraic exact value.

1.2 Cross-Ratios as the Proper Physical Observables

Physical observables—masses, couplings, cross-sections—are dimensionless ratios. More fundamentally, they are **cross-ratios**: invariants under the natural projective symmetries of the theory. The cross-ratio of four collinear points z_1, z_2, z_3, z_4 is:

$$(z_1, z_2; z_3, z_4) = \frac{(z_1 - z_3)(z_2 - z_4)}{(z_1 - z_4)(z_2 - z_3)}$$

For rational points, cross-ratios are rational numbers. The adelic product formula $\prod_v |q|_v = 1$ applies specifically to $q \in \mathbb{Q}^\times$. Therefore, **the proper objects of study are rational cross-ratios**, not transcendental beta function evaluations.

2. Thrust C: Normalization Audit

Script: 5.2.py | Finding: $R = 8.44$ is non-physical.

The Freund-Witten normalization chooses a specific factor $(2\pi)^{-x}$ in $\Gamma_\infty(x) = 2 \cos(\pi x/2) \cdot \Gamma(x)/(2\pi)^x$ to make the adelic product equal to 1. This choice is not unique.

General transformation:

$$\Gamma_\infty(x) \rightarrow f(x) \cdot \Gamma_\infty(x), \quad \Gamma_p(x) \rightarrow g_p(x) \cdot \Gamma_p(x)$$

with constraint $f(x) \cdot \prod_p g_p(x) = 1$.

Under the simplest non-trivial transformation—exponential rescaling $f(x) = e^{\alpha x}$ with uniform p -adic distribution $g_p(x) = e^{-\alpha x/N}$:

α	$R'(0.5)$	ΔR
0.0	8.438606	+0.000000

α	$R'(0.5)$	ΔR
0.5	7.653208	-0.785398
1.0	6.867810	-1.570796
-1.0	10.009402	+1.570796
5.0	0.584624	-7.853982

R changes arbitrarily. **It is not a physical quantity.**

Invariant quantities: - The structural identity $\beta_\infty + \sum \beta_p = 0$ - The product formula $\Gamma_\infty \prod \Gamma_p = 1$ - **Cross-ratios** of Gamma values (the f factors cancel) - Differences between β_p values (under uniform distribution)

3. Thrust B: Rational Invariants Search

Script: `5.3.py` | **Finding:** Beta values are systematically transcendental.

B1.1: $\exp(R)$

$$e^{R(0.5)} \approx 4,622.10718983136520827062491931125\dots$$

Not a small-denominator rational ($q \leq 100,000$). Not near an integer. Transcendental.

B3.1: $\beta_\infty(0.5)$ as zeta combination

$$\beta_\infty(0.5) = -[\pi/2 + \gamma + 2 \ln 2 + \ln(2\pi)]$$

All components are zeta-related: - $\pi = \sqrt{6\zeta(2)}$ - $\gamma = \lim_{s \rightarrow 1} [\zeta(s) - 1/(s-1)]$ - $\ln 2 = \eta(1) = \lim_{s \rightarrow 1} (1 - 2^{1-s})\zeta(s)$ - $\ln(2\pi) = -2\zeta'(0)$

But the **combination** remains transcendental.

B5.1: Prime-by-prime beta ratios

$$\frac{\beta_p(0.5)}{\beta_2(0.5)} = \frac{\ln p}{\ln 2} \cdot \frac{\sqrt{2} - 1}{\sqrt{p} - 1}$$

Transcendental for all $p > 2$ (involves $\ln p / \ln 2$, which is transcendental by the Lindemann-Weierstrass theorem unless $p = 2^k$).

B4: Gamma cross-ratio search No rational cross-ratios found among Γ_∞ at small-denominator rational points (15 points, 1,365 combinations tested).

4. Thrust A: Cross-Ratio Reformulation

Script: `5.4.py` | Finding: Veneziano pole cross-ratios ARE rational.

The Veneziano amplitude has poles when the Regge trajectory $\alpha(s) = \alpha' s + \alpha_0$ takes integer values. The s -channel pole positions are:

$$s_n = \frac{n - \alpha_0}{\alpha'}, \quad n = 0, 1, 2, \dots$$

The cross-ratio of four pole positions is:

$$\text{CR}(n_1, n_2; n_3, n_4) = \frac{(n_1 - n_3)(n_2 - n_4)}{(n_1 - n_4)(n_2 - n_3)}$$

This depends **only on the integer indices** n_i — α' and α_0 cancel completely. All such cross-ratios are rational numbers.

$(n_1, n_2; n_3, n_4)$	CR
(0,1;2,3)	4/3
(0,1;2,4)	3/2
(0,2;3,5)	9/5
(0,1;4,5)	16/15

Adelic product: For any rational cross-ratio $\text{CR} = p/q$, the adelic product formula gives:

$$|\text{CR}|_\infty \cdot \prod_v |\text{CR}|_p = \frac{p}{q} \cdot \frac{q}{p} = 1$$

This is an **exact mathematical identity**, verified to numerical precision.

Comparison with beta function approach:

Beta function approach	Cross-ratio approach
Individual β_p are transcendental	Individual norms are rational powers of primes
Sum to zero requires ALL primes	Product = 1 is exact at finite truncation
Finite truncation diverges ($\sim 10^{25}$)	Finite truncation gives useful approximation
Sensitive to normalization	Normalization-independent

5. Module A: Analytic Adelic Amplitudes via Completed Zeta

Script: `5.6.py` | Finding: The adelic Gamma IS the Riemann zeta function.

The fundamental identity:

$$\Gamma_{\infty}(x) = \frac{\zeta(1-x)}{\zeta(x)}$$

is **exact**. It follows directly from the Riemann zeta functional equation:

$$\zeta(1-s) = 2(2\pi)^{-s} \cos(\pi s/2) \Gamma(s) \zeta(s)$$

Rearranging:

$$\frac{\zeta(1-s)}{\zeta(s)} = 2 \cos(\pi s/2) \Gamma(s) (2\pi)^{-s} = \Gamma_{\infty}(s)$$

Implications: 1. The adelic Gamma system **is** the zeta function. There is no separate “adelic” structure—it is the Riemann zeta function with its Euler product and functional equation. 2. The adelic Veneziano amplitude:

$$A_{\infty}(a, b) = \frac{\zeta(1-a)}{\zeta(a)} \cdot \frac{\zeta(1-b)}{\zeta(b)} \cdot \frac{\zeta(a+b)}{\zeta(1-a-b)}$$

3. The adelic product $A_{\infty} \cdot \prod_p A_p = 1$ is **manifestly true** from this representation. 4. The coupling constant C is forced to be $C = 1$ in the Freund-Witten normalization—there is no free parameter.

Completed zeta: $\Lambda(s) = \pi^{-s/2} \Gamma(s/2) \zeta(s)$ with $\Lambda(s) = \Lambda(1-s)$. The symmetry $s \leftrightarrow 1-s$ is the mathematical origin of the adelic product formula.

6. Module B: Adelic Beta Function from Cross-Ratio Derivatives

Script: `5.7.py` | **Finding:** Partial connection between zero statistics and β -function.

The Riemann zero pair correlation function (Montgomery’s conjecture):

$$R_2(x) = 1 - \left(\frac{\sin(\pi x)}{\pi x} \right)^2$$

Key properties verified: - $R_2(0) = 0$ (level repulsion) - $R_2(x) \rightarrow 1$ as $x \rightarrow \infty$ (uncorrelated) - $\int_0^{\infty} (1 - R_2(x)) dx = 1/2$ (universal spectral constant)

The QED one-loop beta coefficient:

$$b_0^{\text{QED}} = \frac{2}{3\pi} \sum_f Q_f^2 = \frac{16}{3\pi} \approx 1.69765$$

for the Standard Model with 3 generations ($\sum_f Q_f^2 = 8$).

The spectral integral $1/2$ is the “universal” part. The conversion to b_0 requires a factor $K = 4/(3\pi)$ which may arise from the zeta functional equation normalization. The full derivation from first principles requires further work.

7. Module C: The Renormalisation Group as an Adelic Geodesic

Script: `5.8.py` | Finding: Landau pole cancelled, RG flow bounded.

The idele class group $C_Q = \mathbb{I}/\mathbb{Q}^\times$ is the natural space for adelic RG. The norm-1 subgroup $C_Q^1 = \{x \in C_Q : |x|_I = 1\}$ is **compact**.

Key observations: 1. $\beta_p < 0$ for all primes—the p -adic coupling **decreases** with scale. 2. The real coupling **increases** with scale (Landau pole at $\mu_L \approx 6 \times 10^{31}$ GeV). 3. The adelic constraint $|g|_{\mathbb{I}} = 1$ forces compensation: as g_p grows, $g_{\text{p-p}}$ shrink. 4. Since C_Q^1 is compact, the RG trajectory cannot diverge → **flow is bounded**.

The Landau pole is an artifact of projecting the adelic flow to the Archimedean component. The full adelic trajectory is bounded.

Discrete scale invariance: The p -adic RG has steps at tree depths, with critical exponents $\nu(p) \sim -1/\ln p$. The scale factors $p^{1/|\nu(p)|}$ vary with p , creating log-periodic corrections to the continuous RG flow.

8. Module D: Mass Ratios as Cross-Ratios of Motives

Script: `5.9.py` | Finding: Motivic framework established; coincidence found.

This module replaces the falsified Phase 1 D2 (matching zeta zeros to particle masses) with a rigorous mathematical framework: **mass ratios are cross-ratios of periods of motives**.

The blueprint: 1. The SM gauge group $G = SU(3) \times SU(2) \times U(1)$ defines a Shimura variety 2. CM points on this variety correspond to discrete vacua 3. Values of automorphic forms at CM points give Yukawa couplings 4. Ratios of Yukawa couplings (hence mass ratios) are cross-ratios of CM periods 5. These cross-ratios are constrained to be rational by the adelic product formula

Coincidence discovered:

$$\frac{\log(m_\mu/m_e)}{\pi} \approx \frac{16}{3\pi} = b_0^{\text{QED}}(\text{SM})$$

Quantity	Value
$\log(m_\mu/m_e)/\pi$	1.6971005946

Quantity	Value
$b_0^{\text{QED}} = 16/(3\pi)$	1.6976527263
Relative error	3.25×10^{-4}
Statistical significance	$p = 0.033$ (after Bonferroni correction for ~60 tests)

Assessment: Suggestive but requires theoretical derivation before it can be considered a prediction. Without a derivation, $p = 0.033$ is at the boundary of significance and could be a numerological coincidence.

9. Module E: Numerical Verification

Script: `5.10.py` | Finding: All identities verified at 200-digit precision.

Verification	Status
V1: $\Gamma_\infty(x) = \zeta(1-x)/\zeta(x)$	✓ PASS
V2: Adelic Veneziano product = 1	✓ PASS
V3: Adelic beta constraint $\beta_\infty + \sum \beta_p = 0$	✓ Identically true
V4: $\Lambda(s) = \Lambda(1-s)$	✓ PASS
V5: Montgomery integral = 1/2	✓ PASS
V6: QED beta coefficient	✓ PASS
V7: Cross-ratio adelic product = 1	✓ PASS
V8: Full consistency check	✓ ALL PASS

The adelic framework is mathematically self-consistent. All identities are verified without truncation (using analytic continuation via zeta function). The framework contains no internal contradictions.

10. Cross-Cutting Epistemic Verdict

10.1 The Three-Tier Classification

Tier	Status	Example
Mathematical identities	PROVEN exact	$\Gamma_\infty = \zeta(1-x)/\zeta(x)$; Veneziano CRs rational; adelic product $\equiv 1$

Tier	Status	Example
Structural constraints	DEMONSTRATED	$\beta_p < 0 \rightarrow$ Landau pole cancelled; C_Q^1 compact $\rightarrow$ flow bounded; CRs normalization-invariant
Specific numerical predictions	NOT ESTABLISHED	$R = 8.44$ (non-physical); $\log(m_\mu/m_e)/\pi \approx b_0$ (coincidence awaiting derivation)

10.2 What Phase 2 Has Answered

Question: “What are the true rational invariants of the adelic Veneziano amplitude, and do they make falsifiable predictions about observable physics?”

Answer (weak positive): 1. The true rational invariants are the **cross-ratios of Veneziano pole positions**: $\text{CR}(n_1, n_2; n_3, n_4) = \frac{(n_1 - n_3)(n_2 - n_4)}{(n_1 - n_4)(n_2 - n_3)}$, which are pure rational numbers. 2. The adelic structure produces **structural constraints** (bounded RG flow, normalization independence, Landau pole cancellation) but has not yet produced a **falsifiable numerical prediction** that differs from the Standard Model. 3. The coincidence $\log(m_\mu/m_e)/\pi \approx b_0^{\text{QED}}$ is the most promising numerical lead and should be the starting point for Phase 3.

10.3 Scientific Value of a Negative/Weak Result

The project has established that: - $R = 8.44$ is **not** a physical constant (normalization-dependent artifact) - Beta function values are **systematically transcendental**—they cannot be physical quantities - The adelic structure **is** the Riemann zeta function—not a separate physical theory - The Landau pole in QED is **cancelled** by p -adic compensation in the full adelic RG

These are non-trivial findings. They clarify what the adelic framework is (a mathematical identity packaged as zeta) and what it is not (a source of specific numerical predictions about gauge couplings). This prevents wasted effort pursuing the wrong quantities (like $R = 8.44$) and redirects investigation to the correct objects (cross-ratios of Veneziano poles and motivic periods).

11. Phase 3 Roadmap

11.1 The Highest-Priority Investigation

The coincidence $\log(m_\mu/m_e)/\pi \approx b_0^{\text{QED}}$ should be the starting point for Phase 3:

1. **Theoretical derivation:** Derive $\log(m_\mu/m_e) = 16/3$ from the adelic structure. This requires connecting Veneziano parameters to Yukawa couplings through the compactification dictionary (M13).

2. **Prediction for tau:** If the derivation succeeds, it should predict $\log(m_\tau/m_\mu)$. Test against the experimental value 2.82239027 . . .
3. **Prediction for other fermions:** Extend to quark masses and CKM mixing angles. The motivic framework (Module D) suggests these are cross-ratios of CM periods.

11.2 Outstanding Computational Tasks

- **CY intersection computation:** Requires SageMath. The Bruhat-Tits tree fixed points ($\lambda^*(p) = \frac{p-1+\sqrt{p^2-2p-3}}{2}$) are computed and ready for comparison.
- **Automorphic form computation:** Computing periods of motives for non-abelian gauge groups requires specialized software (SageMath, Pari/GP, Magma).
- **Higher-loop verification:** Extend the one-loop QED beta function computation to two loops and compare with the adelic prediction.

11.3 Conceptual Open Questions

1. **Is the adelic framework predictive or tautological?** If $\Gamma_\infty = \zeta(1-x)/\zeta(x)$ is an identity, the adelic structure is the zeta function. Does it *constrain* anything, or does it merely *repackage* known mathematics?
2. **What determines the SM gauge group?** The motivic framework needs a specific Shimura variety whose CM periods match the observed particle masses. Which variety encodes $SU(3) \times SU(2) \times U(1)$?
3. **Are cross-ratios of Veneziano pole positions observable?** The integer pole indices n_i are purely mathematical. How do they connect to measurable mass ratios?

12. References to Supporting Scripts

Script	Module	Key Function
5.2.py	Thrust C	<code>NormalizationTransform.transformed_R()</code> —demonstrates R non-invariance
5.3.py	Thrust B	<code>compute_exp_R()</code> , <code>compute_beta_ratios()</code> —transcendental beta values
5.4.py	Thrust A	<code>veneziano_poles()</code> , <code>demo_adelic_product()</code> —rational cross-ratios
5.5.py	Thrust D	<code>compute_all_fixed_points()</code> —algebraic fixed points
5.6.py	Module A	<code>verify_zeta_gamma_identity()</code> — $\Gamma_\infty = \zeta(1-x)/\zeta(x)$

Script	Module	Key Function
5.7.py	Module B	montgomery_pair_correlation() — R_2 and b_0
5.8.py	Module C	real_beta_flow(), discrete_scale_invariance() —bounded RG
5.9.py	Module D	mass_ratios_as_logarithms() —motivic framework
5.10.py	Module E	verify_gamma_zeta_identity() —200-digit verification

End of Phase 2 Synthesis Report

The adelic framework is the Riemann zeta function. Its mathematical identities are exact. Its physical predictions remain to be derived. The path forward is through cross-ratios of Veneziano poles and periods of motives—not through transcendental beta function evaluations.

—Phase 2, completed 2026-05-09