Joint Mean-Variance Inference for Nonlinear Mixed-Effects Models with Measurement Errors and Outliers

March, 2022

Ye et al.

1 Introduction

Nonlinear mixed-effects (NLME) models are widely used in many biological applications, such as pharmacokinetic analysis, HIV viral dynamics, and studies of growth and decay. Many NLME models are derived based on the underlying data-generation mechanisms, so they are also called *scientific models* or *mechanistic models*. They may be more desirable than empirical models such as linear mixed effects (LME) models and nonparametric mixed models, since they may provide more reliable predicted values for (unobserved) missing/censored values or future values. In many longitudinal studies, the within-individual repeated measurements exhibit large variations and these variations may change over time. It is important to understand the nature of the within-individual systematic and random variations in order to provide good predictions and efficient statistical inference.

Figure 1 shows the profiles of HIV viral load (HIV-1 RNA copies) longitudinal measurements on a sub-sample of individuals from an AIDS study during an anti-HIV treatment. We see that viral loads decline rapidly in the initial period (first few weeks), roughly following an exponential pattern. The variations of the within-individual repeated measurements appear to *change* over time. Moreover, there seem some *outliers* in the data, i.e., some data do not seem to follow the general patterns of the whole data. It is interesting to see if these large and time-varying within-individual variations may be partially explained by covariates such as CD4 cell counts. Based on some biological arguments, as well as a set of differential equations which describe the viral production and elimination processes, Wu and Ding (1999) showed that the viral load trajectories during the treatment may well be modelled by an NLME model.

/home/lang/Figures/ch9outlier.pdf

Figure 1: *Viral loads trajectories (in log_{10} scale) during an anti-HIV treatment.*

To see if the large and time-varying within-individual variations in longitudinal viral load data may be partially explained by time-varying covariates such as CD4 cell counts, we may model the within-individual variations with time-varying covariates and incorporate the large within-individual variations through random effects in the variance model. In other words, we may jointly model the mean and variance of the longitudinal viral load data: we model the mean based on an NLME model and model the variance using another mixed effects model. The advantages of modelling the within-individual variations, compared to NLME models without modelling the variance, are as follows: (a) we may gain good understanding of the nature of the within-individual variation and if this variation may be partially explained by covariates; (b) we may obtain more efficient estimates of the main parameters of interest; (c) as will be described in Section 2, modeling variance also allows us to conduct robust inference to address outliers in the data; and (d) we may provide more accurate predictions for the missing/censored response values or future values.

For example, by modelling the variance, we may obtain an more efficient estimate of initial viral decay rates, which allows us to better evaluate and compare different treatments since the initial viral decay rates may reflect the efficacies of the treatments (Wu and Ding, 1999). Understanding the nature of the within-individual variations in viral load also allows us to choose good timing for treatment interruption.

There is some literature on joint modelling mean and variance for longitudinal data (e.g., Lin et al. 1997; Pourahmadi 1999; Hedeker et al. 2008; German et al. 2022). To our knowledge, all previous work focused on *linear* models, either with random effects or without random effects. In this article, we extend previous work to NLME models, and we also address covariate measurement errors and outlying responses. NLME models are *nonlinear* in the random effects and the dimension of the random effects may be high, so computation for the joint model can be challenging. In this article, we propose a computationally efficient approximate method for approximate likelihood inference.

The article is organized as follows ...

2 The Models

2.1 The mean and variance models

Let y_{ij} be the response value for individual i at time t_{ij}, and let $\mathbf{y}_i = (y_{i1}, y_{i2}, \cdots, y_{in_i})^T$, $i = 1, 2, \cdots, n;\ j = 1, 2, \cdots, n_i$. We consider the following nonlinear mixed effects (NLME) model

$$\mathbf{y}_i = g(\mathbf{x}_i, \boldsymbol{\beta}, \mathbf{u}_i) + \mathbf{e}_i, \qquad i = 1, 2, \cdots, n;\ j = 1, 2, \cdots, n_i, \tag{1}$$

where $g(\cdot)$ is a known nonlinear function for the mean $E(\mathbf{y}_i)$, $\mathbf{x}_i$ contains covariates including time, vector $\boldsymbol{\beta} = (\beta_1, \beta_2, \cdots, \beta_p)$ contains (fixed) mean parameters, vector $\mathbf{u}_i$ contains random effects for the mean function, and $\mathbf{e}_i = (e_{i1}, e_{i2}, \cdots, e_{in_i})^T$ are within-individual errors. We assume that $\mathbf{u}_i \sim N(0, D)$, and $e_{i1}, \cdots e_{in_i}$ are conditionally independent given $\mathbf{u}_i$, with $e_{ij} \sim N(0, \sigma_{ij}^2)$. Here we allow the variances of the within-individual random errors, σ_{ij}^2, to be individual-specific and time-dependent. This allows great flexibility in modeling complicated longitudinal data, as discussed in the previous section. Let $\boldsymbol{\sigma}_i = (\sigma_{i1}^2, \cdots, \sigma_{in_i}^2)$.

To understand the nature of the within-individual variations and check if there are some systematic components in these variations, we investigate if the within-individual variances σ_{ij}^2 may be partially explained by time-varying covariates (say) $\mathbf{z}_i = (z_{i1}, \cdots, z_{in_i})$, where z_{ij} is the value of covariate z for individual i at time t_{ij}. Note that covariate $\mathbf{z}_i$ may be different from covariates $\mathbf{x}_i$. For simplicity, here we only consider one covariate, but the method can be easily extended to more than one covariates. Moreover, the within-individual variations σ_{ij}^2 may vary substantially across individuals. Thus, we model the within-individual variations and introduce additional random effects to account for the associations within an individual and variations across individuals, i.e., we consider the following model for the variance

$$\log(\sigma_{ij}^2) = \alpha_1 + \alpha_2 z_{ij} + \eta a_i, \qquad i = 1, \cdots, n, \quad j = 1, 2, \cdots, n_i, \tag{2}$$

where vector $\boldsymbol{\alpha} = (\alpha_1, \alpha_2)$ and η are fixed parameters, and a_i is a random effect for the variance model. We may assume that $\exp(a_i)$ follows an inverse gamma distribution or $a_i \sim N(0, 1)$. In the variance model (2), we model the log-transformation of the variances, $\log(\sigma_{ij}^2)$, since the variances σ_{ij}^2 are strictly positive and the range of the log-transformation $\log(\sigma_{ij}^2)$ is $(-\infty, \infty)$, which allows us to introduce arbitrary covariate in the model.

In the above mean and variance models, we jointly model the conditional mean $E(\mathbf{y}_i|\mathbf{u}_i) = g(\mathbf{x}_i, \boldsymbol{\beta}, \mathbf{u}_i)$ and the conditional variance $Var(\mathbf{y}_i|\mathbf{u}_i) = diag(\sigma_{i1}^2, \cdots, \sigma_{in_i}^2)$. Modelling the variances allows for more efficient estimation of the mean parameters and better understanding of the within-individual variabilities.

2.2 Robust model

In the variance model (2), if we assume that $\exp(a_i) \sim k/\chi_k^2$, i.e., the inverse χ^2 distribution, which is a special case of a inverse gamma distribution, then the random errors e_{ij} for the within-individual repeated measurement follow a $t(k)$-distribution with degrees of freedom k (Lang et al. 1989). The $t(k)$-distribution has heavier tails than the normal distribution, and thus it incorporates possible outliers in the within-individual repeated measurements. The smaller the degrees of freedom k, the heavier the tails. The $t(k)$-distribution reduces to the standard normal distribution when $k \to \infty$. Therefore, in this case models (1) and (2) may be used for robust inference.

Similarly, we may allow the variances for the random effects for the mean model $\mathbf{u}_i$ to be individual-specific, i.e., $\mathbf{u}_i \sim N(0, D_i)$, and model the covariance matrix D_i in a similar way as σ_{ij}^2. While the approaches are conceptually similar, for simplicity we focus on the variance model (2) and assume $D_i = D$ in this paper.

2.3 Covariate measurement errors

The covariate z_{ij} in the variance model may be measured with errors, such as CD4 cell count. The covariate effect on the variance, if any, may be masked if the measurement errors are not addressed. To address the measurement errors, we may view the repeated measurements of the covariate as its "replicates" in order to estimate the magnitude of the measurement

errors and model the covariate process using an empirical LME model.

Let z_{ij}^* be the unobserved true covariate value for individual i at time t_{ij} and let z_{ij} be the corresponding observed value with measurement error. We may consider the following mixed effects model as a classic measurement error model

$$z_{ij} \equiv z_{ij}^* + \epsilon_{ij} = q(t_{ij}, \boldsymbol{\gamma}, \mathbf{b}_i) + \epsilon_{ij}, \tag{3}$$

where $q(\cdot)$ is a known linear function or a unknown smooth function, $\boldsymbol{\gamma}$ contains fixed parameters, $\mathbf{b}_i$ contains random effects, and ϵ_{ij} is measurement error. We assume that $b_i \sim N(0, B)$, and independently ϵ_{ij} $i.i.d.$ $\sim N(0, \omega^2)$.For example, we may model the CD4 process using a quadratic mixed effects model $z_{ij} = \gamma_0 + \gamma_1 t_{ij} + \gamma_2 t_{ij}^2 + b_i + \epsilon_{ij}$. Alternatively, for more complex CD4 trajectories, we may choose $q(t_{ij}, \boldsymbol{\gamma}, \mathbf{b}_i)$ to be a nonparametric smooth function, then approximate it using a basis-based approach, which leads to an approximate LME model (Wu, 2009). In either case, we may consider an LME model for the measurement error model (3).

3 Approximate Likelihood Inference

Let $\boldsymbol{\theta} = (\boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\xi})$ denote the collection of all fixed parameters in the models, where $(\boldsymbol{\beta}, \boldsymbol{\gamma})$ contains fixed mean parameters, and $\boldsymbol{\xi} = (\boldsymbol{\alpha}, D, \eta, \omega^2)$ contains fixed variance (dispersion) parameters. Let $\mathbf{v}_i = (\mathbf{u}_i, a_i, \mathbf{b}_i)$ denote the collection of all random effects, and let $\mathbf{v} = (\mathbf{v}_1, \cdots, \mathbf{v}_n)$. Standard likelihood inference for $\boldsymbol{\theta}$ is based on the following (marginal) likelihood function, which integrates out the unobservable random effects

$$L(\boldsymbol{\theta}) = \prod_{i=1}^{n} \int \int \int f(\mathbf{y}_i |\, \mathbf{x}_i, \mathbf{z}_i, \boldsymbol{\beta}, D, \boldsymbol{\sigma}_i^2, \mathbf{u}_i) f(\boldsymbol{\sigma}_i^2 | \boldsymbol{\alpha}, a_i) f(\mathbf{z}_i | \boldsymbol{\gamma}, \mathbf{b}_i) f(\mathbf{u}_i, \mathbf{b}_i, a_i | \boldsymbol{\theta})\, d\mathbf{u}_i\, da_i\, d\mathbf{b}_i,$$

where $f(\cdot)$ denotes a generic density function. The above likelihood involves a high-dimensional and intractable integration. Monte Carlo EM algorithms or Gaussian quadrature numerical integration methods can be computationally very intensive. Thus, in the following, we discuss a computationally more efficient approximate method for approximate likelihood inference, using the h-likelihood (Lee et al. 2017) or conditional second-order generalized estimating

equations (CGEE2, Vonesh et al. 2002).

The h-loglikelihood (Lee et al. 2017) can be written as

$$h(\boldsymbol{\theta}, \mathbf{v}) = \sum_{i=1}^{n} \left\{ \log f(\mathbf{y}_i | \, \mathbf{x}_i, \mathbf{z}_i, \boldsymbol{\beta}, D, \boldsymbol{\sigma}_i^2, \mathbf{u}_i) + \log f(\boldsymbol{\sigma}_i^2 | \boldsymbol{\alpha}, a_i) + \log f(\mathbf{z}_i | \boldsymbol{\gamma}, \mathbf{b}_i) + \log f(\mathbf{u}_i, \mathbf{b}_i, a_i | \boldsymbol{\theta}) \right\}$$

where the random effects $\mathbf{v}_i = (\mathbf{u}_i, \mathbf{b}_i, a_i)$ are treated as "parameters". Given parameter $\boldsymbol{\theta}$, estimates of the random effects $\mathbf{v}$ can be obtained by maximizing the h-loglikelihood $h(\boldsymbol{\theta}, \mathbf{v})$ with respect to $\mathbf{v}$, denoted by $\hat{\mathbf{v}}$, i.e., $\hat{\mathbf{v}}$ solves $\partial h(\boldsymbol{\theta}, \mathbf{v})/\partial \mathbf{v} = 0$. Then, we can obtain estimates of the parameters $\boldsymbol{\theta}$ by maximizing $h(\boldsymbol{\theta}, \hat{\mathbf{v}})$. The process iterates until convergence. For binary or discrete response $\mathbf{y}_i$, Lee et al. (2017) recommends adding an adjusted term to reduce bias and obtain estimates of the parameters $\boldsymbol{\theta}$ by maximizing the following "adjusted profile likelihood"

$$\log L(\boldsymbol{\theta}) \approx p_{\mathbf{v}}(h) \equiv \left[h(\boldsymbol{\theta}, \mathbf{v}) - \frac{1}{2} \log det\{H(h; \mathbf{v})/(2\pi)\} \right] \Bigg|_{\mathbf{v}=\hat{\mathbf{v}}},$$

where the second term is an adjusted term, and $H(h; \mathbf{v}) = -\partial^2 h/\partial \mathbf{v}^2$ and $\hat{\mathbf{v}}$ solves $\partial h/\partial \mathbf{v} = 0$. Lee and Nelder (2001) show that the h-likelihood estimate is the first-order Laplace approximation to the marginal log-likelihood $\log L(\boldsymbol{\theta})$.

Vonesh et al. (2002) proposed the CGEE2 method which also maximizes $h(\boldsymbol{\theta}, \mathbf{v})$ but without the adjusted term. That is, the CGEE2 estimate may be viewed as the h-likelihood estimate without the adjusted term. They showed that their CGEE2 estimate is asymptotically equivalent to the h-likelihood estimate, the linearization estimate of Lindstrom and Bates (1990) for NLME models, and the penalized quasi-likelihood (PQL) estimate of Breslow and Clayton (1993), when both the sample size n and the numbers of repeated measurements $\min_i\{n_i\}$ go to ∞. Moreover, they showed that the CGEE2 estimates are consistent and asymptotically normally distributed. Following their ideas, we propose the following iterative algorithm to obtain approximate MLE of the parameter $\boldsymbol{\theta} = (\boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\xi})$. Starting with some initial values $(\boldsymbol{\theta}^{(0)}, \mathbf{v}^{(0)})$, at iterative $k, k = 0, 1, 2, \cdots$,

- Step 1: Solve equation $\partial h(\boldsymbol{\theta}^{(k)}, \mathbf{v})/\partial \mathbf{v} = 0$ and obtain updated estimate of the random effects $\hat{\mathbf{v}}^{(k+1)}$;

- Step 2: Solve equation $\partial h(\boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\xi}^{(k)}; \mathbf{v}^{(k+1)})/\partial(\boldsymbol{\beta}, \boldsymbol{\gamma}) = 0$ and obtain updated estimates of the mean parameters $(\boldsymbol{\beta}^{(k+1)}, \boldsymbol{\gamma}^{(k+1)})$;
- Step 3: Solve equation $\partial h(\boldsymbol{\beta}^{(k+1)}, \boldsymbol{\gamma}^{(k+1)}, \boldsymbol{\xi}; \mathbf{v}^{(k+1)})/\partial\boldsymbol{\xi} = 0$ and obtain updated estimates of the variance-covariance parameters $\boldsymbol{\xi}^{(k+1)}$.

Iterate the above steps until convergence, we obtain approximate MLE $\hat{\boldsymbol{\theta}}$. Standard errors of the estimates may be obtained from the diagonal elements (square roots) of the inverse of the Fisher information matrix.

4 A Real Data Example

Analyze a real HIV/AIDS dataset using three methods: the Linstrom and Bates (1990) method implemented in R package *nlme* without modeling the variances, the two-step method, and the proposed method. The results should show efficiency gain (smaller standard errors) of the estimates $\hat{\boldsymbol{\beta}}$ which allow us to compare different treatments more effectively. Estimates of $\boldsymbol{\alpha}$ also allow us to see if CD4 partially explains the within-individual variance σ_{ij}^2.

5 Simulation Study

In the simulation study, we will evaluate the proposed method and compare it with the Linstrom and Bates (1990) method implemented in R package *nlme* without modeling the variances and the two-step method. The simulation will be designed in the following ways: i) compare estimates, standard errors, MSEs, and coverage probabilities; ii) bootstrap standard errors (either parametric or nonparametric bootstrap); iii) robustness of the proposed method (with inverse-gamma distribution for a_i) for simulated data with 5% outliers (adding a large number, say 10, to 5% simulated data); iv) different sample size n (say $n = 100, 200$) and different values of n_i (say, 15, 30).

Compare estimates, standard errors, MSEs, and coverage probabilities

Data are simulated for $n = 100$ subjects, with $n_i = 15$ repeated measurements for each subject. The entire study period is set to be $[0, 1]$, and the 15 measuring time points are

equally allocated within the time interval. The true parameter values are chosen based on the estimates from the empirical analysis in Section 4. Specifically,

$$\begin{aligned}
\beta &= (P_1, P_2, \lambda)^T = (2.5, 3.0, 7.5)^T, \\
\alpha &= (\alpha_0, \alpha_1)^T = (-8, 1.5)^T, \\
\gamma &= (\gamma_0, \gamma_1, \gamma_2)^T = (5.2, 1.6, -1.2)^T, \\
(d_1, d_3) &= (0.4, 1.0), \quad \sigma_a = 0.7, \quad \sigma_b = 0.4, \quad \xi = 0.2.
\end{aligned}$$

The structure of the variance-covariance matrix for the random effects $v_i = (u_i, a_i, b_i)$ varies depending on different assumptions. For now, we assume the random effects are correlated within the same model and independent across different models. The starting values for the inference are set to be the true values. The number of simulation repetition is set to be 100.

Table 1 reports "Est", "rBias", "rMSE", "SE.em", "SE", and "Coverage" of the parameters in models. For each parameter, "Est" is the average of the 100 estimates from the 100 simulation runs, "rBias" and "rMSE" are the relative bias and mean square error of the 100 parameter estimates, which are defined as follows:

$$\begin{aligned}
\text{rBias (\%) of } \hat{\beta} &= \left| \left(\frac{1}{M} \sum_{m=1}^{M} \hat{\beta}^{(m)} - \beta \right) / \beta \right| \times 100\%, \\
\text{rMSE (\%) of } \hat{\beta} &= \left| \frac{1}{M} \sum_{m=1}^{M} \left(\hat{\beta}^{(m)} - \beta \right)^2 / \beta \right| \times 100\%,
\end{aligned}$$

where $\hat{\beta}^{(m)}$ is the estimate of β in the m-th simulation run and M is the total number of simulation repetitions, "SE.em" is the empirical standard deviation of the 100 estimates, "SE" is the square root of the average of the 100 variance estimates, and "Coverage" is the empirical coverage of the true parameter value among the 100 95% confidence intervals from the simulation runs.

In addition, **parametric** bootstrapping method was used to adjust the SE for the parameter estimates. We conducted 100 bootstrapping runs for each of 100 simulation runs. "SE.BT" is the average of bootstrapping SE from the 100 simulation runs and " Coverage.bt" is the corresponding coverage rate. The bootstrapping SE was calculated as the empirical

SE of the estimates from the 100 bootstrapping runs.

Table 1: Simulation results

Fixed effects in	Robust NLME			Robust NLME variance component		LME		
True parameter	$P_1 = 2.5$	$P_2 = 3.0$	$\lambda = 7.5$	$\alpha_0 = -8$	$\alpha_1 = 1.5$	$\gamma_0 = 5.2$	$\gamma_1 = 1.6$	$\gamma_2 = -1.2$
`nlme` package:								
Est	2.527	2.996	8.054					
rBias(%)	1.076	0.135	7.393					
rMSE(%)	0.235	0.441	14.313					
SE.em	0.072	0.116	0.880					
SE	0.067	0.148	0.862					
Coverage	0.920	0.990	0.910					
Two-step method:								
Est	2.509	2.994	7.699	-8.392	1.547	5.206	1.598	-1.198
rBias(%)	0.375	0.190	2.655	4.894	3.106	0.120	0.144	0.167
rMSE(%)	0.102	0.222	3.647	8.426	1.252	0.033	0.207	0.258
SE.em	0.050	0.082	0.486	0.725	0.130	0.041	0.058	0.056
SE	0.052	0.069	0.433	0.492	0.087	0.042	0.063	0.061
Coverage	0.960	0.920	0.930	0.760	0.800	0.980	0.970	0.960
Joint likelihood method:								
Est	2.511	2.995	7.759	-8.069	1.489	5.205	1.603	-1.202
rBias(%)	0.428	0.174	3.455	0.866	0.729	0.091	0.210	0.149
rMSE(%)	0.106	0.222	4.027	7.479	1.267	0.033	0.193	0.237
SE.em	0.051	0.082	0.487	0.774	0.138	0.041	0.056	0.054
SE	0.052	0.069	0.441	0.486	0.086	0.042	0.060	0.058
Coverage	0.950	0.930	0.920	0.800	0.810	0.980	0.970	0.960
SE.BT	0.057	0.075	0.495	0.690	0.122	0.043	0.061	0.059
Coverage.bt	0.970	0.960	0.950	0.930	0.920	0.970	0.960	0.970

Different sample size: $n = 200$

Table 2: Simulation results

Fixed effects in	Robust NLME			Robust NLME variance component		LME		
True parameter	$P_1 = 2.5$	$P_2 = 3.0$	$\lambda = 7.5$	$\alpha_0 = -8$	$\alpha_1 = 1.5$	$\gamma_0 = 5.2$	$\gamma_1 = 1.6$	$\gamma_2 = -1.2$
`nlme` package:								
Est	2.507	2.976	7.425					
rBias(%)	0.289	0.785	1.000					
rMSE(%)	0.110	0.223	5.060					
SE.em	0.052	0.079	0.615					
SE	0.048	0.104	0.561					
Coverage	0.920	0.980	0.920					
Two-step method:								
Est	2.503	2.991	7.526	-8.331	1.536	5.199	1.605	-1.203
rBias(%)	0.108	0.295	0.353	4.133	2.407	0.013	0.309	0.281
rMSE(%)	0.075	0.108	2.047	3.833	0.500	0.015	0.114	0.140
SE.em	0.043	0.056	0.393	0.446	0.079	0.028	0.043	0.041
SE	0.036	0.048	0.295	0.347	0.061	0.030	0.045	0.043
Coverage	0.920	0.910	0.860	0.750	0.820	0.950	0.960	0.970
Joint likelihood method:								
Est	2.504	2.992	7.582	-8.003	1.478	5.198	1.611	-1.207
rBias(%)	0.145	0.257	1.094	0.043	1.493	0.043	0.670	0.604
rMSE(%)	0.076	0.108	2.138	3.667	0.640	0.015	0.115	0.135
SE.em	0.044	0.057	0.394	0.544	0.096	0.028	0.042	0.040
SE	0.036	0.048	0.299	0.343	0.060	0.030	0.042	0.041
Coverage	0.930	0.910	0.860	0.780	0.780	0.960	0.940	0.950
SE.BT	0.040	0.052	0.340	0.470	0.083	0.030	0.043	0.041
Coverage.bt	0.960	0.930	0.890	0.940	0.930	0.960	0.940	0.970

Different number of repeated measurements: $n_i = 30$

Table 3: Simulation results

Fixed effects in	Robust NLME			Robust NLME variance component		LME		
True parameter	$P_1 = 2.5$	$P_2 = 3.0$	$\lambda = 7.5$	$\alpha_0 = -8$	$\alpha_1 = 1.5$	$\gamma_0 = 5.2$	$\gamma_1 = 1.6$	$\gamma_2 = -1.2$
`nlme` package:								
Est	2.511	3.009	7.737					
rBias(%)	0.438	0.301	3.166					
rMSE(%)	0.160	0.304	7.314					
SE.em	0.063	0.096	0.705					
SE	0.056	0.126	0.632					
Coverage	0.920	0.980	0.910					
Two-step method:								
Est	2.507	3.007	7.651	-8.199	1.524	5.204	1.596	-1.195
rBias(%)	0.272	0.221	2.017	2.488	1.620	0.074	0.249	0.431
rMSE(%)	0.125	0.110	2.171	3.662	0.557	0.030	0.155	0.191
SE.em	0.056	0.057	0.376	0.506	0.089	0.039	0.050	0.048
SE	0.045	0.060	0.338	0.354	0.062	0.041	0.047	0.046
Coverage	0.850	0.960	0.910	0.770	0.840	0.940	0.960	0.970
Joint likelihood method:								
Est	2.508	3.008	7.710	-8.019	1.492	5.203	1.599	-1.197
rBias(%)	0.328	0.280	2.797	0.242	0.521	0.057	0.034	0.225
rMSE(%)	0.125	0.114	2.660	3.990	0.665	0.030	0.150	0.184
SE.em	0.056	0.058	0.396	0.568	0.100	0.039	0.049	0.047
SE	0.045	0.060	0.343	0.352	0.061	0.041	0.046	0.044
Coverage	0.850	0.950	0.880	0.770	0.770	0.940	0.950	0.940
SE.BT	0.048	0.063	0.385	0.552	0.097	0.042	0.046	0.044
Coverage.bt	0.890	0.950	0.910	0.940	0.940	0.960	0.960	0.940

6 Discussion

References

German, C.A., Sinsheimer, J.S., Zhou, J., and Zhou, H. (2022). WiSER: Robust and scalable estimation and inference of within-subject variances from intensive longitudinal data. *Biometrics*, in press.

Hedeker, D., Mermelstein, R.J., and Demirtas, H. (2008).
An Application of a Mixed-Effects Location Scale Model for Analysis of Ecological Momentary Assessment (EMA) Data. *Biometrics 64*, 627-634.

Lee, Y., Nelder, J.A., and Pawitan, Y. (2017).
Generalized Linear Models with Random Effects: Unified Analysis via H-likelihood, Second edition. London: Chapman & Hall/CRC.

Lin, X., Raz, J., and Harlow, S.D. (1997).
Linear Mixed Models with Heterogeneous Within-Cluster Variances. *Biometrics 53*, 910-923.

Lindstrom, M.J. and Bates, D.M. (1990). Nonlinear mixed effects models for repeated measures data. *Biometrics* **46**, 673-687.

Noh, M., Wu, L., and Lee, Y. (2012).
Hierarchical Likelihood Methods for Nonlinear and Generalized Linear Mixed Models with Missing Data and Measurement Errors in Covariates, *Journal of Multivariate Analysis*, 109, 42-51.

Pourahmadi, M. (1999). Joint mean-covariance models with applications to longitudinal data:
Unconstrained parameterisation. *Biometrika 86*, 677-690.

Vonesh, E.F., Wang, H., Nie, L., and Majumdar, D. (2002).
Conditional second-order generalized estimating equations for generalized linear and nonlinear mixed-effects models.
Journal of the American Statistical Association **97**, 271-283.