

# Causal Inference

## Seminar #1

Juraj Medzihorsky

SGIA & RMC  
Durham University

16 January 2024

# Matching

## Selection on Observables

- Suppose

$$\begin{aligned}D_i &\longrightarrow Y_i \\Y_i(0), Y_i(1) &\not\perp\!\!\!\perp D_i\end{aligned}$$

that is, there is selection into treatment, and also

$$\begin{aligned}\mathbf{X}_i &\longrightarrow D_i \\\mathbf{X}_i &\longrightarrow Y_i\end{aligned}$$

- Intuitively: If we are able to account for all of the  $\mathbf{X}$  variables that affect both selection into treatment,  $D$ , **and** variation in the outcome,  $Y$ , then we can claim independence between  $D$  and potential outcomes.

## Selection on Observables

- Suppose

$$\begin{aligned}D_i &\longrightarrow Y_i \\Y_i(0), Y_i(1) &\not\perp\!\!\!\perp D_i\end{aligned}$$

that is, there is selection into treatment, and also

$$\begin{aligned}\mathbf{X}_i &\longrightarrow D_i \\\mathbf{X}_i &\longrightarrow Y_i\end{aligned}$$

- Intuitively: If we are able to account for all of the  $\mathbf{X}$  variables that affect both selection into treatment,  $D$ , **and** variation in the outcome,  $Y$ , then we can claim independence between  $D$  and potential outcomes.
- The big problem: We have to be able to observe and measure all of the  $X$  variables that affect selection into treatment and variation in outcomes.

## Selection on Observables

- Suppose

$$\begin{aligned}D_i &\longrightarrow Y_i \\Y_i(0), Y_i(1) &\not\perp\!\!\!\perp D_i\end{aligned}$$

that is, there is selection into treatment, and also

$$\begin{aligned}\mathbf{X}_i &\longrightarrow D_i \\\mathbf{X}_i &\longrightarrow Y_i\end{aligned}$$

- Intuitively: If we are able to account for all of the  $\mathbf{X}$  variables that affect both selection into treatment,  $D$ , **and** variation in the outcome,  $Y$ , then we can claim independence between  $D$  and potential outcomes.
- The big problem: We have to be able to observe and measure all of the  $X$  variables that affect selection into treatment and variation in outcomes.

## Matching

- More precisely *matching on observables*
- Suppose
  - Binary treatment, i.e., only two options: treatment and control
  - We know and observe all potential confounders
- Then we can use the untreated observations to construct a control group that will be as similar as possible to the treated observations
- This helps in estimating the average treatment effect *on the treated* (ATT) which may not be the same as the average treatment effect (ATE)
- Many options: different matching algorithms

# Identification under Selection on Observables

## Identification Assumptions

1. *Selection on Observables*

# Identification under Selection on Observables

## Identification Assumptions

### 1. *Selection on Observables*

- ATE Version:  $(Y(1), Y(0)) \perp\!\!\!\perp D | X$

# Identification under Selection on Observables

## Identification Assumptions

### 1. *Selection on Observables*

- ATE Version:  $(Y(1), Y(0)) \perp\!\!\!\perp D | X$ 
  - ▶ *There exists a set of observable covariates,  $X$ , such that after controlling for  $X$ , potential outcomes are independent of treatment status*

# Identification under Selection on Observables

## Identification Assumptions

### 1. *Selection on Observables*

- ATE Version:  $(Y(1), Y(0)) \perp\!\!\!\perp D | X$ 
  - ▶ There exists a set of observable covariates,  $X$ , such that after controlling for  $X$ , potential outcomes are independent of treatment status
- ATT Version:  $Y(0) \perp\!\!\!\perp D | X$

# Identification under Selection on Observables

## Identification Assumptions

### 1. Selection on Observables

- ATE Version:  $(Y(1), Y(0)) \perp\!\!\!\perp D | X$ 
  - ▶ There exists a set of observable covariates,  $X$ , such that after controlling for  $X$ , potential outcomes are independent of treatment status
- ATT Version:  $Y(0) \perp\!\!\!\perp D | X$ 
  - ▶ There exists a set of observable covariates,  $X$ , such that after controlling for  $X$ , counterfactual outcomes for treated units and observed outcomes for untreated units are independent of treatment status

# Identification under Selection on Observables

## Identification Assumptions

### 1. Selection on Observables

- ATE Version:  $(Y(1), Y(0)) \perp\!\!\!\perp D | X$ 
  - ▶ There exists a set of observable covariates,  $X$ , such that after controlling for  $X$ , potential outcomes are independent of treatment status
- ATT Version:  $Y(0) \perp\!\!\!\perp D | X$ 
  - ▶ There exists a set of observable covariates,  $X$ , such that after controlling for  $X$ , counterfactual outcomes for treated units and observed outcomes for untreated units are independent of treatment status

### 2. Common Support

# Identification under Selection on Observables

## Identification Assumptions

### 1. Selection on Observables

- ATE Version:  $(Y(1), Y(0)) \perp\!\!\!\perp D | X$ 
  - ▶ There exists a set of observable covariates,  $X$ , such that after controlling for  $X$ , potential outcomes are independent of treatment status
- ATT Version:  $Y(0) \perp\!\!\!\perp D | X$ 
  - ▶ There exists a set of observable covariates,  $X$ , such that after controlling for  $X$ , counterfactual outcomes for treated units and observed outcomes for untreated units are independent of treatment status

### 2. Common Support

- ATE Version:  $0 < \Pr(D = 1 | X = x) < 1$

# Identification under Selection on Observables

## Identification Assumptions

### 1. Selection on Observables

- ATE Version:  $(Y(1), Y(0)) \perp\!\!\!\perp D | X$ 
  - ▶ There exists a set of observable covariates,  $X$ , such that after controlling for  $X$ , potential outcomes are independent of treatment status
- ATT Version:  $Y(0) \perp\!\!\!\perp D | X$ 
  - ▶ There exists a set of observable covariates,  $X$ , such that after controlling for  $X$ , counterfactual outcomes for treated units and observed outcomes for untreated units are independent of treatment status

### 2. Common Support

- ATE Version:  $0 < \Pr(D = 1 | X = x) < 1$ 
  - ▶ For each value of  $X$ , there is a positive probability of observing both treated and untreated units

# Identification under Selection on Observables

## Identification Assumptions

### 1. Selection on Observables

- ATE Version:  $(Y(1), Y(0)) \perp\!\!\!\perp D | X$ 
  - ▶ There exists a set of observable covariates,  $X$ , such that after controlling for  $X$ , potential outcomes are independent of treatment status
- ATT Version:  $Y(0) \perp\!\!\!\perp D | X$ 
  - ▶ There exists a set of observable covariates,  $X$ , such that after controlling for  $X$ , counterfactual outcomes for treated units and observed outcomes for untreated units are independent of treatment status

### 2. Common Support

- ATE Version:  $0 < \Pr(D = 1 | X = x) < 1$ 
  - ▶ For each value of  $X$ , there is a positive probability of observing both treated and untreated units
- ATT Version:  $\Pr(D = 1 | X = x) < 1$ , with  $\Pr(D = 1) > 0$

# Identification under Selection on Observables

## Identification Assumptions

### 1. Selection on Observables

- ATE Version:  $(Y(1), Y(0)) \perp\!\!\!\perp D | X$ 
  - ▶ There exists a set of observable covariates,  $X$ , such that after controlling for  $X$ , potential outcomes are independent of treatment status
- ATT Version:  $Y(0) \perp\!\!\!\perp D | X$ 
  - ▶ There exists a set of observable covariates,  $X$ , such that after controlling for  $X$ , counterfactual outcomes for treated units and observed outcomes for untreated units are independent of treatment status

### 2. Common Support

- ATE Version:  $0 < \Pr(D = 1 | X = x) < 1$ 
  - ▶ For each value of  $X$ , there is a positive probability of observing both treated and untreated units
- ATT Version:  $\Pr(D = 1 | X = x) < 1$ , with  $\Pr(D = 1) > 0$ 
  - ▶ For each value of  $X$  observed for a treated unit, there should exist an untreated unit with the same value of  $X$

## Matching

When  $X$  is continuous we can estimate  $\alpha_{ATT}$  by “imputing” the missing potential outcome of each treated unit using the observed outcome from the “closest” control unit:

$$\hat{\alpha}_{ATT} = \frac{1}{N_1} \sum_{D_i=1} (Y_i - Y_{j(i)})$$

where  $Y_{j(i)}$  is the outcome of an untreated observation such that  $X_{j(i)}$  is the closest value to  $X_i$  among the untreated observations.

## Matching

When  $X$  is continuous we can estimate  $\alpha_{ATT}$  by “imputing” the missing potential outcome of each treated unit using the observed outcome from the “closest” control unit:

$$\hat{\alpha}_{ATT} = \frac{1}{N_1} \sum_{D_i=1} (Y_i - Y_{j(i)})$$


where  $Y_{j(i)}$  is the outcome of an untreated observation such that  $X_{j(i)}$  is the closest value to  $X_i$  among the untreated observations.

We can also use the average for  $M$  closest matches:


$$\hat{\alpha}_{ATT} = \frac{1}{N_1} \sum_{D_i=1} \left\{ Y_i - \left( \frac{1}{M} \sum_{m=1}^M Y_{j_m(i)} \right) \right\}$$

Works well when we can find good matches for each treated unit.


## Exact Matching: 2D Example



## Exact Matching: 2D Example



## Coarsened Exact Matching: Example



## Matching: Distance Metric

“Closeness” is often defined by a distance metric that projects the distance between the multivariate covariate vectors  $X_i = (X_{i1}, X_{i2}, \dots, X_{ik})'$  and  $X_j = (X_{j1}, X_{j2}, \dots, X_{jk})'$  onto a univariate scale

## Matching: Distance Metric

“Closeness” is often defined by a distance metric that projects the distance between the multivariate covariate vectors  $X_i = (X_{i1}, X_{i2}, \dots, X_{ik})'$  and  $X_j = (X_{j1}, X_{j2}, \dots, X_{jk})'$  onto a univariate scale

A commonly used distance is the Mahalanobis distance:

$$D_M(X_i, X_j) = \sqrt{(X_i - X_j)' S^{-1} (X_i - X_j)} ,$$

where  $S$  is the sample variance-covariance-matrix.

## Matching: Distance Metric

“Closeness” is often defined by a distance metric that projects the distance between the multivariate covariate vectors  $X_i = (X_{i1}, X_{i2}, \dots, X_{ik})'$  and  $X_j = (X_{j1}, X_{j2}, \dots, X_{jk})'$  onto a univariate scale


A commonly used distance is the Mahalanobis distance:

$$D_M(X_i, X_j) = \sqrt{(X_i - X_j)' S^{-1} (X_i - X_j)} ,$$

where  $S$  is the sample variance-covariance-matrix.


Propensity score is defined as the selection probability conditional on the confounding variables:  $\pi(X) = P(D = 1|X)$ .

## Propensity Score Matching: Example




Source: Gary King

## Propensity Score Matching: Example




## Propensity Score Matching: Example




Source: Gary King

## Propensity Score Matching: Example




Source: Gary King

## Propensity Score Matching: Example



Source: Gary King

## Propensity Score Matching: Example



## Practical Steps in Matching

1. Like all data analysis exercises, explore your raw data before making a lot of assumptions, transformations, etc.

## Practical Steps in Matching

1. Like all data analysis exercises, explore your raw data before making a lot of assumptions, transformations, etc.

In matching exercises, prior to constructing a matched sample, this can involve...

- Simple difference-of-means tests on the outcome variable
- Regression-adjusted difference-of-means tests on the outcome variable using pre-treatment covariates

## Practical Steps in Matching

### 2. Construct the matched sample

- Whether using distance-matching or propensity scores, try multiple specifications, using different choices for covariates, numbers of control units for each treatment unit, matching with and without replacement, etc.
- Check for balance on pre-treatment covariates (e.g.,  $t$ -tests for balance on pre-treatment covariates, graphical overlays of covariate densities between treatment groups)
- For propensity-score matching, explore the distribution of the propensity score across treatment groups

## Practical Steps in Matching

### 2. Construct the matched sample

- Whether using distance-matching or propensity scores, try multiple specifications, using different choices for covariates, numbers of control units for each treatment unit, matching with and without replacement, etc.
- Check for balance on pre-treatment covariates (e.g.,  $t$ -tests for balance on pre-treatment covariates, graphical overlays of covariate densities between treatment groups)
- For propensity-score matching, explore the distribution of the propensity score across treatment groups
- Choose your method based on balance, not treatment effects!  
For best practices, do not even look at treatment effects before the matching method is chosen

## Practical Steps in Matching

### 3. Estimate treatment effects

- Simple difference-of-means tests between experimental groups
- Regression-adjusted difference of means tests using pre-treatment variables as controls
- Same types of graphical techniques used for difference-of-means and regressions
- Comparisons to unmatched analyses
- If possible, comparisons to an experimental benchmark


## Practical Steps in Matching

4. Evaluate the selection-on-observables assumption with sensitivity analysis
  - Rosenbaum bounds are probably the most widely used test
  - Usually will take the form of asking: How bad will our inferences be if we left out an important covariate?

?


Thank you!

## Coarsened Exact Matching: Example




Source: Gary King

## Coarsened Exact Matching: Example




## Coarsened Exact Matching: Example




Source: Gary King


## Coarsened Exact Matching: Example



## Coarsened Exact Matching: Example



## Mahalanobis Distance Matching: Example



Education (years)

Source: Gary King

## Mahalanobis Distance Matching: Example

