




ARMA Models

Katalin Varga (Ph.D.)

BUTE

2016 Autumn







What is a time series?

Notation: $\{x_1, x_2, \dots, x_T\}$ or $\{x_t\}$, $t = 1, 2, \dots, T$


Example 1.: $y_t = \beta x_t + \epsilon_t$, $E(\epsilon_t | x_t) = 0$

Example 2.: $x_t = \epsilon_t$, $\epsilon_t \sim N(0, \sigma^2)$





Real GDP Growth and Unemployment Rate of the USA





Linear Time Series Models

- 1. Building block:

White noise: $x_t = \epsilon_t$, $\epsilon_t \sim i.i.d. N(0, \sigma^2)$

1. $E(\epsilon_t) = E(\epsilon_t | \epsilon_{t-1}, \epsilon_{t-2}, \dots) = E(\epsilon_t | \text{all information at } t-1) = 0$
2. $E(\epsilon_t \epsilon_{t-j}) = cov(\epsilon_t \epsilon_{t-j}) = 0$
3. $var(\epsilon_t) = var(\epsilon_t | \epsilon_{t-1}, \epsilon_{t-2}, \dots) = var(\epsilon_t | \text{all information at } t-1) = \sigma^2$

- 2. ARMA models:

$$AR(1): x_t = \varphi x_{t-1} + \epsilon_t$$

$$AR(p): x_t = \varphi_1 x_{t-1} + \varphi_2 x_{t-2} + \dots + \varphi_p x_{t-p} + \epsilon_t$$

$$MA(1): x_t = \epsilon_t + \theta \epsilon_{t-1}$$

$$MA(q): x_t = \epsilon_t + \theta_1 \epsilon_{t-1} + \dots + \theta_q \epsilon_{t-q}$$

$$ARMA(p,q): x_t = \varphi_1 x_{t-1} + \dots + \epsilon_t + \theta_1 \epsilon_{t-1} + \dots$$



Lag Operators and Lag Polynomials

- Lag operator moves the index back one time unit:

$$Lx_t = x_{t-1} \rightarrow L^2x_t = x_{t-2} \rightarrow L^{-j}x_t = x_{t+j}$$

- So we can define lag polynomials eg.:

$$a(L)x_t = (a_0L^0 + a_1L^1 + a_2L^2)x_t = a_0x_t + a_1x_{t-1} + a_2x_{t-2}$$

- Using this notation we can rewrite ARMA models:

$$\text{AR: } a(L)x_t = \epsilon_t$$

$$\text{MA: } x_t = b(L)\epsilon_t$$

$$\text{ARMA: } a(L)x_t = b(L)\epsilon_t$$



AR-MA Substitution

$$\text{AR}(1): x_t = \varphi x_{t-1} + \epsilon_t$$



$$x_t = \varphi x_{t-1} + \epsilon_t = \varphi(\varphi x_{t-2} + \epsilon_{t-1}) + \epsilon_t = \varphi^2 x_{t-2} + \varphi \epsilon_{t-1} + \epsilon_t$$

$$x_t = \varphi^k x_{t-k} + \varphi^{k-1} \epsilon_{t-k+1} + \dots + \varphi^2 x_{t-2} + \varphi \epsilon_{t-1} + \epsilon_t$$

If $|\varphi| < 1$ then $\lim_{k \rightarrow \infty} \varphi^k x_{t-k} = 0$, so $x_t = \sum_{j=0}^{\infty} \varphi^j \epsilon_{t-j}$.

It means that AR(1) can be expressed as MA(∞).



The Autocovariance Function

- Definition: The autocovariance of the series x_t is defined as

$$\gamma_j = cov(x_t, x_{t-j})$$

We assume that $E(x_t) = 0$ for all our models, since we are specializing to ARMA models without constant terms.

$$\gamma_j = E(x_t, x_{t-j})$$

Note that $\gamma_0 = var(x_t)$, the autocorrelation function is

$$\rho_j = \gamma_j / var(x_t) = \gamma_j / \gamma_0$$



Autocovariance of ARMA Processes

- White Noise: since $x_t = \epsilon_t$, $\epsilon_t \sim i.i.d. N(0, \sigma^2)$, it is obvious

$$\gamma_0 = \sigma_\epsilon^2, \gamma_j = 0 \text{ for } j \neq 0$$
$$\rho_0 = 1, \rho_j = 0 \text{ for } j \neq 0$$

- MA(1): $x_t = \epsilon_t + \theta\epsilon_{t-1}$, the autocovariance

$$\gamma_0 = \text{var}(x_t) = \text{var}(\epsilon_t + \theta\epsilon_{t-1}) = (1 + \theta^2)\sigma_\epsilon^2$$
$$\gamma_1 = E(x_t x_{t-1}) = \text{var}(\epsilon_t + \theta\epsilon_{t-1})(\epsilon_{t-1} + \theta\epsilon_{t-2}) = \theta\sigma_\epsilon^2$$
$$\gamma_2 = \text{var}(x_t x_{t-2}) = \text{var}(\epsilon_t + \theta\epsilon_{t-1})(\epsilon_{t-2} + \theta\epsilon_{t-3}) = 0$$
$$\gamma_3 = 0$$

$$\rho_1 = \theta / (1 + \theta^2), \quad \rho_{2,\dots} = 0$$



Autocovariance of ARMA Processes

- AR(1): $x_t = \varphi x_{t-1} + \epsilon_t$ the autocovariance

$$(1 - \varphi L)x_t = \epsilon_t \rightarrow x_t = (1 - \varphi L)^{-1}\epsilon_t = \sum_{j=0}^{\infty} \varphi^j \epsilon_{t-j}$$

$$\gamma_0 = \left(\sum_{j=0}^{\infty} \varphi^{2j} \right) \sigma_{\epsilon}^2 = \frac{1}{1 - \varphi^2} \sigma_{\epsilon}^2, \rho_0 = 1$$

$$\gamma_1 = \left(\sum_{j=0}^{\infty} \varphi^j \varphi^{j+1} \right) \sigma_{\epsilon}^2 = \varphi \left(\sum_{j=0}^{\infty} \varphi^{2j} \right) \sigma_{\epsilon}^2 = \frac{\varphi}{1 - \varphi^2} \sigma_{\epsilon}^2, \rho_1 = \varphi$$

$$\gamma_1 = E(x_t x_{t-1}) = E((\varphi x_{t-1} + \epsilon_t)(x_{t-1})) = \varphi \sigma_x^2, \rho_1 = \varphi$$

$$\gamma_2 = E(x_t x_{t-2}) = E((\varphi^2 x_{t-2} + \varphi \epsilon_{t-1} + \epsilon_t)(x_{t-2})) = \varphi^2 \sigma_x^2, \rho_2 = \varphi^2$$

$$\gamma_k = E(x_t x_{t-k}) = E((\varphi^k x_{t-k} + \epsilon_t)(x_{t-k})) = \varphi^k \sigma_x^2, \rho_k = \varphi^k$$



Strong and Weak Stationarity

- We used that the moments do not depend on time:

$$\begin{aligned} E(x_t) &= E(x_s) \text{ for all } t \text{ and } s \\ E(x_t x_{t-j}) &= E(x_s x_{s-j}) \text{ for all } t \text{ and } s \end{aligned}$$

- This property is true for invertible ARMA models. But reflects a much more important property:
- Definitions:
 - A process is strongly or strictly stationary if the joint probability $\{x_{i_1+k}, x_{i_2+k}, \dots, x_{i_n+k}\}$ is independent of k .
 - A process is weakly or covariance stationary if $E(x_t)$ and $E(x_t^2)$ are finite and $E(x_t x_{t-j})$ does not depend on t .



Conditions for Stationary ARMA's

1. Strong stationarity does not \rightarrow weak stationarity. $E(x_t^2)$ must be finite.
2. Strong stationarity plus $E(x_t)$ and $E(x_t^2) < \infty \rightarrow$ weak stationarity.
3. Weak stationarity does not \rightarrow strong stationarity.
4. Weak stationarity plus normality \rightarrow strong stationarity.

Conditions for stationary ARMA:

1. MA process: $x_t = \sum_{j=0}^{\infty} \theta_j \epsilon_{t-j}$, $var(x_t) = \sum_{j=0}^{\infty} \theta_j^2 \sigma^2$, so second moments exist iff the MA coefficients are square summable:

$$\text{stationary MA} \leftrightarrow \sum_{j=0}^{\infty} \theta_j^2 < \infty$$

If second moments exist it is easy to see that they are independent of t .

2. AR process: for the variance to be finite the AR lag polynomial must be invertible:

$$A(L)x_t = (1 - \mu_1 L)(1 - \mu_2 L) \dots x_t = \epsilon_t$$



Conditions for Stationary ARMA's

For the variance to be finite, the AR lag polynomial must be invertible or $|\mu_i| < 1$ for all i . Common way to say this:

$$A(L) = \text{constant}(L - \rho_1)(L - \rho_2) \dots$$

Where ρ_i are the roots of the lag polynomial, rewriting the above equation:

$$A(L) = \text{constant}(-\rho_1) \left(1 - \frac{1}{\rho_1} L\right) (-\rho_2) \left(1 - \frac{1}{\rho_2} L\right) \dots$$

Thus $\frac{1}{\mu_i} = \rho_i$, hence the rule $|\mu_i| < 1$ means $\rho_i > 1$, or since the roots of the lag polynomial can be complex:

AR stationary if all the roots of the lag polynomial lie outside the unit circle, ie. The lag polynomial is invertible.



Literature

1. John H. Cochrane: Time Series for Macroeconomics and Finance
2. G. Koop, D. Korobilis: Bayesian Multivariate Time Series Methods
3. F. Schorfheide, M. Del Negro: Bayesian Macroeconomics
4. Bolla, Krámlí: Statisztikai következtetések elmélete
5. Durbin, Koopman: Time Series Analysis by State Space Methods
6. Tusnády, Ziermann: Idősorok analízise



Websites with Data Sources

1. US data:

- <http://www.bea.gov/>
- <https://fred.stlouisfed.org/>

2. World data:

- <http://www.imf.org/en/data>
- <https://www.bis.org/>

3. European data:

- <https://www.ecb.europa.eu/stats/keyind/html/index.en.html>