

CSE250B Homework 4

Qiao Zhang

February 12, 2016

1 regression problem

1.1

Given $L(w)$, we can calculate the $H(w)$

$$\begin{aligned} H_{jk} &= \frac{\partial^2 L}{\partial w_k \partial w_j} \\ &= \frac{\partial}{\partial w_k} (-2 \sum_{i=1}^n x_j^{(i)} (y^{(i)} - w \cdot x^{(i)})) \\ &= -2 \sum_{i=1}^n x_j^{(i)} \frac{\partial}{\partial w_k} (y^{(i)} - w \cdot x^{(i)}) \\ &= 2 \sum_{i=1}^n x_j^{(i)} x_k^{(i)} \\ &= 2 x_j \cdot x_k \end{aligned}$$

where $x_j = [x_j^{(1)}, x_j^{(2)}, \dots, x_j^{(n)}]$, $x_k = [x_k^{(1)}, x_k^{(2)}, \dots, x_k^{(n)}]$

We can apply matrix decomposition to $H(w)$ by

$$H = 2 \begin{bmatrix} - & - & x_1 & - & - \\ - & - & x_2 & - & - \\ & & \dots & & \\ - & - & x_p & - & - \end{bmatrix} \begin{bmatrix} | & | & \dots & | \\ x_1 & x_2 & \dots & x_p \\ | & | & \dots & | \end{bmatrix} = V V^T$$
$$V = \sqrt{2} \begin{bmatrix} - & - & x_1 & - & - \\ - & - & x_2 & - & - \\ & & \dots & & \\ - & - & x_p & - & - \end{bmatrix}$$

Thus H is positive semi-definite, which implies that $L(w)$ is convex.

1.2

$$H_{jk} = \frac{\partial L}{\partial w_j} = -2 \sum_{i=1}^n x_j^{(i)} (y^{(i)} - w \cdot x^{(i)})$$

$$\nabla L(w) = -2 \sum_{i=1}^n x^{(i)} (y^{(i)} - w \cdot x^{(i)})$$

$$w_{t+1} = w_t - \eta_t \nabla L(w_t)$$

$$w_{t+1} = w_t + 2\eta_t \sum_{i=1}^n x^{(i)} (y^{(i)} - w_t \cdot x^{(i)})$$

1.3

$$w_{t+1} = w_t - \eta_t H^{-1}(w_t) \nabla L(w_t)$$

Because $H(w)$ is positive semi-definite, $H(w)$ can be written as

$$H(w) = Q \Lambda Q^T$$

Thus,

$$H^{-1}(w) = Q \Lambda^{-1} Q^T$$

where $H^{-1}(w_t)$ can be easily obtained.

$$w_{t+1} = w_t - \eta_t \frac{L(w)}{\nabla L(w_t)} = w_t - \eta_t \frac{L(w)}{2 \sum_{i=1}^n x^{(i)} (y^{(i)} - w_t \cdot x^{(i)})}$$

2 Convexity

2.1

$$\begin{aligned}
 H_{jk} &= \frac{\partial^2 f}{\partial x_j \partial x_k} \\
 &= \frac{\partial}{\partial x_j} \left(\frac{\partial x^T M x}{\partial x_k} \right) \\
 &= \frac{\partial}{\partial x_j} \left(\frac{\partial \sum_{i,j} M_{ij} x_i x_j}{\partial x_k} \right) \\
 &= \frac{\partial}{\partial x_j} (\sum_i M_{ik} x_i + \sum_i M_{ki} x_i) \\
 &= M_{jk} + M_{kj}
 \end{aligned}$$

Therefore,

$$H = M + M^T$$

Because M is positive semi-definite, M^T and H are also positive semi-definite. Thus f is convex.

2.2

$$\begin{aligned}
 H_{jk} &= \frac{\partial^2 f}{\partial x_j \partial x_k} \\
 &= \frac{\partial}{\partial x_j} \left(\frac{\partial e^{u \cdot x}}{\partial x_k} \right) \\
 &= \frac{\partial}{\partial x_j} (u_k e^{u \cdot x}) \\
 &= u_k u_j e^{u \cdot x}
 \end{aligned}$$

$$H = \begin{bmatrix} - & - & u_1 e^{u \cdot x/2} & - & - \\ - & - & u_2 e^{u \cdot x/2} & - & - \\ & & \dots & & \\ - & - & u_p e^{u \cdot x/2} & - & - \end{bmatrix} \begin{bmatrix} | & | & \dots & | \\ u_1 e^{u \cdot x/2} & u_2 e^{u \cdot x/2} & \dots & u_p e^{u \cdot x/2} \\ | & | & \dots & | \end{bmatrix} = V V^T$$

$$V = \begin{bmatrix} - & - & u_1 e^{u \cdot x/2} & - & - \\ - & - & u_2 e^{u \cdot x/2} & - & - \\ & & \dots & & \\ - & - & u_p e^{u \cdot x/2} & - & - \end{bmatrix}$$

Thus H is positive semi-definite, which implies that f is convex.

2.3

$\because g$ and h are convex

$$\therefore \forall a, b \in R^p, \theta \in [0, 1] \quad g(\theta a + (1 - \theta)b) \leq \theta g(a) + (1 - \theta)g(b)$$

$$\therefore \forall a, b \in R^p, \theta \in [0, 1] \quad h(\theta a + (1 - \theta)b) \leq \theta h(a) + (1 - \theta)h(b)$$

$$\therefore \forall a, b \in R^p, \max[g(\theta a + (1 - \theta)b), h(\theta a + (1 - \theta)b)] \leq \theta \max[g(a), h(a)] + (1 - \theta) \max[g(b), h(b)]$$

$$\therefore \forall a, b \in R^p, \theta \in [0, 1] \quad f(\theta a + (1 - \theta)b) \leq \theta f(a) + (1 - \theta)f(b)$$

$\therefore f$ is convex

3 Logistic regression using gradient descent

3.1 b

For part b, a function named logistic is implemented with parameters of input data, input label, max_iteration and step size.

Four samples are shown below in order to show the influence of step size and maximum iteration.

```
w = logistic(X, Y, 100000, 0.1)
```

```
iteration = 29513
```

```
w = [ 24.28620473 2.59508653 -89.11172933]
```




Figure 1: decision boundary and six points

```
w = logistic(X, Y, 100000, 0.05)
iteration = 100000
w = [ 8.81024328 0.64992294 -27.95754635]
```




Figure 2: decision boundary and six points

```
w = logistic(X, Y, 100000, 0.2)
iteration = 26769
w = [ 50.02437432 4.63249848 -193.38130935]
```




Figure 3: decision boundary and six points

```
w = logistic(X, Y, 500, 0.2)
iteration = 500
w = [ 16.61027093 5.03188882 -65.61743524]
```




Figure 4: decision boundary and six points

3.2 c

After scaling down the x2-axis, the number of iterations needed for convergence increases. But the corresponding margin also increases, in other words, the decision boundary divides the points in a better way.

```
X = [[2,0.1,1], [1,2,1], [1,0.5,1], [4,0.1,1], [1,4,1], [3,3,1]]
```

```
Y = [-1, -1, -1, 1, 1, 1]
```

```
w = logistic(X, Y, 100000, 0.1)
```

```
iteration = 100000
```

```
w = [ 9.55167574 7.04368379 -30.3522845 ]
```




Figure 5: decision boundary and six points

3.3 d

I used two bi-variate Gaussians, each generate 50 random samples.

```
a = np.random.multivariate_normal([0,0], [[1,0],[0,2]], 50)
```

```
b = np.random.multivariate_normal([3,3], [[3,1],[1,2]], 50)
```

```
w = logistic(X, Y, 100000, 0.07)
```

```
iteration = 134
```

```
w = [-1.1394483 -1.55500097 3.77205524]
```

There are many mis-classified points as a result of overlapping classes.




Figure 6: decision boundary and six points