

## Matrix Numerov method for solving Schrödinger's equation

Mohandas Pillai, Joshua Goglio, and Thad G. Walker

Citation: *American Journal of Physics* **80**, 1017 (2012); doi: 10.1119/1.4748813

View online: <https://doi.org/10.1119/1.4748813>

View Table of Contents: <https://aapt.scitation.org/toc/ajp/80/11>

Published by the *American Association of Physics Teachers*

---

### ARTICLES YOU MAY BE INTERESTED IN

[Computer Solutions to the Schrödinger Equation](#)

*American Journal of Physics* **40**, 730 (1972); <https://doi.org/10.1119/1.1986627>

[The renormalized Numerov method applied to calculating bound states of the coupled-channel Schroedinger equation](#)

*The Journal of Chemical Physics* **69**, 4678 (1978); <https://doi.org/10.1063/1.436421>

[Getting started with Numerov's method](#)

*Computers in Physics* **11**, 514 (1997); <https://doi.org/10.1063/1.168593>

[New numerical methods applied to solving the one-dimensional eigenvalue problem](#)

*The Journal of Chemical Physics* **67**, 4086 (1977); <https://doi.org/10.1063/1.435384>


[The Fourier grid Hamiltonian method for bound state eigenvalues and eigenfunctions](#)

*The Journal of Chemical Physics* **91**, 3571 (1989); <https://doi.org/10.1063/1.456888>

[What is the limit  \$\hbar \rightarrow 0\$  of quantum theory?](#)

*American Journal of Physics* **80**, 1009 (2012); <https://doi.org/10.1119/1.4751274>

---



VIRTUAL WINTER MEETING  
JANUARY 9-12

# Matrix Numerov method for solving Schrödinger's equation

Mohandas Pillai, Joshua Goglio, and Thad G. Walker<sup>a)</sup>

Department of Physics, University of Wisconsin-Madison, Madison, Wisconsin 53706

(Received 16 May 2012; accepted 15 August 2012)

We recast the well-known Numerov method for solving Schrödinger's equation into a representation of the kinetic energy operator on a discrete lattice. With just a few lines of code in a high-level programming environment such as MATHEMATICA, it is simple to calculate and plot accurate eigenvalues and eigenvectors for a variety of potential problems. We illustrate the method by calculating high-accuracy solutions for the  $|x|$  potential. © 2012 American Association of Physics Teachers. [http://dx.doi.org/10.1119/1.4748813]

## I. INTRODUCTION

With modern high-level programming and visualization environments such as MATHEMATICA, MATLAB, and PYTHON, it is possible and desirable to use computational methods to illustrate and illuminate many basic physics principles with a minimum of programming overhead. This latter point is a key: if the programming is too difficult and/or time consuming, the focus shifts from the physics to the programming.

A case in point is the solution of boundary value problems for the one-dimensional Schrödinger equation. One typically starts at one boundary with an assumed value for the energy, then integrates to the other boundary where the boundary conditions are tested. A new guess for the energy is generated, and the process is repeated until the desired level of accuracy is obtained.<sup>1</sup> Using the Numerov method, the numerical integration can be done with relatively high accuracy even with large step sizes.<sup>1-3</sup> Though straightforward, this process is tedious to program when one needs to solve for a large number of eigenstates.

An alternative approach that gives a large number of eigenstates simultaneously is to expand the wave function in a set of orthogonal basis states that satisfy the appropriate boundary conditions. By truncating the basis set, the Hamiltonian can be diagonalized with built-in matrix routines. The resulting eigenvectors can be used to generate a superposition of the basis states to represent the spatial wave function.

A particularly attractive hybrid approach is to discretize the wave function on a (linear) lattice. This is equivalent to expanding in a basis set of Dirac  $\delta$  functions centered at the lattice points. The eigenvectors are then simply lists of the values of the various eigenfunctions at the lattice points. This allows straightforward visualization of the eigenfunctions, as demonstrated recently on time-dependent problems.<sup>4</sup>

In the discretized approach, the potential energy operator is simply a diagonal matrix of the potential energy evaluated at each lattice point. The kinetic energy operator, being a differential operator, is more difficult to realize. The purpose of this paper is to show how the Numerov method can be used to represent the kinetic energy operator on the lattice in a straightforward manner, allowing for high-accuracy solutions to be obtained with very straightforward programs using matrix diagonalization.

## II. MATRIX NUMEROV REPRESENTATION OF THE HAMILTONIAN

The Numerov method<sup>1-3</sup> is a specialized integration formula for numerically integrating differential equations of the form

$$\psi''(x) = f(x)\psi(x). \quad (1)$$

For the time-independent 1-D Schrödinger equation,  $f(x) = -2m(E - V(x))/\hbar^2$ . On a lattice of points  $x_i$  evenly spaced by a distance  $d$ , the integration formula is

$$\psi_{i+1} = \frac{\psi_{i-1}(12 - d^2 f_{i-1}) - 2\psi_i(5d^2 f_i + 12)}{d^2 f_{i+1} - 12} + O(d^6), \quad (2)$$

where, for example,  $\psi_i = \psi(x_i)$ . This can be rearranged into the form

$$\begin{aligned} & -\frac{\hbar^2}{2m} \frac{(\psi_{i-1} - 2\psi_i + \psi_{i+1}))}{d^2} \\ & + \frac{V_{i-1}\psi_{i-1} + 10V_i\psi_i + V_{i+1}\psi_{i+1}}{12} \\ & = E \frac{(\psi_{i-1} + 10\psi_i + \psi_{i+1}))}{12}. \end{aligned} \quad (3)$$

Now, if we represent  $\psi$  as the column vector  $(\dots\psi_{i-1}, \psi_i, \psi_{i+1}, \dots)$  and define matrices  $A = (\mathbb{I}_{-1} - 2\mathbb{I}_0 + \mathbb{I}_1)/d^2$ ,  $B = (\mathbb{I}_{-1} + 10\mathbb{I}_0 + \mathbb{I}_1)/12$ ,  $V = \text{diag}(\dots V_{i-1}, V_i, V_{i+1}, \dots)$ , where  $\mathbb{I}_p$  is a matrix of 1s along the  $p$ th diagonal, and zeros elsewhere, this becomes the matrix equation

$$-\frac{\hbar^2}{2m} A\psi + BV\psi = EB\psi. \quad (4)$$

Multiplying by  $B^{-1}$ , we get

$$-\frac{\hbar^2}{2m} B^{-1}A\psi + V\psi = E\psi. \quad (5)$$

The first term is the Numerov representation of the kinetic energy operator.

On an  $N$ -point grid, the boundary conditions are implemented by taking  $N \times N$  submatrices of  $A$  and  $B$ . This corresponds to the condition  $\psi_0 = \psi_{N+1} = 0$ ; effectively we have placed the potential of interest inside an infinite-walled box. Alternatively, one can use periodic boundary conditions, with  $A_{1,N} = A_{N,1} = 1/d^2$  and  $B_{1,N} = B_{N,1} = 1/12$ .

We choose the grid in the following manner, valid for finding bound-state solutions to attractive potentials. It is easy to extend the following advice to other cases. Suppose we wish to find all the states with  $E < E_m$  above the potential minimum. The minimum local de Broglie wavelength is, therefore,  $\lambda = h/\sqrt{2mE_m}$ . We have found that sufficient

```

(*Potential, desired max energy*)
V[s_] := Abs[s]; em = 20.;
(*determine grid*)
st = FindRoot[V[s] == em, {s, em}][[1, 2]];
ds = 1 / Sqrt[2 em];
n = Round[2 (st / ds + 4 Pi)];
s = Table[-ds (n + 1) / 2 + ds i, {i, n}];
(*calculate KE matrix*)
one[n_, d_] :=
  DiagonalMatrix[1 + 0 Range[n - Abs[d]], d];
B = (one[n, -1] + 10 one[n, 0] + one[n, 1]) / 12.;
A = (one[n, -1] - 2 one[n, 0] + one[n, 1]) / ds^2;
KE = -Inverse[B].A / 2;
(*Hamiltonian*)
H = KE + DiagonalMatrix[V[s]];
(*Energies, wavefunctions*)
{eval, evec} = Eigensystem[H];

```

Fig. 1. Full MATHEMATICA code for solving the potential problem  $V = |s|$ . The energy of the  $m$ th state can be accessed by the command `eval[[m]]`, and the corresponding list of wave function values at the grid points  $s$  is `evec[[m]]`.

accuracy is generally obtained by taking the grid spacing  $d$  corresponding to about one point per radian, i.e.,  $d = \lambda/2\pi$ . The number of grid points needed can be estimated by finding the outer turning points  $x_t$  such that  $V(x_t) = E_m$ , and allowing for an extra  $2\lambda$  in the classically forbidden region. Thus,  $N = 2(x_t/d + 4\pi)$  rounded to the nearest integer.

### III. EXAMPLE: $|x|$ POTENTIAL

The “linear” potential  $V(x) = b|x|$  is analytically solvable<sup>5</sup> and so is useful for comparison to the numerical calculations. Introducing scaled variables  $s = x(mb/\hbar^2)^{1/3}$  and  $\epsilon = E(m/b^2\hbar^2)^{1/3}$ , we get

$$-\frac{1}{2}B^{-1}A\psi + |s|\psi = \epsilon\psi. \quad (6)$$

Following the guidelines for selecting the grid, we pick a grid spacing  $d_s = 1/\sqrt{2\epsilon_m}$  in order to find accurate results for states up to energy  $\epsilon_m$ . The turning point is  $s_t = \epsilon_m$ , so with the additional two de Broglie wavelengths outside the turning point, we get  $N = 2(4\pi + s_t/d_s)$ . A MATHEMATICA code for solving this problem is shown in Fig. 1. The grid used is  $d_s = 0.158$ ,  $N = 278$ , and the program runs in typically 0.1 s of CPU time.

A comparison of the exact and matrix Numerov results for some of the energy levels is given in Table I. It is remarkable that such high-accuracy results can be achieved with such a simple program. An even simpler program might be obtained

Table I. Comparison of exact and numerical results for the quantized energies (in scaled units, see text) of the  $|x|$  potential. The numerical results are calculated on an  $N = 278$  point grid with a spacing of 0.158 in scaled distance units.

	$n = 1$	2	3	4	10	20	50
Exact	0.8086	1.8558	2.5781	3.2446	6.3053	10.182	18.947
Numerov	0.8099	1.8557	2.5785	3.2445	6.3049	10.181	18.936
3-pt	0.8089	1.8529	2.5728	3.2358	6.2717	10.094	18.634




Fig. 2. The lower graph shows the analytical Airy function solution to the Schrödinger equation for the  $n = 50$  state of the  $|x|$  potential, compared to the Numerov method eigenfunction (dots). The upper graph shows the difference between the analytical and Numerov wave functions.

by using a simple three-point approximation to the second derivative, equivalent to setting  $B = 1$ . The results are also shown in Table I and are of similar quality to the Numerov method for small  $n$  but give clearly less accurate results for large  $n$ . Figure 2 compares the Numerov and exact wave functions for  $n = 50$ .

We have used the matrix Numerov method to solve a variety of problems. These include the harmonic oscillator, particle in a box, hydrogen atom, and the partner super potential to the particle in a box.<sup>6</sup> With periodic boundary conditions, we have solved the cosine potential and the periodic square well. We have also simulated a double-well potential to demonstrate tunneling. Among these, the hydrogen atom is the most challenging due to its singularity at  $x = 0$  and the rapid increase of the classical turning point with principal quantum number. Codes for a few of these examples are available online.<sup>7</sup>

### IV. CONCLUSIONS

While there are other related methods, examples being b-splines<sup>8</sup> and the Fourier grid representation,<sup>9</sup> that may be superior for accurate calculation of energies and wave functions by experts, we feel that the simplicity of the matrix Numerov method introduced here makes it ideal for classroom and course work settings as a tool for helping students grow comfortable with the notion of the wave function and eigenvalue problems. The method, though illustrated here for the prominent 1-D Schrödinger equation, is of course useful for any Numerov-type problem. We have used it to solve for diffusion modes of simple geometries, for example.

We have also experimented with variable-mesh grids, which can be useful for certain problems such as the hydrogen atom where such grids can improve the accuracy. However, this adds new complexities that detract from what we feel is the main advantage of the method: attainment of very high accuracy for minimum programming complexity and computer time.

## ACKNOWLEDGMENTS

This work was supported by the National Science Foundation. This paper is an outgrowth of work done by two of the authors (M.P. and J.G.) as part of the Atomic and Quantum Physics course at the University of Wisconsin-Madison.

<sup>a)</sup>Electronic mail: [tgwalker@wisc.edu](mailto:tgwalker@wisc.edu)

<sup>1</sup>J. M. Blatt, "Practical points concerning the solution of the Schrödinger equation," *J. Comput. Phys.* **1**, 382–396 (1967).

<sup>2</sup>P. C. Chow, "Computer solutions to the Schrödinger equation," *Am. J. Phys.* **40**, 730–734 (1972).

<sup>3</sup>R. H. Landau, M. J. Paez, and C. C. Bordeianu, *A Survey of Computational Physics* (Princeton U.P., Princeton, NJ, 2008).

<sup>4</sup>M. Belloni and W. Christian, "Time development in quantum mechanics using a reduced Hilbert space approach," *Am. J. Phys.* **76**, 385–392 (2008).

<sup>5</sup>J. J. Sakurai and J. J. Napolitano, *Modern Quantum Mechanics*, 2nd ed. (Addison Wesley, San Francisco, 2010).

<sup>6</sup>A. Gangopadhyaya, J. V. Mallow, and C. Rasinariu, *Supersymmetric Quantum Mechanics: An Introduction* (World Scientific, Hackensack, NJ, 2011).

<sup>7</sup>See supplementary material at <http://dx.doi.org/10.1119/1.4748813> or <http://www.physics.wisc.edu/~tgwalker/NumerovExamples/> for the code of Fig. 1 and some worked examples.

<sup>8</sup>A. Derevianko, E. Luc-Koenig, and F. Masnou-Seeuws, "Application of B-splines in determining the eigenspectrum of diatomic molecules: Robust numerical description of halo-state and Feshbach molecules," *Can. J. Phys.* **87**, 67–74 (2009).

<sup>9</sup>C. C. Marston and G. G. Balint-Kurti, "The Fourier grid Hamiltonian method for bound state eigenvalues and eigenfunctions," *J. Chem. Phys.* **91**, 3571–3576 (1989).



### Arc Lamp

From the 1916 catalogue of the L.E. Knott Apparatus Co. of Boston: "Demonstration Arc Lamp with Reflector for use with any strong source of battery current, but excellent for use with our [hand-cranked dynamo] in demonstrating the principle of arc lamps. It is a near approach to the automatic form, adjustment by counterpoise gravity feed. ... A simple reflector attached greatly adds to the effectiveness of the design, \$3.30" As the upper carbon was consumed, the gap between the electrodes was adjusted by moving the counterweight in. The end of the upper carbon has been hollowed out, indicating that it was connected to the positive side of the power supply. This example is at Creighton University. (Notes and photograph by Thomas B. Greenslade, Jr., Kenyon College)