

Martin Lindquist

Department of Biostatistics
Johns Hopkins
Bloomberg School of Public Health

Tor Wager

Department of Psychology and
Neuroscience and the
Institute for Cognitive Science
University of Colorado, Boulder


The General Linear Model: Introduction

Statistical Analysis

- There are multiple goals in the statistical analysis of fMRI data.
- They include:
 - localizing brain areas activated by the task;
 - determining networks corresponding to brain function; and
 - making predictions about psychological or disease states.




Data Processing Pipeline



Overview of the GLM analysis process

Typically a two-level hierarchical analysis

1. Within-subject (individual)
 2. Across-subject (group)
- Can be done in stages (1st, 2nd level)
 - Hierarchical models combine stages into one model





University of Colorado **Boulder**




INTRODUCING THE GLM FAMILY

General Linear Model

- The **general linear model** (GLM) approach treats the data as a linear combination of model functions (predictors) plus noise (error).
- The model functions are assumed to have **known** shapes (i.e., straight line, or known curve), but their amplitudes (i.e., slopes) are **unknown** and need to be estimated.
- The GLM framework encompasses many of the commonly used techniques in fMRI data analysis (and data analysis more generally).



The GLM Family



Simple regression

ANOVA

Multiple regression

General linear model

- Mixed effects/hierarchical
- Timeseries models (e.g., autoregressive)
- Robust, penalized regression (LASSO, Ridge)



Generalized linear models

- Non-normal errors
- Binary/categorical outcomes (logistic regression)

Closed form solution

Iterative solutions

Simple regression: A simple linear model with one predictor



Simplification:
“linear relationship”


Find slope,
intercept

Test slope:
P-value


Meaning of
relationship?




The GLM Family



The GLM Family



The multiple regression model basic model for the GLM



- Structural model for regression

Variables DV Pred1 Pred2 Predk

$$y_i = \beta_0 + \beta_1 x_{i1} + \beta_2 x_{i2} + \cdot + \beta_k x_{ik} + \varepsilon_i$$

Parameters **intercept** Slope 1 Slope 2 Slope k Error

• solve for beta vector

Matrix notation

$$y = X\beta + \varepsilon$$

- minimize sum of squared residuals



Matrix Notation

- Alternatively, we can write

$$\mathbf{Y} = \mathbf{X}\beta + \varepsilon$$


as

$$\begin{bmatrix} Y_1 \\ Y_2 \\ \vdots \\ Y_n \end{bmatrix} = \begin{bmatrix} 1 & X_{11} & \cdots & X_{1p} \\ 1 & X_{21} & \cdots & X_{2p} \\ \vdots & \vdots & & \vdots \\ 1 & X_{np} & \cdots & X_{np} \end{bmatrix} \times \begin{bmatrix} \beta_0 \\ \beta_1 \\ \vdots \\ \beta_p \end{bmatrix} + \begin{bmatrix} \varepsilon_1 \\ \varepsilon_2 \\ \vdots \\ \varepsilon_n \end{bmatrix}$$

Observed Data
Design matrix
Model parameters
Residuals

A non-fMRI example

- Does exercise predict life-span?
- Made-up (not real data)
- Control for other variables that might be important, i.e., gender (M/F)




A non-fMRI example

$$\mathbf{Y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\varepsilon}$$

$$\begin{bmatrix} Y_1 \\ Y_2 \\ \vdots \\ Y_n \end{bmatrix} = \begin{bmatrix} 1 & X_{11} & \cdots & X_{1p} \\ 1 & X_{21} & \cdots & X_{2p} \\ \vdots & \vdots & & \vdots \\ 1 & X_{np} & \cdots & X_{np} \end{bmatrix} \times \begin{bmatrix} \beta_0 \\ \beta_1 \\ \vdots \\ \beta_p \end{bmatrix} + \begin{bmatrix} \varepsilon_1 \\ \varepsilon_2 \\ \vdots \\ \varepsilon_n \end{bmatrix}$$

Outcome Data Design matrix Model parameters Residuals




End of Module



@fMRIstats

A standard processing pipeline




General Linear Model

- General model
 - Categorical or continuous predictor
- Linear model
 - Parameters that are
 - not multiplied by other parameters, e.g., not $Y = abX$
 - only first-power, e.g., NOT $Y = b^2X$
 - not exponents, e.g., NOT $Y = X^b$
 - Variables do not need to satisfy the above criteria
 - transformation workaround, e.g., $Y = bX^2$ can be rewritten as $Y = bZ$
 - Not necessary straight-line relationship



Human Brain Mapping

- The most common use of fMRI to date has been to **localize** areas of the brain that activate in response to a certain task.
- These types of **human brain mapping** studies are necessary for the development of biomarkers and increasing our understanding of brain function.



The regression model

- Vector notation: y , x , and e are vectors of N values corresponding to N observations

$$y = b + mx + e$$

Outcome (DV)	Intercept (constant)	slope Predictor value	Error (residual)
-----------------	-------------------------	-----------------------------	---------------------


For point i :

$$y_i = b + mx_i + e_i$$



Fitting and residual variance

- Minimize squares of vertical distance to lines.
- Minimize: $\sum e_i^2 = \sum (y_i - \hat{y}_i)^2$
- The line is being pulled by vertical rubber bands attached to each point. Vector of red lines is e



Matrix Notation

$$\begin{bmatrix} Y_1 \\ Y_2 \\ \vdots \\ Y_n \end{bmatrix} = \begin{bmatrix} 1 & X_{11} & \cdots & X_{1p} \\ 1 & X_{21} & \cdots & X_{2p} \\ \vdots & \vdots & & \vdots \\ 1 & X_{np} & \cdots & X_{np} \end{bmatrix} \times \begin{bmatrix} \beta_0 \\ \beta_1 \\ \vdots \\ \beta_p \end{bmatrix} + \begin{bmatrix} \varepsilon_1 \\ \varepsilon_2 \\ \vdots \\ \varepsilon_n \end{bmatrix}$$

Observed Data
Design matrix
Model parameters
Residuals

Is the same as

$$\begin{bmatrix} Y_1 \\ Y_2 \\ \vdots \\ Y_n \end{bmatrix} = \beta_0 \begin{bmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{bmatrix} + \beta_1 \begin{bmatrix} X_{11} \\ X_{21} \\ \vdots \\ X_{np} \end{bmatrix} + \beta_p \begin{bmatrix} X_{1p} \\ X_{2p} \\ \vdots \\ X_{np} \end{bmatrix} + \begin{bmatrix} \varepsilon_1 \\ \varepsilon_2 \\ \vdots \\ \varepsilon_n \end{bmatrix}$$



Design matrix

- In fMRI the design matrix specifies how the factors of the model change over time.
- The design matrix is an $n \times p$ matrix where n is the number of observations over time and p is the number of model parameters

$$\mathbf{X} = \begin{bmatrix} 1 & X_{11} & \cdots & X_{1p} \\ 1 & X_{21} & \cdots & X_{2p} \\ \vdots & \vdots & & \vdots \\ 1 & X_{np} & \cdots & X_{np} \end{bmatrix}$$

