

OSSD - Calculus and Vector

MCV4U

Qinghao Hu

February 11, 2026

Unit 1: Rate of change. In the first unit, we will discuss the concept of Rate of Change, Continuity and Limits. Also, the the introduction to the Derivatives will be discussed.

Contents

1	Slope	3
1.1	Measuring Rates of Change on a Table or a Graph	3
1.1.1	Interval of the function	3
1.1.2	Determine approximation for the slope of the tangent . .	3
1.1.3	Trends and increasing/decreasing	4
1.2	Measuring Rates of Change Using a Defined Relation	5
1.2.1	Difference Quotient	5
1.2.2	Approximating the slope of the tangent	5
2	Reference Sheet	4
2.1	Assume Knowledge	4
2.1.1	Log Laws	4
2.1.2	Trig Laws	4

1 Slope

1.1 Measuring Rates of Change on a Table or a Graph

Definition 1.1 (slope/Rate of Change). the ratio of the vertical distance to the horizontal distance between two points.

In Grade 9, we all studied the slope formula:

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

m represents the slope because linear function is written as:

$$f(x) = mx + b$$

1.1.1 Interval of the function


When you are asked to give the interval of a function, one must response in one of these forms:

$$x \in (-\infty, a) \cap (b, \infty)$$

Remark. We look the function from $-\infty$ to ∞

Example 1.2. Here is a question by Mr.Hoken

8. Given the graph below, determine the regions on the function where:



a) the y values are increasing

$$x \in (-\infty, -3) \text{ or } -3 < x < 2, x \in \mathbb{R}$$

b) the rate of change (r.o.c.) is increasing

$$x < -3 \cup x > 5, x \in \mathbb{R}$$

c) the y values are negative and the slopes are positive

$$-3 < x < 2, x \in \mathbb{R}$$

d) the slopes are negative and the r.o.c. is decreasing

$$x \in (2, 5), x \in \mathbb{R}$$

The answer for a) should be $x \in (-\infty, -3)$

The answer for b) should be $x \in (-\infty, -3) \cup (5, \infty)$

1.1.2 Determine approximation for the slope of the tangent

To start off, we need to two points which have equal difference of x to the point that tangent line cross over.

Let's assume we have points 1, 2, and 3. Point 2 touches the tangent line. Solve for the slope between point 1 and point 2:

$$m_1 = \frac{y_2 - y_1}{x_2 - x_1}$$

Solve for the slope between point 2 and point 3:

$$m_2 = \frac{y_3 - y_2}{x_3 - x_2}$$

The slope at point 2 that touches tangent line should be:

$$m = \frac{m_2 + m_1}{2}$$

1.1.3 Trends and increasing/decreasing

Here are the summary Mr. Hoken made for this class

- An increasing trend in y values corresponds to positive slopes of tangents in that interval
- A decreasing trend in y values corresponds to negative slopes of tangents in that interval
- "Rate of Change" in y values refers to your slope values. We use standard expressions to describe the trend in Slope values
 - * "The rate of change is increasing" means that the slope values are climbing.
ex. (0.5, 3, 4, 5, 6, 20, ...) or (-10, -9, ..., 1, 2)
 - * "The rate of change is decreasing" means that the slope values are dropping.
ex. (25, 11, 9, 5, ..., -3, -7, -20)

1.2 Measuring Rates of Change Using a Defined Relation

1.2.1 Difference Quotient

Definition 1.3 (Difference Quotient). a measure of the average rate of change of the function over an interval. In other words, it is a simplified expression for the slope of a secant. It can be expressed as

$$m = \frac{f(x+h) - f(x)}{h}$$

where $h \neq 0$

Let's proof the difference Quotient

Proof. We have the formula of the slope:

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

Assume we have two points: $(x, f(x))$ and $(x+h, f(x+h))$, sub these two points into the formula of slope.

$$\begin{aligned} m &= \frac{f(x+h) - f(x)}{(x+h) - x} \\ m &= \frac{f(x+h) - f(x)}{h} \end{aligned}$$

□

Let's look at an example.

Example 1.4. Determine a general expression for the slope given the function $f(x) = -x^2 + 4x + 5$

$$\begin{aligned} m &= \frac{f(x+h) - f(x)}{h} \\ m &= \frac{-(x+h)^2 + 4(x+h) + 5 - (-x^2 + 4x + 5)}{h} \\ m &= \frac{-x^2 - 2xh - h^2 + 4x + 4h + 5 + x^2 - 4x - 5}{h} \\ m &= \frac{-2xh - h^2 + 4h}{h} \\ m &= -2x + 4 - h, h \neq 0 \end{aligned}$$

1.2.2 Approximating the slope of the tangent

Recall:

- The slope of a secant line will give the average rate of change of the 'y' values over a given interval of 'x'.
- The slope of a tangent line will give the instantaneous rate of change of the 'y' value at a given value of 'x'.

For difference quotient

$$\frac{f(x+h) - f(x)}{h}$$

if h is extremely small, we can approximate the slope of the tangent line.

In later sections, we will discuss about principle of limits. At that time, you will have a better understanding about today's topic.

2 Reference Sheet

2.1 Assume Knowledge

2.1.1 Log Laws

$$\log_b A + \log_b B = \log_b (A \times B)$$

$$\log_b \left(\frac{M}{N}\right) = \log_b M - \log_b N$$

$$\log_b M^k = k \log_b M$$

$$\log_b b^k = k$$

$$b^{\log_b k} = k$$

2.1.2 Trig Laws

$$\tan \theta = \frac{\sin \theta}{\cos \theta}$$

$$\cot \theta = \frac{\cos \theta}{\sin \theta}$$

$$\csc \theta = \frac{1}{\sin \theta}$$

$$\sec \theta = \frac{1}{\cos \theta}$$

$$\sin^2 \theta + \cos^2 \theta = 1$$

$$1 + \cot^2 \theta = \csc^2 \theta$$

$$\tan^2 \theta + 1 = \sec^2 \theta$$

$$\sin(A \pm B) = \sin A \cos B \pm \sin B \cos A$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(\alpha \pm \beta) = \frac{\tan \alpha \pm \tan \beta}{1 \mp \tan \alpha \tan \beta}$$

sin Law:

$$\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}$$

Cosine Law:

$$a^2 = b^2 + c^2 - 2bc \cos A$$

Unit Circle:

