

Math 135

Please give me feedback! I am at the first row.

Qinghao Hu

September 16, 2026

Contents

1	Admin	3
1.1	General Information	3
1.1.1	Contact	3
1.1.2	Teaching Style	3
1.1.3	Graded Assignments	3
1.1.4	Late Submission Policy	3
1.1.5	Office Hours	3
2	Introduction to the Language of Mathematics	4
2.1	Introduction to Proof	4
2.2	Sets	4
2.3	Statement and Negation	4
2.4	Quantifiers and Quantified Statements	5
2.4.1	Universal Quantifier	5
2.4.2	Existential quantifier	5
2.4.3	Nested Quantifier	5
3	Logical Analysis of Mathematical Statements	6
3.1	Truth Table and Negation	6
3.2	Conjunction and Disjunction	6
3.3	Implication	8
3.4	Converse and Contrapositive	8
3.4.1	Converse	8
3.4.2	Contrapositive	9
3.4.3	If and Only If	9

1 Admin

1.1 General Information

1.1.1 Contact

For mathematics questions: `abeltaos@uwaterloo.ca`

For course logistics: `math135@uwaterloo.ca`

1.1.2 Teaching Style

The LEARN website contains several useful resources. In particular:

- (i) The icons at the top of the page provide quick access to important resources, including the course schedule.
- (ii) Crowdmark is used for submitting assignments.

1.1.3 Graded Assignments

There are three assignments each week.

1.1.4 Late Submission Policy

Late submissions are accepted with a penalty of 1% per minute for the entire assignment.

1.1.5 Office Hours

Monday and Wednesday 2:00 to 3:30 in (MC 6503) or by appointment (email the instructor to arrange).

2 Introduction to the Language of Mathematics

2.1 Introduction to Proof

Definition 2.1 (Proof). A proof is a logically valid argument that establishes the truth of a mathematical statement using definitions, axioms, and previously proved results.

Proposition 2.2. *For every positive integer n , the number $n^2 + 1$ is not a perfect square.*

Proof. Let n be a positive integer. The consecutive perfect squares surrounding $n^2 + 1$ are n^2 and $(n + 1)^2$. Indeed,

$$n^2 < n^2 + 1 < n^2 + 2n + 1 = (n + 1)^2,$$

where the second inequality follows from $n > 0$.

Therefore, $n^2 + 1$ lies strictly between two consecutive perfect squares, so it cannot itself be a perfect square. ■

2.2 Sets

Definition 2.3 (Set). A **set** is a well-defined, unordered collection of distinct objects. Each object that appears in this collection is called an **element** of the set.

For a finite set, we can denote it by listing its elements, separated by commas, surrounding the entire list by braces (or by $\{\}$).

Example 2.4. The set containing the elements 2, 7, and -3 can be denoted by

$$\{7, -3, 2\}.$$

Example 2.5. Here are some examples:

- $\{2, 4, 6, 8\}$
- $\{1, 2, \{1, 2, 3\}\}$
- $\{a, b, c, d\}$
- $\mathbb{N} = \{1, 2, 3, \dots\}$
- $\mathbb{Z} = \{\dots, -1, 0, 1, \dots\}$
- $\mathbb{Q} = \{\frac{a}{b} : a, b \in \mathbb{Z}, b \neq 0\}$

Fact: $|\mathbb{N}| = \aleph_0$, $|\mathbb{R}| = \aleph_1$ $\aleph_1 = 2^{\aleph_0}$.

Definition 2.6. The **parity** of an integer is **even** if the integer is a multiple of 2, and **odd** otherwise.

2.3 Statement and Negation

Definition 2.7 (Statement). A **statement** is a sentence that has a definite state of being either true or false.

Definition 2.8 (Negation). Suppose that A is a statement. Then the negation of A , denoted by $\neg A$, is the statement asserting the opposite truth value to A . That is, $\neg A$ is false when A is true, and $\neg A$ is true when A is false.

2.4 Quantifiers and Quantified Statements

2.4.1 Universal Quantifier

$\forall$ means "for all", and it is called the universal quantifier.

Example 2.9.

$$\forall x \in \mathbb{Z}, x \geq 5, 2^n > n^2.$$

The negation of universal quantifier.

$$\neg(\forall x \in S, P(x)) \equiv (\exists x \in S, \neg P(x))$$

2.4.2 Existential quantifier

$\exists$ means "there exists", and it is called the existential quantifier.

Example 2.10.

$$\exists a \in \mathbb{Z}, a^2 + 29a + 209 \leq 0.$$

The negation of an existentially quantifier statement (such as $\exists x \in S, P(x)$) is:

$$\neg(\exists x \in S, P(x)) \equiv (\forall x \in S, \neg P(x)).$$

Remark. A quantified statement contains four parts:

- variable;
- quantifier;
- domain;
- an open sentence involving the variable (that is either true or false whenever a value of the variable is chosen from the domain).

2.4.3 Nested Quantifier

Consier this statement

$$\forall x \in \mathbb{R}, \exists y \in \mathbb{R}, x > y.$$

That is a true statement. (Given $x \in \mathbb{R}$, you could provide $y = x - 1$ to prove the existential statement)

Remark. Order matters! Consider the statement: $\exists y \in \mathbb{R}, \forall x \in \mathbb{R}, x > y$. This says there is a real number y which is smaller than every real number! False!!!

To negative a nested quantifier statement, think about it in pieces. For example:

$$\begin{aligned} \neg(\forall x \in S, \exists y \in T, P(x, y)) &\equiv \\ \exists x \in S, \forall y \in T, \neg P(x, y). \end{aligned}$$

3 Logical Analysis of Mathematical Statements

3.1 Truth Table and Negation

Definition 3.1. The truth value for "not A ", written symbolically as $\neg A$, is defined by the truth table given below.

A	$\neg A$
T	F
F	T

We also refer to $\neg A$ as the negation of A .

We can use the truth table to prove that an argument is equal to another argument.

Example 3.2. Given the truth table for $A, \neg A, \neg(\neg A)$,

A	$\neg A$	$\neg(\neg A)$
T	F	T
F	T	F

This truth table serves as a proof that $A \equiv \neg(\neg A)$.

Remark. "A" is considered a statement variable.

3.2 Conjunction and Disjunction

Definition 3.3 (Compound statement). A compound statement is a statement composed of several individual statements, each of which is called a component statement.

Definition 3.4. The truth value for $A \wedge B$, is defined by the truth table

A	B	$A \wedge B$
T	T	T
T	F	F
F	T	F
F	F	F

We also refer to $A \wedge B$ as the conjunction of A and B .

If we have three component statement, we need 2^3 rows to cover all the possibility.

Example 3.5. The compound statement

$$(2 + 3 = 6) \wedge (\pi + 2 > 5)$$

is false, since the statement $2 + 3 = 6$ is false.

Definition 3.6. The truth value for A or B , written as $A \vee B$, is defined by the truth table

A	B	$A \vee B$
T	T	T
T	F	T
F	T	T
F	F	F

We also refer to $A \vee B$ as the **disjunction** of A and B .

Remark. The disjunction $A \vee B$ is the *inclusive* or: it is true whenever at least one of A , B is true, including the case where both are true. This differs from the *exclusive* or (xor), written $A \oplus B$, which is true when exactly one of A , B is true and false when both are true. Thus $A \vee B$ and $A \oplus B$ agree in every case except $A = B = \text{T}$, where $A \vee B$ is true but $A \oplus B$ is false.

Example 3.7. The Compound statement:

$$(2 + 3 = 6) \vee (\pi + 2 \geq 5),$$

is true since one of the statements is true.

Example 3.8. Let us use a truth table to prove that

$$\neg(A \vee B) \equiv (\neg A \wedge \neg B)$$

A	B	$A \vee B$	$\neg(A \vee B)$	$\neg A$	$\neg B$	$\neg A \wedge \neg B$
T	T	T	F	F	F	F
T	F	T	F	F	T	F
F	T	T	F	T	F	F
F	F	F	T	T	T	T

A similar equivalence is

$$\neg(A \wedge B) \equiv ((\neg A) \vee (\neg B)).$$

These things together are called DeMorgan's Law.

Theorem 3.9 (DeMorgan's Law). *For statement variable A and B , we have*

$$\neg(A \vee B) \equiv (\neg A \wedge \neg B)$$

$$\neg(A \wedge B) \equiv (\neg A \vee \neg B)$$

Proposition 3.10. *For statements A, B, C , the following rules hold for the logical operations $\wedge$ and $\vee$:*

Commutative Laws

$$A \vee B \equiv B \vee A$$

$$B \wedge A \equiv A \wedge B$$

Associative Laws

$$A \vee (B \vee C) \equiv (A \vee B) \vee C$$

$$A \wedge (B \wedge C) \equiv (A \wedge B) \wedge C$$

Distributive Laws

$$A \vee (B \wedge C) \equiv (A \vee B) \wedge (A \vee C)$$

$$A \wedge (B \vee C) \equiv (A \wedge B) \vee (A \wedge C)$$

3.3 Implication

Definition 3.11. The truth value for *A implies B*", written symbolically as $A \implies B$, is defined by the truth table

A	B	$A \implies B$
T	T	T
T	F	F
F	T	T
F	F	T

We also refer to $A \implies B$ as an implication.
Just for fun. Instructor does not talk about it.

$$A \implies B \equiv \neg(A \wedge \neg B) \equiv \neg A \vee B.$$

Hypothesis: Assume to be true: A

Conclusion: B

Example 3.12.

$$(\forall x \in \mathbb{R}) [(x > 2) \implies (x^2 > 4)]$$

Remark. Interesting.

$$(A \vee B) \implies C \equiv (A \implies C) \wedge (B \implies C)$$

3.4 Converse and Contrapositive

3.4.1 Converse

The $B \implies A$ is called the converse of $A \implies B$.

A	B	$A \implies B$	$B \implies A$
T	T	T	T
T	F	F	T
F	T	T	F
F	F	T	T

Remark. $(A \implies B) \not\equiv (B \implies A)$ (not logical equivalent)

3.4.2 Contrapositive

The implication $(\neg B) \implies (\neg A)$ is called the contrapositive of $A \implies B$.

A	B	$\neg A$	$\neg B$	$A \implies B$	$(\neg B) \implies (\neg A)$
T	T	F	F	T	T
T	F	F	T	F	F
F	T	T	F	T	T
F	F	T	T	T	T

Remark. $(A \implies B) \equiv ((\neg B) \implies (\neg A))$

An implication is also logically equivalent to another type of compound statement, as shown in the following truth table:

A	B	$\neg A$	$A \implies B$	$\neg A \vee B$
T	T	F	T	T
T	F	F	F	F
F	T	T	T	T
F	F	T	T	T

Remark. The negation of an implication is not an implication.

3.4.3 If and Only If

Definition 3.13. The truth value for "A if and only if B", written symbolically as $A \iff B$, is defined by the truth table.

A	B	$A \iff B$
T	T	T
T	F	F
F	T	F
F	F	T