

Day 7

1. Environments

a list of binding, clear that tell different variable has different values
<http://ku-sldg.github.io/plih//ids/2-Adding-Environments.html>

Approximating the semantics of `let` has proved tricky:

- We can approximate the cbn reduction relation, but only at the cost of performing much of the substitution that we might hope to avoid
- We have even less success with the cbv reduction, as we can't substitute approximations into terms

Our approach to this is to reconsider the role that substitutions play in evaluation. Rather than applying substitutions immediately to terms, we'll **preserve substitutions as *environments* in the reduction relation.**

2. CBV with Environments

We define ***environments*** to be mappings from terms to values: $H \in \mathcal{X} \rightarrow \mathcal{V}$.

(Aside: H is supposed to be a Greek capital eta, not a Latin H. What difference does it make? None.)

Now we can define a 3-place evaluation relation $\Downarrow \in (\mathcal{X} \rightarrow \mathcal{V}) \times \mathcal{T} \times \mathcal{V}$

$$\frac{H, t_1 \Downarrow v_1 \quad H, t_2 \Downarrow v_2}{H, t_1 + t_2 \Downarrow v_1 + v_2} \quad \frac{H, t_1 \Downarrow v_1 \quad H, t_2 \Downarrow v_2}{H, t_1 \div t_2 \Downarrow \lfloor t_1/t_2 \rfloor} (v_2 \neq 0)$$

$$\frac{}{H, x \Downarrow_{\text{cbv}} H(x)} \quad \frac{H, t_1 \Downarrow_{\text{cbv}} v_1 \quad H[x \mapsto v_1], t_2 \Downarrow_{\text{cbv}} v_2}{H, \text{let } x = t_1 \text{ in } t_2 \Downarrow_{\text{cbv}} v_2}$$

- We could introduce a new evaluation symbol (or new subscript) for the three-place version of the evaluation relation... but the context will always make it clear which version we mean.
- **The constant rules behave the same** in call-by-name and call-by-value
- We write $H(x)$ to denote the value that **x is mapped to in H** , and $H[x \mapsto v]$ to denote **extending a partial function**... formally:

$$H[x \mapsto v](y) = \begin{cases} v & \text{if } x = y \\ H(y) & \text{otherwise} \end{cases}$$

We have some simple derivations:

$$\frac{\frac{}{\emptyset, 4 \Downarrow_{\text{cbv}} 4} \quad \frac{\frac{\{x \mapsto 4\}, x \Downarrow_{\text{cbv}} 4 \quad \{x \mapsto 4\}, x \Downarrow_{\text{cbv}} 4}{\{x \mapsto 4\}, x \div x \Downarrow_{\text{cbv}} 1}}{\emptyset, \text{let } x = 4 \text{ in } x \div x \Downarrow_{\text{cbv}} 1}}$$

$$\frac{\frac{\overline{\emptyset, 4 \Downarrow_{\text{cbv}} 4} \quad \overline{\emptyset, 4 \Downarrow_{\text{cbv}} 4}}{\overline{\emptyset, 4 \div 4 \Downarrow_{\text{cbv}} 1}} \quad \frac{\overline{\{x \mapsto 1\}, x \Downarrow_{\text{cbv}} 1} \quad \overline{\{x \mapsto 1\}, x \Downarrow_{\text{cbv}} 1}}{\overline{\{x \mapsto 1\}, x + x \Downarrow_{\text{cbv}} 2}}}{\overline{\emptyset, \text{let } x = 4 \div 4 \text{ in } x + x \Downarrow_{\text{cbv}} 2}}$$

- Starting with the **empty environment** (\emptyset), and $\emptyset[x \mapsto 1] = \{x \mapsto 1\}$.
- Results of evaluation equivalent to substitution version: same final value, same number of operations. **However, we replace substitution with variable lookup.** Complexity implications?

3. CBN with Environments

Intuitively:

- CBV evaluates *before* substituting
- CBN **evaluates after** substituting

To **map this intuition to environments**, we have:

- CBV environments store *values*
- CBN environments store *terms*

So for CBN, we define $H \in \mathcal{X} \rightarrow \mathcal{T}$, and have evaluation rules

$$\frac{H, H(x) \Downarrow_{\text{cbn}} v}{H, x \Downarrow_{\text{cbn}} v} \quad \frac{H[x \mapsto t_1], t_2 \Downarrow_{\text{cbn}} v}{H, \text{let } x = t_1 \text{ in } t_2 \Downarrow_{\text{cbn}} v}$$

Again, we can consider some simple derivations:

$$\frac{\frac{\overline{\{x \mapsto 4 \div 4\}, 4 \Downarrow_{\text{cbn}} 4} \quad \overline{\{x \mapsto 4 \div 4\}, 4 \Downarrow_{\text{cbn}} 4}}{\overline{\{x \mapsto 4 \div 4\}, 4 \div 4 \Downarrow_{\text{cbn}} 1}} \quad \frac{\overline{\{x \mapsto 4 \div 4\}, 4 \Downarrow_{\text{cbn}} 4} \quad \overline{\{x \mapsto 4 \div 4\}, 4 \Downarrow_{\text{cbn}} 4}}{\overline{\{x \mapsto 4 \div 4\}, 4 \div 4 \Downarrow_{\text{cbn}} 1}}}{\overline{\{x \mapsto 4 \div 4\}, x \Downarrow_{\text{cbn}} 1} \quad \overline{\{x \mapsto 4 \div 4\}, x \Downarrow_{\text{cbn}} 1}}{\overline{\{x \mapsto 4 \div 4\}, x + x \Downarrow_{\text{cbn}} 2}}{\overline{\emptyset, \text{let } x = 4 \div 4 \text{ in } x + x \Downarrow_{\text{cbn}} 2}}$$

代表
整体

直接引用所用项t2

and:

$$\frac{\overline{\{x \mapsto 4 \div 0\}, 3 \Downarrow_{\text{cbn}} 3}}{\overline{\emptyset, \text{let } x = 4 \div 0 \text{ in } \Downarrow_{\text{cbn}} 3}}$$

- Same properties of evaluation: evaluation repeated for each use of a variable, **but unused variables don't stop evaluation.** 变量名不一样，不进行带入或替换

4. Approximating Evaluation

We build an **approximation** of evaluation with environments following the same approach we've used for earlier evaluation relations.

Let environments $\Gamma \in \mathcal{X} \rightarrow \mathcal{P}(\mathcal{S})$ **map variables to approximations of values.**

近似值，或者叫types

Now, we define approximation by:

$$\begin{array}{c}
\frac{\Gamma, t_1 \Downarrow_{\pm} S_1 \quad \Gamma, t_2 \Downarrow_{\pm} S_2}{\Gamma, t_1 + t_2 \Downarrow_{\pm} \bigcup \{s_1 \hat{+} s_2 \mid s_1 \in S_1, s_2 \in S_2\}} \quad \frac{\Gamma, t_1 \Downarrow_{\pm} S_1 \quad \Gamma, t_2 \Downarrow_{\pm} S_2}{\Gamma, t_1 \div t_2 \Downarrow_{\pm} \bigcup \{s_1 \hat{\div} s_2 \mid s_1 \in S_1, s_2 \in S_2\}} \\
\frac{}{\Gamma, x \Downarrow_{\pm} H(x)} \quad \frac{\Gamma, t_1 \Downarrow_{\pm} S_1 \quad \Gamma[x \mapsto S_1], t_2 \Downarrow_{\pm} S_2}{\Gamma, \text{let } x = t_1 \text{ in } t_2 \Downarrow_{\pm} S_2}
\end{array}$$

As we would hope, the approximation relation now exactly follows the pattern of the evaluation relation!

This is still an **over-approximation** of the meanings of terms:

$$\frac{\frac{\overline{\emptyset, 5 \Downarrow_{\pm} \{+\}} \quad \overline{\emptyset, 5 \Downarrow_{\pm} \{+\}}}{\overline{\emptyset, 5 - 5 \Downarrow_{\pm} \{-, 0, +\}}} \quad \frac{\overline{\{x \mapsto \{-, 0, +\}\}, 5 \Downarrow_{\pm} \{+\}} \quad \overline{\{x \mapsto \{-, 0, +\}\}, x \Downarrow_{\pm} \{-, 0, +\}}}{\overline{\{x \mapsto \{-, 0, +\}\}, 5 \div x \Downarrow_{\pm} \{-, +\}}}{\overline{\emptyset, \text{let } x = 5 - 5 \text{ in } 5 \div x \Downarrow_{\pm} \{-, +\}}}$$

But this is no different than our previous approximation for $5 \div (5 - 5)$, so we should not be surprised.