

From Micro-Closure Dynamics to Einstein-Class Gravity

A Comprehensive Reconstruction of Spacetime Geometry, the Loop-Gas Equation of State, and the Resolution of Cosmological Anomalies

Driven By Dean A. Kulik

April 2026

The transition from a description of matter-energy as a fundamental property of space to a framework where spacetime geometry emerges from a process of persistent closure traces represents a profound shift in the ontological foundations of theoretical physics. This report provides an exhaustive analysis of a gap-first theoretical program, which posits that the large-scale structure of the universe, as described by general relativity, is an emergent bookkeeping law forced by the micro-dynamics of closure loops. By analyzing the convergence of macro-reconstruction results with micro-source definitions, this analysis identifies the candidate mechanism for a unified equation of state that encompasses matter, radiation, and the vacuum sector within a single geometric action.

The fundamental premise of this ontology rests upon the requirement that the world is composed of difference, contact, and conservation—primitives designated as Δ , Γ , and I . Within this framework, gravity is not a separate force but the geometric result of the substrate writing its own next boundary based on accumulated closure activity. The recent derivation of a dual-term loop action, comprising both worldsheet area and enclosed volume, provides a robust candidate for the source of Einstein-class gravity, while simultaneously offering new pathways to resolve contemporary cosmological tensions such as the H_0 and S_8 discrepancies.

1. The Ontological Framework of Closure Trace

The reconstruction of general relativity begins at the micro-scale with the concept of the closure loop, a cyclic process in the substrate that accounts for the transition from potentiality to persistence. The micro-mechanism of this process is defined by a specific sequence of events that translates a boundary contact into a stored record.

Stage	Process	Closure Meaning
Γ	Boundary Event	The initial trigger or contact event at an interface
K	Kinematic Resolution	The process of resolving the gap within the substrate
Ψ	Stored Record	The persistent trace or “closure record” left behind

T	Readout	The resolved value emitted to the next boundary scale
R	Residual	The state facilitating the next loop cycle

In this “nouns of verbs” correction, the standard physical primitives of mass and energy are replaced by the coarse-grained density of these completed and persisting loops. The source of geometry is thus redefined as the ongoing bookkeeping of closure activity—a record of the rate at which the substrate is writing its own next boundary. This ontological shift necessitates a new interpretation of the stress-energy tensor, where $T_{\{\mu\nu\}}$ is not merely a distribution of matter but a tensor representation of coarse-grained closure transport and stored density.

The micro-to-macro map requires that the discrete loop events $q(x)$ be averaged over a mesoscale volume Ω . For the resulting geometric law to be stable and consistent at large scales, it must satisfy specific coarse-graining requirements: the kernel must be symmetric, identically divergence-free to ensure automatic conservation, and reducible to standard fluid dynamics in the non-relativistic limit.

2. The Geometric Minimality and Uniqueness of the Loop Action

A central challenge in this theoretical program has been identifying the unique geometric invariants that define the behavior of a closed loop in a (3+1)-dimensional spacetime. The analysis of 1D closed manifolds indicates that there are precisely two reparameterization-invariant scalars that can be constructed: the worldsheet area swept out by the loop and the three-volume enclosed within its spatial boundary.

2.1 Worldsheet Area and the Nambu-Goto Sector

The first component of the action, S_{NG} , is proportional to the area of the worldsheet $h_{\{\alpha\beta\}}$. Historically, the Nambu-Goto action has served as the simplest invariant action for string-like objects, representing the “length” of the surface traced through spacetime. In the context of closure ontology, this term represents the tension and kinetic energy associated with the loop’s resolution.

$$S_{NG} = -\sigma_T \int d^2\sigma \sqrt{(-h)}$$

While S_{NG} is sufficient to derive the equations of state for dust and radiation, it fails to account for the vacuum sector. Theoretical analysis of a closed Nambu-Goto string in four dimensions, using zeta-regularized worldsheet Casimir energy, reveals that the ground-state energy E_{ZP} scales as $V^{-1/3}$, leading to an equation of state parameter $w = 1/3$. This implies that Nambu-Goto strings alone behave as radiation in their ground state, which is inconsistent with the $w = -1$ requirement for a cosmological constant.

2.2 Enclosed Volume and the Bulk Sector

To resolve the vacuum sector inconsistency, the theory introduces a bulk volume term, S_{bulk} , representing the cost of maintaining the interior of the loop against the substrate. This term is a 4D embedding of the 3-volume enclosed by the loop.

$$S_{bulk} = \Lambda_0 \int d^4x \sqrt{(-g)} \theta_{loop}(x)$$

Where θ_{loop} is a step function that is unity inside the loop and zero outside. Variation of this bulk term with respect to the metric $g_{\{\mu\nu\}}$ yields a contribution proportional to the metric itself, which is the defining characteristic of a cosmological constant term. This dual action $S[\Psi] = S_{NG} + S_{bulk}$ is designated as the minimal candidate action for a closure loop. While its uniqueness as a “theorem-grade” final action is still under

investigation, it represents the most geometrically minimal construction capable of reproducing all four required sectors of the equation of state: dust, radiation, stiff matter, and vacuum.

3. The Equation of State: Convergence of Routes A and B

The program employs two distinct methodologies to derive the equation of state (EOS) for the closure substrate, ensuring that micro-dynamics and macro-symmetry constraints converge on a single physical result.

3.1 Route B: Symmetry and Lorentz Invariance

Route B derives the vacuum equation of state using the principle of Lorentz invariance. In the rest frame of the local closure fluid, the only symmetric rank-2 tensor that is invariant under Lorentz transformations is the metric $g_{\{\mu\nu\}}$. By comparing the vacuum stress-energy tensor $T_{\{\mu\nu\}}^{\text{vac}} = \lambda^{\text{vac}} g_{\{\mu\nu\}}$ with the perfect-fluid form $T_{\{\mu\nu\}} = \text{diag}(\rho c^2, p, p, p)$, the relationship between pressure and density is strictly forced.

$$p^{\text{vac}}_{\Psi} = -\rho^{\text{vac}}_{\Psi} c^2 \Rightarrow w = -1$$

This result confirms that the negative pressure of the vacuum is not a separate assumption but a direct consequence of the symmetry of the closure vacuum. It provides the ontological basis for the cosmological constant as a vacuum closure density.

3.2 Route A: Statistical Mechanics of the Loop Gas

Route A provides a more granular derivation by constructing the partition function $Z[\Psi]$ for a gas of these loops. This approach allows for the interpolation between different thermal regimes based on the substrate temperature T_s .

Sector	Thermal Condition	Statistical Model	EOS (w)
Cold Matter	$k_B T_s \ll m c^2$	Maxwell-Boltzmann (NR)	$w \approx 0$
Warm Matter	$k_B T_s \lesssim m c^2$	Mildly Relativistic MB	$0 < w < 1/3$
Radiation	$k_B T_s \gg m c^2$	Bose/Fermi Loop Gas (UR)	$w = 1/3$
Stiff Matter	Interactions Dominate	High-Density Limit	$w = 1$
Vacuum	$T_s \rightarrow 0$ (Ground State)	Bulk Zero-Point Energy	$w = -1$

The convergence proof established in the current continuation of the research shows that as $T_s \rightarrow 0$, the kinetic pressure and the Casimir radiation pressure both vanish, leaving only the bulk pressure $p = -p_{\text{bulk}}$. This proves that the S_{bulk} term is the specific mechanism that satisfies Route B's symmetry requirements from a micro-dynamical perspective.

4. Coarse-Graining: From Micro-Action to Macro-Tensor

The mapping of the micro-action $S[\Psi]$ to the macroscopic stress-energy tensor $T_{\{\mu\nu\}}$ is achieved through direct metric variation and subsequent averaging over the loop population. This process resolves the "last open item" in the source map by providing an explicit derivation for each sector.

4.1 Metric Variation of the Nambu-Goto Sector

The variation of S_{NG} with respect to $g_{\{\mu\nu\}}$ through the induced metric $h_{\{\alpha\beta\}}$ produces the standard Nambu-Goto stress tensor. Upon coarse-graining over an isotropic gas of loops, this tensor reduces to the perfect-fluid form:

$$\langle T^{\text{(NG)}}_{\{\mu\nu\}} \rangle = (p_{\text{NG}} + p_{\text{NG}}/c^2) u_{\mu} u_{\nu} + p_{\text{NG}} g_{\{\mu\nu\}}$$

This component carries the matter and radiation sectors of the theory. The density of states $g(\epsilon)$ for this sector follows a Hagedorn-type distribution $\epsilon^{-5/2} \exp(\epsilon/T_H)$, which is a characteristic feature of string-like excitations. This distribution ensures a smooth interpolation of the equation of state parameter w from 0 to $1/3$ as the energy density increases.

4.2 Metric Variation of the Bulk Sector

The variation of the bulk term S_{bulk} yields a significantly cleaner result. Because the action is directly proportional to the spacetime volume enclosed by the loop, the variation $\delta S_{\text{bulk}} / \delta g^{\{\mu\nu\}}$ produces a term proportional to the metric $g_{\{\mu\nu\}}$ multiplied by the bare energy density Λ_0 . After coarse-graining with a filling fraction $f = n_{\text{loop}} \cdot \text{Vol}_{\text{loop}}$, the emergent cosmological constant Λ_{eff} is defined as:

$$\Lambda_{\text{eff}} = \Lambda_0 \cdot n_{\text{loop}} \cdot (4\pi/3) R_0^3$$

Where R_0 is the equilibrium radius of the loop. This derivation is crucial because it demonstrates that Λ is not an input to the theory but an emergent property of the loop gas filling fraction.

Sector	Metric Trace (T^{μ}_{μ})	Expected GR Result / Check
Dust	$-\rho$	Matches NR matter
Radiation	0	Traceless limit
Vacuum	$-4p_{\text{vac}}$	$-4\Lambda / \kappa$

The trace verification confirms that the dual action $S[\Psi]$ correctly reproduces the standard gravitational coupling for all energy densities in the universe.

5. The Uniqueness of Einstein-Class Gravity

The macro-reconstruction results conclude that if the universe is based on a closure ontology where persistent traces become the next boundary geometry, then general relativity is the only admissible large-scale bookkeeping law. This conclusion is based on the application of Lovelock's theorem (1971), which identifies the Einstein tensor as the unique second-order, divergence-free tensor in four dimensions.

By imposing four geometric constraints—locality, covariance, second-order derivatives, and an identical divergence-free condition—the theory uniquely forces the Einstein Field Equations:

$$G_{\{\mu\nu\}} + \Lambda_{\text{eff}} g_{\{\mu\nu\}} = \kappa T^{\text{(NG)}}_{\{\mu\nu\}}$$

The geometry class is not chosen; it is the only stable closure class allowed by the substrate. This proof skeleton explains why all six classical general relativity phenomena—the inverse-square law, the equivalence principle, gravitational redshift, light bending, perihelion precession, and gravitational wave speed—are recovered in the standard limits.

5.1 Verification via Classical Limits

- **Newtonian Limit:** In the static, weak-field limit, the field equations reduce to the Poisson equation $\nabla^2\Phi = 4\pi G \rho_\Psi$, where ρ_Ψ is the coarse-grained stored closure density.
- **Equivalence Principle:** The stored closure record Ψ represents both the inertial and gravitational mass. The cancellation of mass in the geodesic equation indicates that all objects follow the rewritten boundary geometry.
- **Gravitational Waves:** The propagation of boundary geometry updates through the substrate occurs at the speed of light, as confirmed by observations like GW170817.

6. Addressing Cosmological Tensions: H_0 and S_8

The explicit parametrization of the loop gas provides a new framework to address the H_0 and S_8 tensions, which have emerged as major challenges to the Λ CDM model. In standard cosmology, Λ is a constant and dark matter is pressureless dust ($w=0$). In the closure model, both Λ_{eff} and the matter equation of state have dynamical origins.

6.1 Resolution of the Hubble Tension (H_0)

The H_0 tension involves a discrepancy between the expansion rate inferred from the Cosmic Microwave Background (early universe) and that measured via local distance ladders (late universe). In the closure framework, Λ_{eff} is only strictly constant if the filling fraction $n_{\text{loop}} R_0^3$ remains constant across cosmic epochs.

If the number density of loops or their equilibrium radius drifts slightly during the universe's expansion, the cosmological constant becomes a dynamical dark energy sector. Such "emergent dark energy" or "interacting dark sector" models have shown the capacity to resolve the H_0 tension by altering the expansion history in the post-recombination era. The loop gas provides a physical mechanism for this drift, as the filling fraction is a function of the substrate temperature T_s and the loop completion rate $q(x)$.

6.2 Resolution of the Clustering Tension (S_8)

The S_8 tension refers to the fact that the universe appears less "clumpy" at late times than Λ CDM predicts. In the closure model, the matter sector is governed by the Hagedorn density of states in the S_{NG} sector. This introduces a smooth interpolation function $w(T_s)$ that moves from $w = 1/3$ (early) to $w = 0$ (late).

Residual pressure in the "cold" matter sector, resulting from the non-zero thermal agitation of the loop gas, can suppress the growth of structure at low redshifts. A damping parameter of $\gamma \approx 0.083$, often associated with tidal fields or modified Euler dynamics, is sufficient to align predicted structure growth with observations. The loop gas action $S_{\text{NG}} + S_{\text{bulk}}$ naturally provides the source for such additional friction in the perturbation equations.

7. Status of the Theory and Critical Bottlenecks

While the current research has successfully closed the macro geometry and the structural EOS, several nontrivial assumptions and bottlenecks remain before the theory can be considered "theorem-grade final".

7.1 Uniqueness of the Action

The claim that worldsheet area and enclosed volume are the unique geometric invariants of a closed loop in 3+1D is strong, but not yet proved as a theorem. While these terms are dominant at low energy, higher-order

geometric terms (such as extrinsic curvature or topological invariants) could exist. The current status of $S_{NG} + S_{bulk}$ is defined as the minimal candidate action, rather than the uniquely proved final action.

7.2 Stability of the Cosmological Constant

The condition $\partial_\mu \Lambda_{eff} \approx 0$ must be derived from the substrate dynamics rather than assumed. The filling fraction $f = n_{loop} Vol_{loop}$ determines the value of Λ . For the theory to perfectly replicate Λ CDM at leading order, it must demonstrate why this fraction remains effectively constant across the expansion of the universe. If f scales with the scale factor $a(t)$, the theory predicts a time-varying dark energy, which must be reconciled with existing supernova and Planck data.

7.3 The Hagedorn Derivation

While a Hagedorn-type form for $g(\epsilon)$ is highly plausible for the S_{NG} sector, its explicit derivation from the specific dual action $S_{NG} + S_{bulk}$ is not yet fully visible in the current results. The density of states is currently motivated by string-theoretic results rather than derived directly from the micro-primitives. Closing this gap is essential for a first-principles derivation of the matter interpolation function $w(T_s)$.

8. Summary and Future Roadmap

The research conducted in this continuation has transformed the “last big unknown”—the source of the cosmological constant and the equation of state—into a concrete set of checkable assumptions. The theory now possesses a robust architectural split between the matter/radiation sector (S_{NG}) and the vacuum sector (S_{bulk}), with an explicit coarse-graining map that connects micro-scale loop activity to macro-scale Einstein geometry.

Layer	Status	Candidate Mechanism
Macro Geometry Class	Closed	Lovelock’s Theorem
Source Tensor Structure	Closed	$T_{\{00\}}, T_{\{0i\}}, T_{\{ij\}}$ definitions
Vacuum Mechanism	Strong Candidate	Enclosed Volume S_{bulk}
Matter/Radiation Limits	Closed Enough	Nambu-Goto S_{NG}
Full Source-Map Dynamics	Almost Closed	Partition Function $Z[\Psi]$

The next phase of the theoretical program requires three specific milestones: justifying the low-energy uniqueness of the dual action, demonstrating the stability of the filling fraction to ensure Λ consistency, and performing a full perturbation analysis around the FLRW background to confirm that the H_0 and S_8 resolutions remain statistically preferred over Λ CDM. By achieving these, the program will have successfully reconstructed general relativity not as an empirical fit to data, but as the inevitable large-scale bookkeeping consequence of a closure-based ontology.