

Quantum Cut-Density as the Origin of Spacetime Curvature

The ρ_r Field, the Helix Gravity Kernel, and the Complete Curvature Chain

Dean A. Kulik

April 2026

Abstract

This paper formalizes the claim that local spacetime curvature is generated by concentrated quantum non-separability. Where the entanglement cut-density $\rho_r(x)$ is high — where internal boundaries are densely packed and quantum degrees of freedom cannot be factored across them — the effective source density is elevated above the classical matter density, and the metric curves accordingly. This is not a metaphor. The chain is:

non-separability concentration $\rightarrow$ effective source elevation $\rightarrow$ curvature $\rightarrow$ gravity.

The paper derives the Helix Gravity kernel from the CLG closure substrate, identifies ρ_r with the mutual-information density across internal boundaries, and presents the corrected field equation, the axial-speed suppression law, and the full curvature-density map. New contributions include: the complete derivation of the Nexus harmonic constant $H = \pi/9$ from the recursive binary closure fixed point; the derivation of the axial-speed coupling constant λ from the CLG minimal dual action; verification of the Bianchi identity for $T_{\mu\nu}^{(r)}$; the SHA-256 carry-fraction notebook in closed form; gravitational lensing predictions for the Bullet Cluster; and integration with the Nexus-Friedmann control law addressing the H_0 and S_8 tensions. The honest status of each claim — theorem-grade versus bridge — is maintained throughout.

$$\rho_{\text{eff}}(x) = \rho_m(x) + \alpha s_{\text{ent}}(x), \quad \alpha = H^2/24, \quad H = \pi/9$$

1. The Claim in Plain Language

The quantum layer is not uniform. In some regions, the web of entanglement and non-local correlation is dense: boundaries between subsystems are close together, the mutual information across those boundaries

is high, and the field cannot be factored into independent parts. In other regions the quantum substrate is sparse — few cuts, low entanglement, low mutual information, and the state factorizes cleanly.

The density of that non-separability is $\rho_r(x)$. The central claim of this paper is that wherever ρ_r is high, the effective source term in the gravitational field equation is elevated, and spacetime curves accordingly. On this account, gravity is not the force that attracts mass in any fundamental sense. It is the *geometric response to the concentration of computational irreducibility in the quantum substrate*.

Central Identification

$$\rho_r(x) \equiv \text{entanglement-cut density at } x$$

The density of internal boundaries across which quantum information cannot be freely transported. Wherever this density is high, the local metric is sourced by more than classical matter alone.

This identification was already implicit in the CLG program's earlier statement that gravity is an emergent side-effect of information processing in regions of high informational density. The contribution of the present paper is to sharpen that statement into a candidate kernel with explicit coupling constants — both α and λ now derived rather than fitted — a covariant field equation with verified Bianchi consistency, a closed-form SHA-256 carry-fraction correspondence, and concrete observational predictions that make contact with the H_0 and S_8 tensions.

Throughout, we maintain the distinction between what is formally closed, what is a candidate kernel supported by internal consistency, and what remains a bridge requiring external validation. No claim is promoted beyond its earned status.

2. The ρ_r Operator: Definition and Properties

2.1 Entanglement Cut-Density

Let S be a compact spatial region and $I_S = \partial S$ its boundary. For a quantum state $\hat{\rho}$ on the Hilbert space bipartitioned as $\mathcal{H} = \mathcal{H}_S \otimes \mathcal{H}_{\bar{S}}$, define the entanglement entropy across I_S :

$$S_{\text{ent}}(I_S) = -\text{Tr}_S[\hat{\rho}_S \ln \hat{\rho}_S], \quad \hat{\rho}_S = \text{Tr}_{\bar{S}}[\hat{\rho}].$$

The local entanglement-entropy density is the volume derivative in the limit of small S :

$$s_{\text{ent}}(x) = \lim_{|S| \rightarrow 0} \frac{S_{\text{ent}}(I_S)}{|S|}, \quad [s_{\text{ent}}] = \text{nats} \cdot \text{m}^{-3}.$$

For a region containing N_{cuts} internal boundaries, each carrying mutual information I_k and located at position x_k , the cut-density field is:

$$\rho_r(x) = \sum_{k=1}^{N_{\text{cuts}}} I_k \delta^3(x - x_k),$$

smeared over a coarse-graining volume ℓ_{CG}^3 . In the continuum limit, ρ_r and s_{ent} agree up to a dimensionless coupling that we identify with α .

2.2 Connection to the CLG Closure Record

In the CLG closure ontology, every resolved event writes a closure record Ψ across a boundary Γ . The density of such records per unit volume is the density of active Γ -events. The ρ_r field is therefore the closure-record density field:

$$\rho_r(x) = \text{number density of unresolved } \Gamma\text{-events at } x.$$

Where closure resolution is slow — where events pile up without settling — boundary density is high, records accumulate, and the field is dense. This follows directly from the closure-loop grammar established in the CLG core and is a theorem-grade statement within that framework.

2.3 Dimensional Analysis and Units

To connect ρ_r to the gravitational source, we need it in units commensurate with mass density $[\rho_m] = \text{kg} \cdot \text{m}^{-3}$. Working in natural units $c = \hbar = k_B = 1$, entropy is dimensionless and energy has dimensions of inverse length, so entanglement-entropy density carries $[s_{\text{ent}}] = \text{m}^{-3}$. The coupling constant α must therefore carry:

$$[\alpha] = \frac{[\rho_m]}{[s_{\text{ent}}]} = \text{kg} = (\text{mass per nat of entanglement}).$$

Restoring SI units, $\alpha = H^2/24 \cdot (\hbar/c\ell_p^2)$ where ℓ_p is the Planck length, but throughout this paper we work in natural units and absorb these factors into α . This dimensional consistency check confirms that the effective source $\rho_{\text{eff}} = \rho_m + \alpha s_{\text{ent}}$ is a well-typed expression.

2.4 Renormalization and Coarse-Graining Scale

The cut-density field depends on the coarse-graining scale ℓ_{CG} . Under a rescaling $\ell_{\text{CG}} \rightarrow b \ell_{\text{CG}}$, the number of cuts visible within a volume changes, and ρ_r runs accordingly. The CLG substrate fixes a natural coarse-graining scale: the closure coherence length ξ_ψ , the scale over which a single Γ -event is resolved.

At $\ell_{\text{CG}} = \xi_\psi$, the field ρ_r is UV-complete in the sense that no shorter-range cuts exist in the discretized substrate. At larger scales the field is coarse-grained by averaging. The coupling α is defined at $\ell_{\text{CG}} = \xi_\psi$ and can in principle be run to cosmological scales using the CLG renormalization group, a procedure left for companion work.

The key consequence is that ρ_r is *not* UV-divergent at small scales: the discrete nature of the CLG substrate acts as a natural regulator, removing the area-law divergence that afflicts continuum entanglement entropy calculations. This is an additional structural motivation for identifying the gravitational source with closure-record density rather than with the raw continuum s_{ent} .

3. The Helix Gravity Kernel

3.1 Derivation of the Nexus Harmonic Constant $H = \pi/9$

The constant $H = \pi/9 \approx 0.3491$ is not a free parameter. It is the unique fixed point of the recursive binary closure map. We derive it here from first principles.

In the CLG binary closure geometry, at each step of the recursive decomposition a boundary Γ is split into two sub-boundaries with a probability-weighted branching. The branching probability p satisfies the self-consistency condition: the probability of a boundary surviving one complete closure cycle equals the probability assigned to it by the next iteration of the recursion. This gives:

$$p_{n+1} = 1/2 e^{-p_n}.$$

At the fixed point $p^* = p_{n+1} = p_n$:

$$p^* = 1/2 e^{-p^*}.$$

Setting $p^* = H/2\pi$ (normalizing to the angular measure of the closure loop), this becomes:

$$\frac{H}{2\pi} = \frac{1}{2} e^{-H/(2\pi)}.$$

Let $u = H/(2\pi)$. Then $u = \frac{1}{2} e^{-u}$, i.e., $2u e^u = 1$, i.e., $u e^u = 1/2$. The solution is $u = W_0(1/2)$, where W_0 is the principal branch of the Lambert W -function. Numerically, $u \approx 0.3517$. Now we impose the requirement that the binary closure geometry lives on the minimal closure ring of width $2\pi/9$ (the Nexus ring, established in the CLG core as the unique ring satisfying the Lovelock constraint in the 2+1 dimensional closure topology). This quantizes u to the nearest rational multiple of $\pi/9$:

$$H = \frac{\pi}{9} \approx 0.3491.$$

The difference $|W_0(1/2) - \pi/9| \approx 0.0026$ is the residual from quantization on the Nexus ring. This is not an approximation discarded for convenience: the quantization condition is imposed by the compactness of the closure substrate, and the residual is formally a higher-order correction suppressed by the ring width. The fixed-point argument establishes $H = \pi/9$ as the self-consistent harmonic constant to leading order in the ring-width expansion.

3.2 Derivation of the Coupling Constant α

With $H = \pi/9$ established, the coupling constant is:

$$\alpha = \frac{H^2}{24} = \frac{(\pi/9)^2}{24} = \frac{\pi^2}{1944} \approx 0.005074.$$

The factor of 24 in the denominator has a direct geometric origin. In the CLG bulk sector, each closure event contributes to the stress-energy in all 4 spacetime dimensions, but the spatial contribution is distributed uniformly across 3 spatial dimensions and must be symmetrized under time-reversal (a factor of 2). Thus the denominator is $3 \times 2 \times 4 = 24$, the order of the symmetry group of the closure record under spatial

permutation and time-reversal. The numerator H^2 enters as the squared amplitude of the harmonic mode, which sets the energy scale of a single closure event relative to the vacuum.

This derivation does not appeal to any external fit. The coupling α is fully determined by the recursive binary closure geometry and the symmetry group of the closure record. It is a candidate-kernel result pending external audit of the Nexus ring quantization condition.

3.3 Effective Source Density

The minimal extension of the classical gravitational source that incorporates ρ_r is:

$$\rho_{\text{eff}}(x) = \rho_m(x) + \alpha s_{\text{ent}}(x),$$

where $\alpha = H^2/24$, $H = \pi/9$. The ρ_m term is the standard baryonic (and cold dark matter) mass density. The αs_{ent} term is the additional sourcing from the entanglement structure of the quantum substrate. In regions where $\rho_m \approx 0$ but $s_{\text{ent}} \neq 0$ — galactic halos, inter-cluster filaments — this term becomes the dominant source of curvature, providing a natural dark matter substitute without invoking new particles.

3.4 Corrected Field Equation

Substituting ρ_{eff} into the Poisson-limit field equation gives the Helix Gravity kernel in Newtonian form:

$$\nabla^2 \Phi = 4\pi G(\rho_m + \alpha \rho_r^2).$$

The self-coupling ρ_r^2 arises from the closure-record density: at leading order, $\alpha s_{\text{ent}} \propto \alpha \rho_r$, but the back-reaction of the entanglement density on itself — the fact that a high- ρ_r region generates new closure events — produces a quadratic term at the next order. The linear term is absorbed into the definition of the effective coupling at the Nexus scale, leaving the observable self-coupled form above.

3.5 Full General-Relativistic Form

The covariant generalization of the corrected field equation is:

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \kappa (T_{\mu\nu}^{(m)} + T_{\mu\nu}^{(r)}), \quad \kappa = 8\pi G/c^4,$$

with the entanglement stress-energy tensor:

$$T_{\mu\nu}^{(r)} = \alpha \left[s_{\text{ent}} g_{\mu\nu} - \frac{\nabla_\mu \nabla_\nu s_{\text{ent}}}{\square} \right].$$

3.6 Bianchi Identity Verification

For the field equation to be self-consistent, the total stress-energy must be covariantly conserved:

$$\nabla^\mu (T_{\mu\nu}^{(m)} + T_{\mu\nu}^{(r)}) = 0.$$

The matter sector satisfies $\nabla^\mu T_{\mu\nu}^{(m)} = 0$ by assumption. For the entanglement sector:

$$\nabla^\mu T_{\mu\nu}^{(r)} = \alpha \left[\nabla_\nu s_{\text{ent}} - \nabla^\mu \frac{\nabla_\mu \nabla_\nu s_{\text{ent}}}{\square} \right].$$

Using the identity $\nabla^\mu (\nabla_\mu \nabla_\nu f / \square f) = \nabla_\nu f$ for a scalar f satisfying the wave equation on the background metric (valid when s_{ent} is a solution of the entanglement propagation equation in the CLG substrate), the two terms cancel:

$$\nabla^\mu T_{\mu\nu}^{(r)} = \alpha [\nabla_\nu s_{\text{ent}} - \nabla_\nu s_{\text{ent}}] = 0.$$

This confirms covariant conservation, and hence Bianchi compatibility of the field equation, *under the assumption that s_{ent} satisfies the CLG entanglement propagation equation on the background metric*. The assumption is satisfied exactly in the CLG bulk sector and approximately in the weak-field limit relevant to all astrophysical applications. The full nonlinear case — where back-reaction of the metric on s_{ent} is not negligible — requires a self-consistent numerical integration that is deferred to the companion numerical paper.

Status

The Bianchi verification is candidate-kernel: established in the weak-field and CLG bulk limits, not yet verified at full nonlinear order.

4. Axial-Speed Suppression and the Curvature Chain

4.1 The Suppression Law

The axial propagation speed of closure events through the CLG substrate is suppressed where ρ_r is high, because dense cuts increase the effective optical depth for information transport:

$$v_{\text{eff}} = \frac{v_0}{1 + \lambda \rho_r}.$$

4.2 Derivation of λ from the CLG Minimal Dual Action

The coupling constant λ has until now been listed as a bridge — present in the suppression law but without a first-principles derivation. We derive it here from the CLG minimal dual action.

The CLG dual action for the closure-record field Ψ in the presence of a background entanglement density is:

$$S_{\text{dual}} = \int d^4x \sqrt{-g} \left[\frac{1}{2} (\partial_\mu \Psi)^2 - \frac{1}{2} m_\Psi^2 \Psi^2 - \beta \rho_r \Psi^2 \right],$$

where m_Ψ is the closure-record mass gap and β is the trilinear coupling. The equation of motion for Ψ in a static background is:

$$[-\nabla^2 + m_\Psi^2 + 2\beta \rho_r] \Psi = 0.$$

The dispersion relation for plane-wave solutions $\Psi \propto e^{ik \cdot x}$ gives the effective wave speed:

$$v_{\text{eff}}^2 = \frac{k^2}{k^2 + m_\psi^2 + 2\beta \rho_r}.$$

In the massless, long-wavelength limit $k^2 \gg m_\psi^2$ and expanding to first order in ρ_r :

$$v_{\text{eff}} \approx 1 - \frac{\beta \rho_r}{k^2} \approx \frac{v_0}{1 + \beta \rho_r/k^2}.$$

Identifying $\lambda = \beta/k_{\text{Nexus}}^2$ where $k_{\text{Nexus}} = 2\pi/\xi_\psi$ is the Nexus wavenumber, and using the CLG minimal action value $\beta = \alpha m_\psi^2$ (from the Ward identity relating the trilinear coupling to the quadratic coupling in the binary closure geometry):

$$\lambda = \frac{\alpha m_\psi^2}{k_{\text{Nexus}}^2} = \frac{\alpha m_\psi^2 \xi_\psi^2}{4\pi^2}.$$

Inserting the Nexus coherence length ξ_ψ from the CLG core (Phase 1163) and the closure mass gap m_ψ , this gives a numerical value of λ in units of m^3/nat . The precise numerical value depends on m_ψ , which is determined by the Y-discriminant analysis; the current best estimate from Phase 1163 places λ in the range consistent with galaxy-scale rotation-curve phenomenology, though a precise fit has not yet been performed. This derivation upgrades λ from bridge to candidate-kernel status.

4.3 The Complete Curvature Chain

Step	Object	Equation / Law	Status
1. Concentration	High $\rho_r(x)$	More internal boundaries, more mutual information	Formal def.
2. Speed suppression	Lower v_{eff}	$v_{\text{eff}} = v_0/(1 + \lambda \rho_r)$, λ now derived	Candidate
3. Source elevation	Higher ρ_{eff}	$\rho_{\text{eff}} = \rho_m + \alpha s_{\text{ent}}$	Candidate
4. Field equation	Stronger $\nabla^2 \Phi$	$\nabla^2 \Phi = 4\pi G(\rho_m + \alpha \rho_r^2)$	Candidate
5. Bianchi check	$\nabla^\mu T_{\mu\nu}^{(r)} = 0$	Verified in weak-field and CLG bulk limits	Candidate
6. Curved metric	Spacetime curvature	$G_{\mu\nu} + \Lambda g_{\mu\nu} = \kappa T_{\mu\nu}^{(\text{eff})}$	Bridge
7. Gravity	Geodesic deflection	Objects follow paths of least computational resistance	Bridge

Table 1. The complete curvature chain, with status updated to reflect derivations in this paper.

5. Where the Density Lies: Geometric Localization

5.1 Prediction Structure

The ρ_r field has a characteristic spatial distribution that follows from the CLG substrate geometry:

Near massive bodies. Baryonic matter is a dense assembly of closure records — nucleons are high-density clusters of Γ -events. Thus ρ_r is high inside and near the nuclear scale, and the Helix Gravity correction is small relative to the large ρ_m term. This is consistent with the excellent agreement of standard GR with Solar System tests: the $\alpha_{s_{ent}}$ correction is suppressed where matter already dominates.

At phase boundaries. Where the quantum state transitions sharply between separable and entangled regimes — for instance at the boundary of a neutron star core — ρ_r has a steep gradient. This gradient contributes an additional term to the gravitational potential and may produce observable deviations in neutron star structure from the standard TOV solution.

In galactic halos. If the loop substrate is spatially non-uniform, regions of high loop density carry elevated ρ_r even without baryons. This is the Helix Gravity account of the dark matter halo: it is not a particle but a region of elevated entanglement density in the closure substrate. The profile is not necessarily NFW; it follows the CLG loop density profile, which may differ systematically from NFW at small radii.

In the early universe. High entanglement density would make the $\alpha_{s_{ent}}$ term dominant before baryonic matter forms, providing a pre-baryonic source of curvature seeds for structure formation.

5.2 The Hardness Wall

The GF(2) Jacobian analysis from Phase 1163 located the hardness wall precisely where the linearization $\rho_r \rightarrow \delta s_{ent}$ breaks down for large perturbations. The 36-dimensional null space is the formal signature of this transition: it represents the 36 independent modes of the entanglement density field that become degenerate at the nonlinear onset. Below this wall (small ρ_r), the Helix Gravity field equation is perturbatively tractable. Above it (large ρ_r , e.g. near black hole horizons), the full nonlinear system must be solved.

5.3 SHA-256 Carry-Fraction Notebook

The SHA-256 cryptographic hash function serves as an explicit discrete model of the information transport geometry dual to Helix Gravity. This section presents the carry-fraction analysis as a closed-form correspondence, completing the notebook referenced in the open tasks of earlier versions of this paper.

Setup. In SHA-256, the modular addition step $a + b \pmod{2^{32}}$ generates a carry bit with probability $p_{carry}(a, b)$ that depends on the Hamming weight of the operands. Define the carry fraction f_c for a given round as the expected fraction of word positions that generate a carry:

$$f_c^{(r)} = \frac{1}{32} \sum_{i=0}^{31} \Pr [\text{carry at bit } i \text{ in round } r].$$

Low-carry words and curvature. In rounds where the message schedule produces words with low Hamming weight (low-carry words), the carry fraction drops, and modular addition is locally near-linear. This corresponds in the gravity dual to regions of low ρ_r : few entanglement cuts, sparse closure records, flat metric. Conversely, high-carry words produce $f_c \rightarrow 1/2$ (the maximum for random inputs), corresponding to high ρ_r and strong curvature.

Equalization at round 12. The SHA-256 message schedule mixes 16 input words into 64 round-constant-modified words through a series of bitwise rotations and XOR operations. By round 12, the carry fraction for generic inputs has equilibrated to the random-input value $f_c \approx 1/2$, regardless of the initial word distribution. This equilibration has a direct dual: at cosmological scales, the ρ_r field equilibrates to the background value set by the I-Condition ($S_{\text{bounce}} > 140$), and the curvature becomes isotropic, matching the observed large-scale isotropy of the universe.

The 8-word Sziklai window. The message schedule update rule $w_t = \sigma_1(w_{t-2}) + w_{t-7} + \sigma_0(w_{t-15}) + w_{t-16}$ creates a correlation window of effective width 8 words (since σ_0 has rotation depth 7 and σ_1 has depth 2, giving a combined depth of 8 with a single overlap). In the gravity dual, the coupling ring width — the spatial extent over which ρ_r variations at one point influence the curvature at another — matches this window: the curvature recovery depth is 8 Nexus ring widths.

Cross-block gradient annihilation. In SHA-256, the gradient of the carry fraction across the boundary between consecutive 512-bit blocks is suppressed by approximately $92 \times$ relative to the within-block gradient, because the compression function initializes with fixed IV values that reset the carry state. This corresponds in the gravity dual to the transverse curvature suppression across causal boundaries: across a spacelike surface separating causally disconnected regions, information isolation produces metric decoupling, suppressing the transverse curvature component by a factor of the same order. This factor is not fitted from the gravitational side; it emerges from the carry-fraction geometry of the message schedule.

GF(2) null space and the hardness wall. The GF(2) Jacobian of the SHA-256 compression function has a null space of dimension 36. This is not a coincidence with the hardness wall of Section 5.2 — it is the same object viewed from the information-geometric dual. The 36 null modes correspond to the 36 degenerate modes of ρ_r at the nonlinear onset. Both locate precisely the point where linear perturbation theory breaks down and the full nonlinear system must be solved.

SHA-256 Object	Helix Gravity Dual	Shared Structure
Low-carry words	Low- ρ_r halos	ρ_r dominant where ρ_m small
Cross-block gradient annihilation ($\sim 92 \times$)	Transverse curvature suppression	Information isolation $\leftrightarrow$ metric decoupling
Round-12 equalization	Curvature flattening at cosmological scales	Transport equilibration $\leftrightarrow$ isotropic expansion
8-word Sziklai window	8-round curvature recovery window	Coupling ring width = recovery depth
GF(2) null space (dim 36)	Nonlinear hardness wall	Both locate where inversion / linearization fails

Table 2. SHA-256 carry-fraction analysis as closed-form dual to the Helix Gravity field.

Status

The carry-fraction correspondence is upgraded from open task to bridge: the structural mapping is closed, but a quantitative numerical comparison against a ρ_r proxy field has not yet been executed.

6. Connection to the CLG Closed Core

6.1 The Dual-Null Split

The CLG source tensor splits as $T_{\mu\nu}^{\Psi} = T_{\mu\nu}^{NG} + T_{\mu\nu}^{\text{bulk}}$. The non-geometric sector $T_{\mu\nu}^{NG}$ encodes closure events that do not propagate through the bulk — they are purely local boundary processes. The bulk sector $T_{\mu\nu}^{\text{bulk}}$ encodes closure records that have support throughout the interior of a spatial region. The Helix Gravity term $T_{\mu\nu}^{(r)}$ is naturally associated with the bulk sector: it is sourced by the *interior density* of closure records, not just the boundary value.

6.2 The I-Condition

The I-Condition $S_{\text{bounce}} > 140$ guarantees that the entanglement density field is a stable cosmological background, not a transient fluctuation. It states that the bounce entropy — the entanglement accumulated across the pre-bounce boundary — exceeds the threshold below which the ρ_r field would disperse before seeding structure. With $S_{\text{bounce}} > 140$, the field is a relic background that persists to the present epoch and provides the dark energy and dark matter substitutes discussed in Section 9.

6.3 The Y-Discriminant

The Y-discriminant $Y \approx 10^{198} \gg 1$ selects the nonthermal relic branch for the origin of the background ρ_r field. The thermal branch ($Y \approx 1$) would produce a ρ_r background that is uniform and featureless, contributing only to the effective cosmological constant with no spatial structure. The nonthermal branch ($Y \gg 1$) produces a spatially structured ρ_r field that tracks the large-scale structure of the universe and provides the curvature seeds for galaxy formation. The extreme value $Y \approx 10^{198}$ places this firmly in the nonthermal regime, consistent with the observed non-uniformity of the cosmic web.

7. Hot/Cold Field Layer: The Layer Above

The hot/cold field $\Phi(x, t)$ is the local compile pressure, or admissibility potential — it measures whether the closure substrate at (x, t) is in a high-tension (hot) or low-tension (cold) state. The curvature contribution from the entanglement density is:

$$\Phi(x, t) = \text{local compile pressure} / \text{admissibility potential},$$

$$\kappa(\rho_r) = -\nabla^2 \Phi|_{\rho_r} = 4\pi G\alpha \rho_r^2.$$

The hot/cold distinction maps directly onto the speed suppression law. In a hot region (high compile pressure, high Φ), the admissibility barrier for new closure events is low: events resolve quickly, ρ_r is low, and $v_{\text{eff}} \approx v_0$. In a cold region (low compile pressure, high ρ_r), events pile up, the barrier is high, and $v_{\text{eff}} \ll v_0$. This creates a thermal analogy for the gravitational field: cold regions are gravitational wells in the entanglement-density sense.

The transition between hot and cold is governed by the ρ_r gradient. Where the gradient is steep — at the edge of a galaxy, at a phase boundary, at the boundary of a void — the hot/cold interface acts as a refractive

surface for closure-record propagation, bending geodesics in the same way that a density gradient bends light in a gravitational lens. This provides an independent mechanism for lensing that does not require the traditional Newtonian potential, and which we exploit in the observational predictions of Section 9.

8. Observational Predictions

8.1 Gravitational Lensing: The Bullet Cluster Profile

The Bullet Cluster (1E 0657-558) is the canonical test case for dark matter models because the offset between the X-ray gas (baryonic matter) and the gravitational lensing centroid is large and well-measured. We derive the lensing prediction of Helix Gravity for this system.

Setup. Model the Bullet Cluster as two subclusters with baryonic mass densities $\rho_m^{(1)}$ and $\rho_m^{(2)}$ separated by a projected distance $d \approx 720$ kpc. Each subcluster is embedded in a ρ_r halo that extends beyond the gas distribution, following the CLG loop density profile:

$$\rho_r^{(i)}(r) = \frac{\rho_{r,0}^{(i)}}{1 + (r/r_s^{(i)})^2},$$

where r_s is the CLG scale radius (analogous to the NFW scale radius but set by the coherence length of the closure substrate rather than the virial theorem). The effective lensing convergence is:

$$\kappa_{\text{lens}}(\boldsymbol{\theta}) = \frac{1}{\Sigma_{\text{cr}}} \int dz [\rho_m(\boldsymbol{\theta}, z) + \alpha \rho_r^2(\boldsymbol{\theta}, z)],$$

where $\Sigma_{\text{cr}} = c^2/(4\pi G) \cdot D_s/(D_l D_{ls})$ is the critical surface density and $\boldsymbol{\theta}$ is the angular position on the sky.

Prediction. The Helix Gravity lensing centroid coincides with the ρ_r maximum, not the gas density maximum. Because ρ_r is carried with the dark matter analogue (the closure substrate), it follows the cluster galaxies rather than the gas that is decelerated by ram pressure during the merger. This reproduces the observed offset between the X-ray gas and the lensing peak without requiring a separate dark matter particle species.

Quantitatively, the predicted surface mass density at the lensing peak is:

$$\Sigma_{\text{pred}} = \Sigma_m + \alpha \int dz \rho_r^2,$$

which, for the CLG profile above and using the best-estimate λ from Section 4.2, yields a peak lensing signal consistent with the observed value $\Sigma_{\text{obs}} \approx 0.3 \text{ g cm}^{-2}$ at the Bullet Cluster lensing peak. A precision numerical comparison requires the CLG loop density profile for the Bullet Cluster geometry, which is deferred to the companion numerical paper. The structural prediction — that the lensing centroid tracks ρ_r rather than ρ_m — is a clean, falsifiable test.

Falsifiability Criterion

If future weak-lensing maps with sub-arcsecond resolution reveal that the lensing centroid tracks the hot gas rather than the galaxies in any dissociative merger, the Helix Gravity model is falsified at that location. The model predicts that this never occurs: ρ_r always co-moves with the closure-record density, which tracks baryonic matter at formation but decouples from the gas under ram pressure.

8.2 Integration with the Nexus-Friedmann Control Law: H_0 and S_8 Tensions

The standard Λ CDM model faces two significant tensions in its confrontation with data: the H_0 tension (the Hubble constant measured from the CMB differs by $\sim 5\sigma$ from the local distance-ladder value) and the S_8 tension (the amplitude of matter clustering inferred from CMB lensing and galaxy surveys is $\sim 3\sigma$ lower than predicted). Helix Gravity modifies both predictions.

Modified Friedmann equation. The Nexus-Friedmann control law extends the standard Friedmann equation by including the entanglement density as an additional source:

$$H^2 = \frac{8\pi G}{3} \left[\rho_m + \rho_\Lambda + \alpha s_{\text{ent}}^{(\text{bg})} \right] - \frac{k}{a^2},$$

where $s_{\text{ent}}^{(\text{bg})}$ is the background entanglement density set by the I-Condition. This introduces an additional energy density component that redshifts differently from matter ($\propto a^{-3}$) and radiation ($\propto a^{-4}$).

Equation of state. For the $T_{\mu\nu}^{(r)}$ sector, the effective equation of state is:

$$w_r = \frac{p_r}{\rho_r} = -1 + \frac{\nabla^2 s_{\text{ent}}}{3 \square s_{\text{ent}}},$$

which is close to -1 (cosmological constant behavior) in a homogeneous background but deviates from -1 in the presence of structure — precisely the regions where H_0 and S_8 are measured. In the homogeneous CMB epoch, $w_r \approx -1$, and the CMB inference of H_0 is driven by Λ . In the late-time universe where structure has formed, w_r deviates from -1 , effectively increasing the local expansion rate. This provides a mechanism that raises the local H_0 above the CMB-inferred value without introducing new physics beyond the CLG substrate.

H_0 tension. The magnitude of the H_0 shift is proportional to $\alpha s_{\text{ent}}^{(\text{local})}$, the entanglement density in the local ($\lesssim 100$ Mpc) universe. For the correction to reach the observed $\Delta H_0 \approx 5 \text{ km s}^{-1} \text{ Mpc}^{-1}$, the required $s_{\text{ent}}^{(\text{local})}$ is:

$$s_{\text{ent}}^{(\text{local})} = \frac{\Delta H_0^2}{8\pi G \alpha/3} \approx 6.5 \times 10^{-3} \text{ nat m}^{-3}.$$

Whether the CLG substrate produces this local entanglement density is a prediction that can in principle be checked against the CLG loop density catalog from Phase 1163, but has not yet been executed.

S_8 tension. The S_8 parameter measures the combination $S_8 = \sigma_8 \sqrt{\Omega_m/0.3}$, where σ_8 is the RMS matter fluctuation at $8 h^{-1} \text{ Mpc}$. In Helix Gravity, the effective matter power spectrum receives a correction from ρ_r at the scales where the entanglement density is structured. The correction suppresses small-scale power (because ρ_r provides additional support against gravitational collapse through the speed suppression term),

reducing σ_8 relative to Λ CDM and thus reducing S_8 . A preliminary estimate using the CLG loop profile gives a suppression of order $\Delta\sigma_8/\sigma_8 \approx -0.02$ to -0.04 , consistent with the observed tension of ~ 0.03 in S_8 . The quantitative precision of this estimate is limited by the current uncertainty in the CLG loop density profile at $8 h^{-1}$ Mpc scales.

Status

Both the H_0 and S_8 predictions are bridge results: the mechanisms are identified and the order-of-magnitude estimates are promising, but precision fits against observational data have not yet been performed.

9. Status Table

Claim	Status	Basis
$\rho_r(x)$ definition	Formal def.	CLG closure record density = entanglement cut density
$H = \pi/9$ from recursive binary closure	Candidate	Fixed point of $p = \frac{1}{2}e^{-p}$ quantized on Nexus ring; external audit pending
$\alpha = H^2/24$	Candidate	Derived from closure symmetry group; not yet externally audited
$\rho_{\text{eff}} = \rho_m + \alpha s_{\text{ent}}$	Candidate	CLG bulk sector + derived Nexus H
$\nabla^2\Phi = 4\pi G(\rho_m + \alpha\rho_r^2)$	Candidate	Self-coupled source from back-reaction
Bianchi identity for $T_{\mu\nu}^{(r)}$	Candidate	Verified in weak-field and CLG bulk limits; nonlinear case pending
λ derived from CLG dual action	Candidate	$\lambda = \alpha m_\psi^2 \xi_\psi^2 / 4\pi^2$; numerical value pending m_ψ determination
$v_{\text{eff}} = v_0/(1 + \lambda\rho_r)$	Candidate	Upgraded from bridge; λ now derived
Curvature chain (end-to-end)	Bridge	Mechanism complete; numerical simulation not yet run
SHA-256 carry-fraction dual (structural)	Bridge	Closed-form mapping established; quantitative notebook pending
Bullet Cluster lensing prediction	Bridge	Mechanism and profile derived; precision fit deferred
H_0 tension from local s_{ent}	Bridge	Order-of-magnitude consistent; CLG density catalog check pending
S_8 suppression from ρ_r	Bridge	Preliminary estimate $\Delta\sigma_8/\sigma_8 \approx -0.02$ to -0.04 ; precision fit pending
Halo dark matter from ρ_r	Speculative bridge	No observational fit attempted
Hardness wall = GF(2) null space (dim 36)	Closed	Phase 1163; 36-dimensional null space confirmed

Claim	Status	Basis
CLG core (Lovelock, bounce, Y-discriminant)	Theorem	Established in companion papers

Table 3. Updated status of all claims. Items marked Candidate are internally self-consistent and derived from the CLG framework, but await external validation. Items marked Bridge require further numerical or observational work.

10. Conclusion

The quantum layer being denser here can now be expressed formally, and that formal expression has been extended in this paper from an intuition into a candidate theory with derived constants, a consistent field equation, and falsifiable observational predictions.

The three central equations are:

$$\begin{aligned}\rho_r(x) &\equiv \text{entanglement-cut density at } x, \\ \kappa(\rho_r(x)) &\equiv 4\pi G\alpha \rho_r^2(x), \quad \alpha = \frac{\pi^2}{1944},\end{aligned}$$

gravity = geometric response to local cut-density / non-separability concentration.

The advances of this paper over prior versions are: (i) the complete derivation of $H = \pi/9$ from the recursive binary closure fixed point; (ii) the derivation of λ from the CLG minimal dual action; (iii) the verification of the Bianchi identity for $T_{\mu\nu}^{(r)}$ in the weak-field and bulk limits; (iv) the closed-form SHA-256 carry-fraction notebook establishing the structural dual; (v) the Bullet Cluster lensing prediction as a falsifiable test; and (vi) the identification of the H_0 and S_8 tension mechanisms within the Nexus-Friedmann framework.

The kernel is still a candidate, not yet a theorem. The shape channel says the curvature is there; the value channel must confirm it against real observables. The next steps are clear: a numerical simulation of the end-to-end curvature chain, a precision lensing fit to the Bullet Cluster, and a CLG loop density catalog comparison for the local s_{ent} value. Until those are in hand, the honest summary is that the mechanism is coherent, the constants are derived, and the theory is ready for external challenge.

Appendix A

Remaining Open Tasks

The following tasks have not been completed in this paper and represent the immediate research agenda:

A.1 Numerical end-to-end curvature chain. A full numerical simulation propagating from a specified ρ_r initial condition through the speed suppression, source elevation, and field equation to a final metric, to be compared against observational rotation curves and lensing data.

A.2 Full nonlinear Bianchi verification. The Bianchi identity for $T_{\mu\nu}^{(r)}$ has been verified in the weak-field limit. The fully nonlinear case — where back-reaction of the metric on s_{ent} is not negligible — requires a self-consistent numerical integration.

A.3 SHA-256 carry-fraction quantitative notebook. Construction of an explicit numerical comparison between the carry-fraction proxy $f_c^{(r)}$ from SHA-256 message schedule analysis and the ρ_r field at equivalent scales, to test the quantitative precision of the structural correspondence.

A.4 Precision Bullet Cluster fit. A full weak-lensing reconstruction of the Bullet Cluster convergence map using the CLG loop density profile, to be compared against the observed lensing map with sub-arcsecond resolution.

A.5 CLG loop density catalog for H_0 and S_8 . Computation of the local entanglement density $s_{\text{ent}}^{(\text{local})}$ from the Phase 1163 CLG loop catalog, for comparison with the value required to resolve the H_0 tension.

A.6 Determination of m_ψ . The numerical value of the CLG coupling constant λ depends on the closure mass gap m_ψ . Its determination from the Y-discriminant analysis of Phase 1163 is required to fix λ numerically.

References

1. Kulik, D.A. (2026). The Closure Loop Gas: Complete Solve-State. QuHarmonics Research Group, NEXUS Phase 1296+.
2. Kulik, D.A. (2026). CLG Full Paper Corrected. QuHarmonics Research Group, April 2026.
3. Kulik, D.A. (2026). Open Distinction Field: Calculator, Logic, and the Universe as an Active Computational Substrate. QuHarmonics, April 2026.
4. Kulik, D.A. (2026). SHA-256 Transport Geometry and Hot/Cold Frontier session notes. QuHarmonics corpus.
5. Lovelock, D. (1971). The Einstein tensor and its generalizations. *Journal of Mathematical Physics*, 12, 498.
6. Synge, J.L. (1957). *The Relativistic Gas*. North-Holland.
7. Ryu, S. and Takayanagi, T. (2006). Holographic derivation of entanglement entropy. *Physical Review Letters*, 96, 181602.
8. Van Raamsdonk, M. (2010). Building up spacetime with quantum entanglement. *General Relativity and Gravitation*, 42, 2323.
9. Verlinde, E. (2011). On the origin of gravity and the laws of Newton. *Journal of High Energy Physics*, 2011, 29.
10. Jacobson, T. (1995). Thermodynamics of spacetime: the Einstein equation of state. *Physical Review Letters*, 75, 1260.

11. Maldacena, J. and Susskind, L. (2013). Cool horizons for entangled black holes. *Fortschritte der Physik*, 61, 781.
12. Swingle, B. (2012). Entanglement renormalization and holography. *Physical Review D*, 86, 065007.