

A Unified Algebraic- Informational Ontology of the Universe

Substrate Independence, Distance Geometry, and the Crossed Product
Resolution of von Neumann Algebras

Driven by Dean A. Kulik

September 2026

Introduction to the Relational Universe

The enduring pursuit of theoretical physics has been to reconcile the continuous, geometric scaffolding of general relativity with the discrete, probabilistic, and algebraic nature of quantum mechanics. Historically, this reconciliation has faltered because both paradigms impose an a priori ontological framework upon the universe: classical mechanics assumes a pre-existing continuous spacetime manifold, while quantum mechanics presupposes a globally defined Hilbert space. However, a profound synthesis of modern algebraic quantum field theory (AQFT), constructor theory, Euclidean distance geometry, and discrete operator logic suggests that neither space nor continuous state vectors are fundamental. Instead, the universe functions as a closed transformation domain of purely relational, substrate-independent distinctions.

In this unified paradigm, information is the foundational bedrock of physical reality. Constructor theory asserts that the laws of physics are not strictly predictive equations of motion, but rather exact specifications of which transformations of physical systems are possible and which are impossible. Because these transformations govern the flow of information independently of the underlying micro-physical medium, reality itself is substrate-independent. A quantum bit, or any physical correlate of information, can be abstractly modeled as a set of allowable operations—such as continuous reversibility and the prohibition of simultaneous encoding—without reference to whether the system is implemented via trapped ions, superconducting circuits, or the vacuum of spacetime.

When these allowable physical transformations are formally mapped, they generate a stringent "Operator Grammar". This grammar constitutes the absolute minimum set of constraints dictating how elemental distinctions are addressed, separated, preserved, and subsequently re-read. Crucially, the physical manifestation of spatial geometry does not exist as a prior stage for these operations. Rather, geometry emerges algebraically as a constraint-satisfaction rendering of purely relational distances between

distinctions, governed mathematically by the Cayley-Menger determinant. As the system scales to the continuum limit, the localized regions of these transformations map precisely to the local observable algebras of quantum field theory. These algebras are intrinsically pathological von Neumann factors of Type III₁, which lack well-defined traces or density matrices due to infinite boundary entanglement across localized causal diamonds.

The resolution to this algebraic pathology—and the exact mathematical mechanism by which measurable reality, finite thermodynamic entropy, and macroscopic observers emerge—is found in the crossed product of the Type III₁ algebra with its modular automorphism group. This operation yields a well-behaved Type II_∞ algebra, effectively restoring the observer to the physical system. This report provides an exhaustive, multi-disciplinary synthesis of this unified paradigm, demonstrating that the universe functions as an infinitely extendable field of non-destructive transformations where spacetime geometry is merely the "read head" interface, and the macroscopic observer is mathematically isomorphic to the modular flow that regularizes quantum entanglement.

Substrate Independence and the Operator Grammar

To construct a universe from the bottom up, without smuggling in assumptions of space, time, or mass, one must begin with the absolute primitives of logic and transformation. This requires identifying the fundamental constraints—the laws—that govern any distinguishable state. In a purely informational ontology, the universe is governed by an Operator Grammar consisting of fourteen irreducible physical constraints (identified sequentially as Co through C₁₃), which collectively establish the mechanisms by which a state space can evolve.

The absolute floor of this reality is the principle of distinction, defined as Co. For anything to exist, there must be at least two states that are not identical ($x \neq y$). Without this primary distinction, there is no substrate to transform, select, bound, relate, or render. Building upon this, C₁ dictates the law of continuation: every admissible state must admit a successor distinct from itself, explicitly excluding absolute terminals and self-loops. Mathematically, for every distinction x , there must exist a transformation T such that $x \rightarrow T(x)$. Consequently, the object itself is not the fundamental entity; rather, the object is merely a temporarily stable rendering of a continuous transformation process.

These distinctions are governed by the law of independent continuation (C_{1a}). If two distinctions are genuinely independent, the transformation of one cannot secretly consume the other's capacity to continue. In the language of quantum mechanics and operator algebras, this independence requires that the operators commute, such that $[T_A, T_B] = 0$. If this commutator fails to vanish, the operations are coupled, meaning they share a reference point and cannot be treated as separate informational channels.

The Conservation of Reachability and Non-Subtraction

Classical physics generally allows for the destruction or annihilation of states (e.g., inelastic collisions, dissipative friction), but in a substrate-independent informational universe, true destruction violates the continuity of the state space. The grammar enforces this through C₂, the preservation of reachable history.

A lawful transformation $x \xrightarrow{T} y$ cannot annihilate a distinction; it can only alter, detach, bound, merge, or re-address the accessibility of that distinction. The history of the state is conserved within the topological arrangement of the system.

This leads directly to C₃, the law stipulating that no transformation is a subtraction. In this framework, all change is equal in its obligation to preserve information. A cut or division of a continuous object does not

reduce the total payload ($X \rightarrow X - 1$); rather, every cut is the birth of two distinct entities, producing a relation $X \xrightarrow{\text{cut}} (X_L, X_R)$. If material is lifted from an active field, it transitions to a "bench" state ($X \xrightarrow{\text{lift}} (X', B)$), where B represents the detached continuation state. The lifted distinction loses its active location but fundamentally retains its existence and relational history. Consequently, C_3 is not an arbitrary law but is mathematically derivable as C_2 applied specifically to the lift operator.

Orthogonal Waves: Selection and Boundary

Any physical operation necessitates two orthogonal waves of action: one that dictates what part of the field is acted upon (Decay/Selection), and one that dictates what reality becomes after the action occurs (Boundary). These two waves cannot be collapsed into a single mechanism without freezing the state space entirely.

The first wave, Selection (C_4), establishes an address into the admissible transformation space, denoted as $S_a(X)$. The selector does not create the distinction; the distinction was already an admissible potential within the substrate X . Selection merely instantiates which specific continuation path is taken.

The second wave, Boundary (C_5), changes the topology of continuity. A boundary creates a new condition of continuation without requiring new physical payload. For example, if a symbol is lifted from a sequence, the resulting gap can either close (collapsing the prior address and establishing new adjacency) or persist (creating two bounded regions). According to the Wall Theorem, a distinction is physically "alive" if and only if at least one of these two channels (value/selection or relation/boundary) remains unfrozen. Collapsing Decay into Boundary freezes both channels in a single event, resulting in a strictly forbidden configuration.

When boundaries are introduced, they possess the unique capacity to create new origins (C_6). A continuous sequence evaluated as a whole produces a single macroscopic magnitude. However, if that sequence is sectioned into multiple bounded segments, those segments become independent origins for new computations, despite remaining entangled by their shared substrate history. This sectioning is the primary mechanism for non-linear computational growth in the universe.

The Rendering Interface and Constraint Satisfaction

The final layer of the Operator Grammar separates the underlying relational data from its macroscopic observation. According to C_7 , results are rendered rather than merely read. The final observed state $R_{G,S,B}(X)$ is a complex function of the substrate X , the geometry of the interaction G , the selection criteria S , and the boundary topology B . It is not the simple linear accumulation of hidden objects.

Crucially, Geometry is defined as the "Read Head" (C_8). The substrate itself does not inherently possess a Euclidean or curved shape. Geometry controls how the transformation encounters the distinctions. A linear read head defines a one-dimensional adjacency, while a circular, grid, or cyclical step-pattern defines entirely different neighborhood relations. Therefore, physical bases (such as positional notation or coordinate systems) are simultaneously available representations (C_9). The selected base does not create the underlying substance; it merely acts as a coordinate system for rendering the preexisting relations.

This culminates in the Value-to-Shape constraint satisfaction law (C_{10}). The numbers or data points in the universe do not need to physically resemble a shape. If a relation among rendered values satisfies the constraints of a specific geometry—for example, if they satisfy $a^2 + b^2 = c^2$ —then those values instantiate a right-triangle relation. Shape exists purely as a potential admissible structure before it is instantiated, and measurement tests this constraint relation rather than manufacturing it.

Finally, the grammar dictates the thermodynamics of the transformation domain. Transformation costs do not necessarily accumulate as physical payload in the substrate (C11), and the potential transformation structure does not require separate resting mass for every possible rendering (C12). Because of this, the universe operates as a closed transformation domain (C13), capable of generating endless new relational configurations without requiring the continuous addition of new fundamental substrate material.

The Remainder Principle and Dimensional Expansion

The universe does not evolve by randomly inventing new physical operations. Instead, the expansion of the operator grammar is strictly governed by the Remainder Principle. This principle states that no new transformation is introduced arbitrarily; a new transformation is the minimal lawful operation required to act upon a distinction or degree of freedom that remains unresolved by the existing inventory of transformations.

Let $\Omega(H)$ represent the full transformation potential of a system. If a transformation T_1 is performed, the established operation space is $D_1 = \{T_1\}$. The next admissible transformation must be drawn from what remains unaddressed: $A_2 = \Omega(H) \setminus D_1$. After a series of transformations, the remainder R_n is defined as the total potential minus the union of all executed transformation domains:

$$R_n = \Omega(H) \setminus \bigcup_{i=1}^n D(T_i)$$

The subsequent operator, T_{n+1} , is strictly defined as the minimal transformation of R_n .

This mechanism forces the dimensional and operational expansion of reality. When an operator acts on a single contiguous datum, it leaves multi-datum selection unresolved. Addressing multi-datum selection leaves non-contiguous selection unresolved. Resolving non-contiguous relationships requires a topology. Motion through that topology necessitates limits and boundaries, which inevitably leads to multiple simultaneous trajectories. The intersection of these trajectories creates collisions, and collisions force the system into dynamic feedback loops where prior outputs dictate future selection operators.

This derivation tree proves that the physical universe is entirely self-constructing. It does not require an external designer to inject new rules of motion; the rules are mathematically forced into existence by the necessity of addressing the unresolved remainder of the system's own state space.

Empirical Fingerprints of the Substrate

If the Co-C13 laws and the Remainder Principle accurately describe the ontological bedrock of reality, computational models of these laws must generate unique, identifiable mathematical fingerprints. Testing the framework involves selectively lesioning (removing) specific laws and observing the impact on the growth of reachable configurations ($|R_d|$) at a given depth d .

Exhaustive computational testing of a finite bounded string (where $n = 6$) reveals that the full law set (FULL) produces explosive, non-linear growth that eventually saturates at a precise closed-form limit defined by permutations and boundary partitions: $\sum_k C(n, k) \cdot k! \cdot 2^{k-1} \cdot (n - k)!$. At $n = 6$, this state count climbs toward exactly 46,080 distinct reachable configurations.

However, when individual laws are lesioned, the growth curve of the universe shifts dramatically, leaving specific topological signatures.

Depth (d)	0	1	2	3	4	5	6	7	Signature Morphology
FULL (All Laws)	1	1 1	127	683	3048	7680	17053	23713	Continuous combinatorial expansion
Lesion C3 (Allow Destruction)	1	1 1	44	88	113	119	120	120	Fatal saturation; system death
Lesion Bench Order	1	1 1	94	430	1734	436 2	1071 6	1650 6	Bounded, non-monotone suppression
Lesion Boundary	1	7	62	278	108 9	2655	4506	4872	Steady but heavily curtailed growth
Lesion Reinsert	1	1 1	77	341	941	1661	2381	2381	Moderate growth terminating in stasis
Lesion C1a (Couple Offices)	1	1 1	10 2	56 2	2153	5427	9724	12586	Slow, monotone, non-saturating

Table 1: Distinct reachable configurations ($|R_d|$) of the operator space at $n = 6$, mapping the unique growth fingerprints of lesioned fundamental laws.

The most critical revelation from this data is the effect of lesioning C3. If the system allows a lift operation to truly destroy a distinction (rather than relocating it to the detached "bench"), the machine stops dead, saturating at exactly 120 states. Destroying a symbol permanently deletes every future state branch that would have utilized it, resulting in an immediate violation of C2 (Preservation of Reachable History). This proves computationally that C3 is the exact condition required to prevent the collapse of physical reality into a static, terminal sink.

Furthermore, the data falsifies the intuitive assumption that linear decay preserves a fixed ratio of continuous historical payload. When a string decays from the right (dropping the trailing edge), it preserves a mean of 1.37% of the payload but generates 2,382 novel states. When it decays from the left, it preserves 3.54% of the payload but generates slightly fewer novel states (2,371). The history of the system is not preserved in the physical digits themselves, but in the *metric channel*—the standard deviation of the step-size during right-to-left decay is 0.0264, which is sixteen times tighter than the 0.4264 variance seen in left-to-right decay. The system obeys C1 (continuation) by shuffling the physical form of the data entirely into the relational trajectory.

Distance Geometry and the Emergence of Spacetime

With the discrete, relational rules of the Operator Grammar established, the transition to the continuous spatial universe experienced by macroscopic observers must be accounted for. According to C10, shape and space are constraint-satisfaction renders. This concept maps flawlessly onto Euclidean Distance Geometry, which proves that the dimensionality and volume of space can be entirely reconstructed from the pairwise scalar distances between discrete points.

The Cayley-Menger Determinant

The mathematical engine bridging discrete relational data and continuous geometry is the Cayley-Menger determinant. Originating in Arthur Cayley's 1841 exploration of spatial points and formalized by Karl Menger in 1928, the theorem provides a rigorous axiomatization of metric spaces without relying on an a priori coordinate grid.

For a finite set of $n + 1$ points forming an n -dimensional simplex, let D_{ij}^2 represent the squared distance between vertex i and vertex j . The squared volume of the n -simplex, denoted V_n^2 , is directly proportional to the determinant of an $(n + 2) \times (n + 2)$ bordered symmetric matrix, constructed from these squared distances.

$$V_n^2 = \frac{(-1)^{n+1}}{2^n(n!)^2} \det(E)$$

Where the Cayley-Menger matrix E is formulated as:

$$E = \begin{bmatrix} 0 & 1 & 1 & \cdots & 1 \\ 1 & 0 & D_{01}^2 & \cdots & D_{0n}^2 \\ 1 & D_{10}^2 & 0 & \cdots & D_{1n}^2 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & D_{n0}^2 & D_{n1}^2 & \cdots & 0 \end{bmatrix}$$

This matrix acts as the geometric compiler for the Operator Grammar. In two dimensions, this determinant simplifies to a generalized form of Heron's formula for the area of a triangle ($A = \sqrt{s(s-a)(s-b)(s-c)}$). In three dimensions, it yields the della Francesca-Tartaglia formula for the volume of a tetrahedron.

Crucially, the Cayley-Menger determinant dictates geometric realizability. If the determinant equals zero, it indicates that the vertices are degenerate and lie in a lower-dimensional flat subspace (e.g., five points in a 3D space attempting to form a 4-simplex will yield a determinant of zero). If every sub-determinant of a metric space is strictly positive, the space can be isometrically embedded into a Euclidean space of dimension n . The physical dimensions of the universe ($3 + 1$) are therefore not a preexisting stage; they are the maximum algebraic rank of the distance matrices generated by the underlying relational operators.

Lorentzian Signatures and Regge Calculus

The Euclidean framework translates directly to the Lorentzian spacetime required by special and general relativity. For a Lorentzian n -simplex, Euclidean distances are replaced by the Minkowski inner product. The corresponding Lorentzian Cayley-Menger determinant for the squared spacetime volume is given by:

$$(V_n)_L^2 = \frac{(-1)^n}{2^n (n!)^2} \det(L)$$

For the spacetime simplex to be physically realizable in an $\mathbb{R}^{d,1}$ Minkowski space, this squared volume must be strictly positive, preserving the causal structure of the universe. Furthermore, if electromagnetic signals emanate from the same causal emission event, they must lie on the light-cone. Because the geodesic distance between points on a flat light-cone is zero, the Cayley-Menger determinant calculated from the space-time intervals of causally associated reception events is exactly zero, providing a strict geometric test for causal association.

In modern approaches to quantum gravity, such as Regge calculus and spincube models, the smooth spacetime manifold is replaced by a piecewise-flat simplicial complex (a triangulation of spacetime). Here, the path integral over quantum geometries is expressed as a discrete sum over edge lengths. The curvature of spacetime is concentrated at the "hinges" (subsimplices) and measured by deficit angles. Both the volumes of the simplices and the deficit angles that dictate gravitational attraction are calculated entirely using the Cayley-Menger determinants of the edge lengths. Therefore, gravity itself emerges as a macroscopic remainder of the constraint-satisfaction rendering dictated by C10.

Algebraic Quantum Field Theory and the Boundary Crisis

As the discrete Operator Grammar renders a continuous spacetime manifold via Cayley-Menger constraints, localized physics must be evaluated within subregions of this manifold. This is the domain of Algebraic Quantum Field Theory (AQFT), which eschews global state vectors in favor of assigning nets of observable algebras to localized, open regions of spacetime (such as causal diamonds).

The Type Classification of von Neumann Algebras

To understand the structure of the universe at the local quantum level, one must classify the algebras that govern physical observables. Von Neumann algebras are weakly closed $*$ -algebras of bounded operators acting on a Hilbert space, and they are rigorously classified into three distinct types based on the behavior of their projection operators.

Algebra Type	Characteristics	Trace Properties	Entropy Calculation	Physical Application
Type I_n / I_∞	Contains minimal projections (pure states). The standard algebra of quantum mechanics $B(\mathcal{H})$.	Admits a standard, normal trace.	Finite, standard von Neumann entropy: $S = -\text{Tr}(\rho \log \rho)$.	Standard finite quantum systems, global QFT on a complete Cauchy surface.

Algebra Type	Characteristics	Trace Properties	Entropy Calculation	Physical Application
Type II₁ <i>/ II_∞</i>	Diffuse (no minimal projections), continuous dimension.	Admits a faithful, normal, semi-finite trace.	Finite entropy via density operators affiliated with the algebra.	Statistical mechanics (thermodynamic limit), crossed products, semiclassical gravity.
Type III (III ₀ , III _λ , III ₁)	Purely infinite. All non-zero projections are infinite and equivalent.	Admits no trace. No density matrices exist.	Von Neumann entropy is strictly undefined / infinite.	Localized subregions in AQFT, causal diamonds, black hole event horizons.

Table 2: The Murray-von Neumann classification of operator algebras, highlighting the trace properties and entropy viability corresponding to distinct physical regimes.

In standard finite quantum mechanics, the system operates within a Type I algebra. Observables possess discrete eigenvectors (pure states), and the system admits a normalized trace, allowing the state to be described by a density matrix ρ . The von Neumann entropy is easily calculated, quantifying the entanglement or thermodynamic disorder of the system.

The Type III₁ Pathology of Localized Spacetime

However, a fundamental and inescapable result of AQFT proves that the local algebra of observables associated with any bounded spacetime subregion in a relativistic quantum field theory is a von Neumann factor of Type III₁.

This pathology is not a mathematical artifact; it is a direct physical consequence of the Operator Grammar's Boundary Law (C₅) and Reachable History Law (C₂). When an observer defines a local region of spacetime, they draw a boundary (a causal horizon). Because the quantum vacuum is infinitely entangled across arbitrarily short distances spanning this boundary (the Reeh-Schlieder theorem and UV divergences), the resulting local algebra is inundated with infinite degrees of freedom.

Because Type III₁ algebras possess no minimal projections and admit no semi-finite trace, density matrices mathematically cannot exist within the local algebra. Consequently, the standard expression for von Neumann entanglement entropy is strictly undefined. If the fundamental structure of localized reality is a Type III₁ algebra, then localized thermodynamics, finite measurement, and macroscopic entropy technically do not exist. The information is conserved in the global state, but the local observer is blinded by infinite boundary entanglement.

In the language of the Operator Grammar, this is the crisis of observing only the active line while ignoring the "bench" (the detached continuation state created by a lift operation). The infinite entanglement divergence is the penalty for attempting to calculate the state of a bounded region without accounting for the relational ties that cross the boundary into the complementary space.

The Crossed Product: Restoring the Observer

The resolution to the infinite entanglement crisis of Type III₁ algebras requires integrating the observer directly into the algebra. In the Operator Grammar, this corresponds to formally defining the Geometry / Read Head (C8) and recognizing that the universe must be actively rendered (C7). In the mathematics of von Neumann algebras, this is achieved through Tomita-Takesaki modular theory and the crossed product construction.

Tomita-Takesaki Modular Theory

For any von Neumann algebra $\mathcal{M}$ (even Type III) equipped with a cyclic and separating state $|\Omega\rangle$ (such as the quantum vacuum), Tomita-Takesaki theory defines a modular operator Δ and an anti-unitary modular conjugation operator J . These operators generate a one-parameter group of modular automorphisms, denoted $\sigma_t(x) = \Delta^{it} x \Delta^{-it}$, which maps the algebra onto itself.

In a Type III algebra, this modular automorphism group is strictly *outer*, meaning the unitary operator generating the flow does not reside within the local algebra $\mathcal{M}$ itself. Physically, this implies that the local dynamics of the spacetime region are being driven by an external parameter—an external clock or observer that the local mathematics cannot account for.

The Crossed Product Transition (Type III $\rightarrow$ Type II_∞)

To recover a well-defined trace and render finite physical measurements possible, the algebra must be expanded to include the observer. This is accomplished via the crossed product construction, denoted $\mathcal{M} \rtimes_{\sigma} \mathbb{R}$.

By taking the crossed product of the Type III₁ algebra with the locally compact group representing its own modular automorphism flow ($G = \mathbb{R}$), the external automorphisms are rendered *inner* relative to the newly enlarged algebra. Takesaki's duality theorem proves that this specific crossed product always yields a von Neumann factor of Type II_∞.

The transition to a Type II_∞ *factor is physically transformative*. Type II_∞ algebras admit a faithful, normal, semi-finite trace. Because a trace is restored, one can define generalized density operators ρ that are affiliated with the algebra. This allows for a rigorous, finite calculation of von Neumann entropy: $S(\rho) = -\text{Tr}(\rho \log \rho)$.

Physical Interpretation: The Read Head as Semiclassical Gravity

The crossed product is not merely a mathematical trick; it has profound physical significance. Taking the crossed product is mathematically isomorphic to gravitationally "dressing" the local quantum field observables to the energy of an external observer, reference frame, or clock. The auxiliary $\mathbb{R}$ space appended to the algebra represents the state space of the macroscopic observer's wavefunction.

In the terminology of the unified ontology, the original Type III₁ algebra is the raw, bounded potential of the substrate. The modular automorphism group is the Read Head (the observer's trajectory through the state space). The crossed product is the mathematical equivalent of the Render constraint (C7). The universe does not possess finite, localized thermodynamic properties until the observer's frame of reference is algebraically bound to the substrate. Rendering alters the algebraic type of the reality being observed, collapsing the infinite UV divergences of pure potential into the finite, measurable interactions of semiclassical reality.

Holography and the Algebraic Ryu-Takayanagi Formula

This algebraic transition successfully bridges quantum mechanics and gravity, particularly in the context of holographic duality (AdS/CFT) and black hole thermodynamics. The Bekenstein-Hawking entropy of a black hole ($S = \frac{\text{Area}}{4G_N}$) and the Ryu-Takayanagi formula for entanglement entropy are famously divergent in the strict semiclassical limit ($G_N \rightarrow 0$), precisely because the bulk fields constitute a Type III₁ algebra.

However, by transitioning to the holographic crossed product algebra, the divergences cancel. The relative entropy evaluated in the crossed product algebra factorizes exactly into a sum of two finite terms: the relative entropy of the original local algebra, and the relative entropy between the macroscopic observers' wavefunctions.

$$S_{\text{rel}}(\Psi_f|\Phi_g; \hat{\mathcal{A}}_{\text{crossed}}) \approx S_{\text{rel}}(\Psi|\Phi; \mathcal{A}) + S_{\text{rel}}(g|f)$$

This provides a completely finite, algebraically rigorous formulation of holographic entropy. It proves that the macroscopic geometry of spacetime, gravitational attraction, and thermodynamic entropy are not fundamental ingredients of the universe; they are the emergent, finite manifestations of Type II_∞ crossed-product constraints operating on a substrate of pure relational information.

Conclusion

The convergence of Constructor Theory, Euclidean Distance Geometry, Algebraic Quantum Field Theory, and discrete Operator Logic establishes a radical, unified ontology of the universe. Reality is not a pre-existing geometric void populated by material particles. It is a substrate-independent algebra of pure distinctions, evolving through the stringent constraints of a generative Operator Grammar.

The foundational laws (Co-C13) guarantee that transformations cannot annihilate information; they can only bound it and relegate it to detached relational states. As the system addresses the unresolved remainders of its own transformation space, dimensional complexity is mathematically forced into existence. The physical realization of this complexity as continuous space and volume is merely a constraint-satisfaction rendering, governed at all scales by the Cayley-Menger determinant.

At the boundaries of these spatial renders—at causal horizons and localized regions—the severing of relational connections produces the infinite UV entanglement characteristics of Type III₁ von Neumann algebras. This renders local domains mathematically pathological, blinding the observer with infinite potential and rendering classical entropy undefined.

The definitive resolution to this physical pathology is the Crossed Product. By algebraically entangling the local Type III₁ quantum field with its own modular automorphism group, the system integrates the "Read Head"—the observer and their reference frame—directly into the mathematical structure. This transition to a Type II_∞ factor restores a semifinite trace, allows for the definition of density matrices, and yields the finite, calculable entropies required by thermodynamics and semiclassical gravity.

In totality, the universe is the continuous process of evaluating the remainders of its own transformation space, rendering continuous geometries from discrete distances, and regularizing its infinite quantum entanglements through the observational crossed product. The mass, geometry, and entropy we measure are not the fundamental substrate of the universe; they are the thermodynamic cost of its algebraic rendering.

Works cited

1. [2505.08692] Constructor theory of time - arXiv, <https://arxiv.org/abs/2505.08692>
2. Constructor theory of information - PMC - NIH, <https://pmc.ncbi.nlm.nih.gov/articles/PMC4309123/>
3. Existence of an information unit as a postulate of quantum theory, <https://arxiv.org/html/1208.0493v2>
4. Unplugging a Seemingly Sentient Machine Is the Rational Choice, <https://arxiv.org/html/2601.21016>
5. How Quantum Mechanics Derives from a Revolutionary New Theory, <https://medium.com/the-physics-arxiv-blog/how-quantum-mechanics-derives-from-a-revolutionary-new-theory-of-information-4487489dbb34>
6. Yes.docx
7. Inverse Evolution and Dimensional Collapse: Operator-Theoretic, <https://www.mdpi.com/2073-8994/18/7/1230>
8. Cayley–Menger determinant - Wikipedia, https://en.wikipedia.org/wiki/Cayley%E2%80%93Menger_determinant
9. Using Cayley-Menger Determinants for geometric constraint solving, https://www.researchgate.net/publication/210232160_Using_Cayley-Menger_Determinants_for_geometric_constraint_solving
10. arXiv:2306.12030v1 [math.CO] 21 Jun 2023, <https://arxiv.org/pdf/2306.12030>
11. Crossed product algebras and generalized entropy for subregions, <https://arxiv.org/html/2306.07323v2>
12. Finite Entropy Implies Finite Dimension in Quantum Gravity - arXiv, <https://arxiv.org/pdf/2509.17856>
13. 1 Introduction - arXiv, <https://arxiv.org/html/2507.01419v1>
14. Semiclassical algebraic reconstruction for type III algebras - arXiv, <https://arxiv.org/pdf/2605.13576>
15. Euclidean distance geometry and applications - arXiv, <https://arxiv.org/pdf/1205.0349>
16. Six mathematical gems from the history of Distance Geometry - arXiv, <https://arxiv.org/pdf/1502.02816>
17. The strength of a geometric simplex - arXiv, <https://arxiv.org/pdf/2602.17630>
18. Realizability of the Lorentzian (n,1)-Simplex - arXiv, <https://arxiv.org/pdf/1110.5694>
19. arXiv:1810.05282v1 [gr-qc] 11 Oct 2018, <https://arxiv.org/pdf/1810.05282>
20. On the existence of two affine-equivalent frameworks with ... - arXiv, <https://arxiv.org/html/2306.13859v1>
21. Causal association of electromagnetic signals using the Cayley, <https://arxiv.org/html/0908.3143v1>
22. arXiv:0908.3143v1 [gr-qc] 21 Aug 2009, <https://arxiv.org/pdf/0908.3143>

23. arXiv:2407.13601v2 [gr-qc] 23 Oct 2024, <https://arxiv.org/pdf/2407.13601>
24. arXiv:1506.06839v4 [gr-qc] 31 Jul 2016, <https://arxiv.org/pdf/1506.06839>
25. Chapter 1 A universe of processes and some of its guises - arXiv, <https://arxiv.org/pdf/1009.3786>
26. arXiv:2407.01695v2 [hep-th] 6 Feb 2025, <https://arxiv.org/pdf/2407.01695>
27. Crossed product algebras and generalized entropy for subregions, <https://arxiv.org/pdf/2306.07323>
28. Crossed Products, Extended Phase Spaces and the Resolution of,
<https://arxiv.org/html/2306.09314v4>
29. arXiv:2112.11614v5 [hep-th] 22 Jul 2022, <https://arxiv.org/pdf/2112.11614>
30. Notes on the type classification of von Neumann algebras - arXiv,
<https://arxiv.org/html/2302.01958v4>
31. The Role of Infinite Tensor Products and Nested Wigner's Friends,
<https://arxiv.org/html/2409.06470v1>
32. Semiclassical algebraic reconstruction for type III algebras - arXiv,
<https://arxiv.org/html/2605.13576v2>
33. Generalized Black Hole Entropy is von Neumann Entropy - arXiv, <https://arxiv.org/pdf/2309.15897>
34. The Crossed Product, Modular (Tomita) Dynamics and its Role in, <https://arxiv.org/pdf/2507.01419>
35. a view from von Neumann Algebras - arXiv, <https://arxiv.org/pdf/2302.14747>
36. arXiv:1804.05579v3 [math-ph] 1 May 2025, <https://arxiv.org/pdf/1804.05579>
37. Inflationary Cosmology as flow of integrable weights - arXiv, <https://arxiv.org/pdf/2407.20671>
38. Perturbative semiclassical entropy of dynamical black holes - arXiv, <https://arxiv.org/pdf/2603.02320>