

On the sum-of-squares degree of symmetric quadratic functions

Troy Lee* Anupam Prakash* Ronald de Wolf† Henry Yuen‡

March 9, 2016

arXiv:1601.02311v2 [cs.CC] 8 Mar 2016

Abstract

We study how well functions over the boolean hypercube of the form $f_k(x) = (|x| - k)(|x| - k - 1)$ can be approximated by sums of squares of low-degree polynomials, obtaining good bounds for the case of approximation in ℓ_∞-norm as well as in ℓ_1-norm. We describe three complexity-theoretic applications: (1) a proof that the recent breakthrough lower bound of Lee, Raghavendra, and Steurer [LRS15] on the positive semidefinite extension complexity of the correlation and TSP polytopes cannot be improved further by showing better sum-of-squares degree lower bounds on ℓ_1-approximation of f_k; (2) a proof that Grigoriev's lower bound on the degree of Positivstellensatz refutations for the knapsack problem is optimal, answering an open question from [Gri01]; (3) bounds on the query complexity of quantum algorithms whose expected output approximates such functions.

*School of Physical and Mathematical Sciences, Nanyang Technological University and Centre for Quantum Technologies, Singapore. This material is based on research supported by the Singapore National Research Foundation under NRF RF Award No. NRF-NRFF2013-13. Email: troyjlee@gmail.com, aprakash@ntu.edu.sg

†QuSoft, CWI and University of Amsterdam, the Netherlands. Partially supported by ERC Consolidator Grant QPROGRESS and by the European Commission FET-Proactive project Quantum Algorithms (QALGO) 600700. rdewolf@cwi.nl

‡MIT, Cambridge, MA. Supported by NSF grants 1218547 and 1452302 and a Simons Graduate Award in Theoretical Computer Science. hyuen@mit.edu