

Multipartite Entanglement in Stabilizer Tensor Networks

Sepehr Nezami[1,2] and Michael Walter[1,3]

[1]*Stanford Institute for Theoretical Physics, Stanford University, Stanford, USA*
[2]*Institute for Quantum Information and Matter and Walter Burke Institute for Theoretical Physics, California Institute of Technology, Pasadena CA 91125, USA*
[3]*Korteweg-de Vries Institute for Mathematics, Institute for Theoretical Physics, Institute for Logic, Language, and Computation, and QuSoft, University of Amsterdam, The Netherlands*

Despite the fundamental importance of quantum entanglement in many-body systems, our understanding is mostly limited to bipartite situations. Indeed, even defining appropriate notions of multipartite entanglement is a significant challenge for general quantum systems. In this work, we initiate the study of multipartite entanglement in a rich, yet tractable class of quantum states called *stabilizer tensor networks*. We demonstrate that, for generic stabilizer tensor networks, the *geometry* of the tensor network informs the multipartite entanglement structure of the state. In particular, we show that the average number of Greenberger-Horne-Zeilinger (GHZ) triples that can be extracted from a stabilizer tensor network is small, implying that tripartite entanglement is scarce. This, in turn, restricts the higher-partite entanglement structure of the states. Recent research in quantum gravity found that stabilizer tensor networks reproduce important structural features of the AdS/CFT correspondence, including the Ryu-Takayanagi formula for the entanglement entropy and certain quantum error correction properties. Our results imply a new operational interpretation of the monogamy of the Ryu-Takayanagi mutual information and an entropic diagnostic for higher-partite entanglement. Our technical contributions include a spin model for evaluating the average GHZ content of stabilizer tensor networks, as well as a novel formula for the third moment of random stabilizer states, which we expect to find further applications in quantum information.

arXiv:1608.02595v4 [quant-ph] 13 Jan 2021

Quantum entanglement is of fundamental relevance for the behavior of quantum mechanical systems in condensed matter and high energy physics. From the perspective of quantum information processing, it is the resource that provides speedups in quantum computing, security in quantum cryptography, and improved performance in quantum sensing. However, the structure of many-body or multipartite entanglement is only poorly understood [1]. In this work, we focus on analyzing multipartite entanglement in an important but tractable class of quantum states known as *stabilizer tensor networks*, i.e., tensor networks that are obtained by contracting stabilizer states. Stabilizer states are an important family of quantum states that can be highly entangled (even maximally so) but still have sufficient algebraic structure to admit an efficient classical description. This makes them a versatile tool in quantum information theory, particularly in the theory of quantum error correction [2]. Of particular import in the present context is that the tripartite entanglement structure of stabilizer states can be precisely quantified – any tripartite stabilizer state is locally equivalent to a collection of bipartite Bell pairs and tripartite GHZ states [3, 4] (cf. [5–11]).

An important additional motivation to study stabilizer tensor networks comes from current research in quantum gravity. In recent years, research in quantum gravity and quantum information theory has been inspired by a fruitful mutual exchange of ideas. Tensor networks in particular provide a common framework, rooted in the similarity between the structure of the tensor network and the bulk geometry in holographic duality [12–14]. A paradigmatic example is the Ryu-Takayanagui formula, $S(A) \simeq |\gamma_A|/4G_N$, which asserts that the entanglement entropy of a boundary region A in a holographic state is in leading order proportional to the area of a corresponding minimal surface γ_A in the bulk geometry [15, 16]. Likewise, in any tensor network, the entanglement entropy of a boundary subsystem can be upper-bounded in terms of the size of a minimal cut through the network [17] (Fig. 1). This bound can be saturated not only through the choice of suitable tensors [18, 19] but is in fact a generic phenomenon in *random tensor networks* with large bond dimension [20, 21], the mechanism of which can be understood in terms of multipartite entanglement distillation. These tensor network models not only reproduce the Ryu-Takayanagi formula for the entanglement entropy, but they also implement several other significant features of holographic duality [18–20]. In many ways, these properties follow from the bipartite entanglement structure and can be therefore reduced to entropic considerations.




Figure 1. **Stabilizer tensor networks.** A tensor network state is obtained by placing random stabilizer states at the bulk vertices (blue) and contracting according to the edges of the graph. In the limit of large bond dimensions, the average entanglement entropy of a boundary region A is proportional to the length of a minimal cut γ_A through the network (dashed line) [20], $S(A) \simeq S_{RT}(A)$, reproducing the Ryu-Takayanagi formula in holography.

In this paper, we initiate a study of *multipartite entanglement* in random tensor network models. Our motivation is twofold: First, recent research in quantum gravity has raised profound questions regarding the multipartite entanglement in holographic states [22–24], in particular with regards to tripartite entanglement of GHZ type [25, 26]. Answers to these questions in the context of tensor network models will likely