arXiv:1611.08571v1 [quant-ph] 25 Nov 2016

On preparing ground states of gapped Hamiltonians: An efficient Quantum Lovász Local Lemma

András Gilyén* Or Sattath†

Abstract

A frustration-free local Hamiltonian has the property that its ground state minimises the energy of all local terms simultaneously. In general, even deciding whether a Hamiltonian is frustration-free is a hard task, as it is closely related to the QMA_1-complete quantum satisfiability problem (QSAT) – the quantum analogue of SAT, which is the archetypal NP-complete problem in classical computer science. This connection shows that the frustration-free property is not only relevant to physics but also to computer science.

The Quantum Lovász Local Lemma (QLLL) provides a sufficient condition for frustration-freeness. A natural question is whether there is an efficient way to prepare a frustration-free state under the conditions of the QLLL. Previous results showed that the answer is positive if all local terms commute.

In this work we improve on the previous constructive results by designing an algorithm that works efficiently for non-commuting terms as well, assuming that the system is "uniformly" gapped, by which we mean that the system *and all its subsystems* have an inverse polynomial energy gap. Also, our analysis works under the most general condition for the QLLL, known as Shearer's bound. Similarly to the previous results, our algorithm has the charming feature that it uses only local measurement operations corresponding to the local Hamiltonian terms.

1 Introduction

Frustration-free Hamiltonians and quantum satisfiability. Most physical systems and models are described by a local Hamiltonian $H = \sum_i H_i$ where each k-local term H_i acts non-trivially only on at most k of its subsystems. Such a Hamiltonian is called *frustration-free* if its ground state is also the ground state of each of the local terms H_i. Frustration-free Hamiltonians appear in various areas, for example: quantum error correcting codes [18], parent Hamitlonians for PEPS (a 2-D generalisation of matrix-product-states) [33], and various models in many-body quantum physics.

An equivalent way to ask whether a Hamiltonian H is frustration-free is whether $H' = \sum_i \Pi_i$ is frustration-free, where Π_i is the projector on the excited states of H_i. The quantum satisfiability problem (QSAT) [1] is to determine whether H' in the above form is frustration-free. QSAT is QMA_1-Complete [8], and therefore intractable in general even for quantum computers (unless $\mathsf{BQP} = \mathsf{QMA}_1$). Finding the ground state of frustration-free Hamiltonians – the challenge we tackle in this work – is, in general, an even harder task.

*QuSoft, CWI and University of Amsterdam, the Netherlands. gilyen@cwi.nl

†The Hebrew University and MIT. sattath@cs.huji.ac.il

[1] For technical reasons, that would not be relevant for this work, there is a promise that if H' is not frustration-free, the minimal energy of H' is at least inverse polynomial in the number of qubits.