

2017–
2021

arXiv:1708.00360v4 [quant-ph] 12 Nov 2018

Disentanglement Cost of Quantum States

Mario Berta[1] and Christian Majenz[2,3]

[1]*Department of Computing, Imperial College London, London*
[2]*Institute for Language, Logic and Computation, University of Amsterdam, Amsterdam*
[3]*QuSoft, Amsterdam*

We show that the minimal rate of noise needed to catalytically erase the entanglement in a bipartite quantum state is given by the regularized relative entropy of entanglement. This offers a solution to the central open question raised in [Groisman *et al.*, PRA 72, 032317 (2005)] and complements their main result that the minimal rate of noise needed to erase all correlations is given by the quantum mutual information. We extend our discussion to the tripartite setting where we show that an asymptotic rate of noise given by the regularized relative entropy of recovery is sufficient to catalytically transform the state to a locally recoverable version of the state.

Introduction Quantifying and classifying quantum correlations is of fundamental importance in quantum information theory [22]. Motivated by Landauer's erasure principle [24] one way to quantify the correlations present in a bipartite quantum state ρ_{AB} is to measure the amount of noise that is required to erase them. In that respect, Groisman *et al.* [19] showed that the optimal asymptotic rate of local noise to bring ρ_{AB} close to a product $\sigma_A \otimes \sigma_B$ is given by the *quantum mutual information*

$$I(A:B)_\rho := D(\rho_{AB}\|\rho_A\otimes\rho_B) = \inf_{\sigma\in\mathrm{PR}} D(\rho_{AB}\|\sigma_A\otimes\sigma_B) \tag{1}$$

with $\mathrm{PR}(A:B)$ the set of product states in $A:B$ and $D(\rho\|\sigma) := \mathrm{Tr}\left[\rho(\log\rho - \log\sigma)\right]$ is the quantum relative entropy. Hence, the quantum mutual information quantifies the total amount of correlations in bipartite states – including both the quantum and classical ones. Alternatively, we can write

$$I(A:B)_\rho = \inf_{\sigma\in\mathrm{PR}} H(AB)_{\sigma\otimes\sigma} - H(AB)_\rho \tag{2}$$

with $H(A)_\rho := -\mathrm{tr}\left[\rho_A \log \rho_A\right]$ the von Neumann entropy. Thus, the cost function $I(A:B)_\rho$ can conveniently be understood as either the quantum relative entropy distance to the next product state as in Eq. (1) or as the amount of entropy injected into the system to reach the next product state as in Eq. (2). This finding was generalized in various directions, including a catalytic analysis of the one-shot case [26], the study of tripartite correlations [4, 37, 38], as well as the study of coherence [31] and more general symmetries [36]. However, it remained open how to quantify the optimal asymptotic rate of local noise to bring ρ_{AB} close to a separable state $\sum_j p_j \sigma_A^j \otimes \sigma_B^j$. In particular, it was unclear if a quantity defined in such a way can be the basis of a proper entanglement measure.

In this note, we solve the problem and give a precise mathematical model for erasing entanglement in bipartite states where the optimal asymptotic rate of local noise needed to get close to a separable state is given by the *regularized relative entropy of entanglement*. In particular, this also gives a new operational interpretation to the distance measure quantum relative entropy.

Entanglement Measures. As discussed the quantum mutual information is a measure for the total amount of correlations and in the following we introduce more refined measures only capturing the quantum correlations. The *relative entropy of entanglement* is given by [33]

$$E(A:B)_\rho := \inf_{\sigma\in\mathrm{SEP}} D(\rho\|\sigma)\,, \tag{3}$$

where $\mathrm{SEP}(A:B)$ denotes the set of separable states in $A:B$. Since the relative entropy of entanglement is in general not additive on tensor product states, it has to be regularized [3, 35]. The regularized relative entropy of entanglement is defined as

$$E^\infty(A:B)_\rho := \lim_{n\to\infty} \frac{1}{n} E(A:B)_{\rho^{\otimes n}} \tag{4}$$

This quantity has an operational interpretation in composite asymmetric quantum hypothesis testing as the asymptotic exponential rate of mistakenly identifying ρ_{AB} instead of a state separable in $A:B$ [8]. As a corresponding one-shot analogue based on the smooth max-relative entropy [16]

$$D^\varepsilon_{\max}(\rho\|\sigma) := \inf_{\bar\rho\approx_\varepsilon\rho} \inf\left\{\lambda:\ 2^\lambda\cdot\sigma - \bar\rho \geq 0\right\} \tag{5}$$

with $\bar\rho \approx_\varepsilon \rho$ in purified distance [32],

we have the *smooth max-relative entropy of entanglement* [15]

$$E^\varepsilon_{\max}(A:B)_\rho := \inf_{\sigma\in\mathrm{SEP}} D^\varepsilon_{\max}(\bar\rho_{AB}\|\sigma_{AB})\,. \tag{6}$$

This is a smoothed version of the logarithm global robustness of entanglement [34]. All the quantities

$$E(A:B)_\rho,\ E^\infty(A:B)_\rho,\ E^\varepsilon_{\max}(A:B)_\rho$$

define proper entanglement measures with mathematical properties as requested by axiomatic entanglement theory (see, e.g., [5, 13] for an overview). We emphasize that these type of information-theoretic entanglement measures have been vastly useful for understanding the entanglement structure of multipartite quantum states in many body physics. In particular, this lead to strong insights about entropic area laws [9, 10, 20, 39] and detecting topological order [23, 25] in condensed matter physics as well as to basics findings in quantum thermodynamics [6, 12, 17, 19].