

arXiv:1711.08039v1 [cs.CC] 21 Nov 2017

Alternating minimization, scaling algorithms, and the null-cone problem from invariant theory

Peter Bürgisser[1], Ankit Garg[2], Rafael Oliveira[3,4], Michael Walter[5,6], and Avi Wigderson[7]

[1]*Institut für Mathematik, Technische Universität Berlin*
[2]*Microsoft Research New England*
[3]*Department of Computer Science, Princeton University*
[4]*Department of Computer Science, University of Toronto*
[5]*QuSoft, Korteweg-de Vries Institute for Mathematics, Institute of Physics, and Institute for Logic, Language and Computation, University of Amsterdam*
[6]*Stanford Institute for Theoretical Physics, Stanford University*
[7]*Institute for Advanced Study, Princeton*

Abstract

Alternating minimization heuristics seek to solve a (difficult) global optimization task through iteratively solving a sequence of (much easier) local optimization tasks on different parts (or blocks) of the input parameters. While popular and widely applicable, very few examples of this heuristic are rigorously shown to converge to optimality, and even fewer to do so efficiently.

In this paper we present a general framework which is amenable to rigorous analysis, and expose its applicability. Its main feature is that the local optimization domains are each a group of invertible matrices, together naturally acting on tensors, and the optimization problem is minimizing the norm of an input tensor under this joint action. The solution of this optimization problem captures a basic problem in Invariant Theory, called the *null-cone problem*.

This algebraic framework turns out to encompass natural computational problems in combinatorial optimization, algebra, analysis, quantum information theory, and geometric complexity theory. It includes and extends to high dimensions the recent advances on (2-dimensional) *operator scaling* [Gur04, GGOW16, IQS17].

Our main result is a fully polynomial time approximation scheme for this general problem, which may be viewed as a multi-dimensional scaling algorithm. This directly leads to progress on some of the problems in the areas above, and a unified view of others. We explain how faster convergence of an algorithm for the same problem will allow resolving central open problems.

Our main techniques come from Invariant Theory, and include its rich *non-commutative duality theory*, and new bounds on the bitsizes of coefficients of *invariant polynomials*. They enrich the algorithmic toolbox of this very computational field of mathematics, and are directly related to some challenges in geometric complexity theory (GCT).