

Matrix product states and the quantum max-flow/min-cut conjectures

Fulvio Gesmundo,[1] J. M. Landsberg,[2] and Michael Walter[3,4]

[1)] *QMATH, University of Copenhagen, Universitetsparken 5, 2100 København Ø, Denmark*[a)]

[2)] *Department of Mathematics, Texas A&M University, College Station, TX 77843-3368, USA*[b)]

[3)] *QuSoft, Korteweg-de Vries Institute for Mathematics, Institute for Theoretical Physics, Institute for Logic, Language and Computation, University of Amsterdam, 1098 XG Amsterdam, Netherlands*

[4)] *Stanford Institute for Theoretical Physics, Stanford University, Stanford, CA 94305, USA*[c)]

In this note we discuss the geometry of matrix product states with periodic boundary conditions and provide three infinite sequences of examples where the quantum max-flow is strictly less than the quantum min-cut. In the first we fix the underlying graph to be a 4-cycle and verify a prediction of Hastings that inequality occurs for infinitely many bond dimensions. In the second we generalize this result to a $2d$-cycle. In the third we show that the $2d$-cycle with periodic boundary conditions gives inequality for all d when all bond dimensions equal two, namely a gap of at least 2^{d-2} between the quantum max-flow and the quantum min-cut.

arXiv:1801.09106v2 [quant-ph] 18 Oct 2018

I. INTRODUCTION

A *tensor network* associated to a graph is a way of constructing tensors in large spaces from smaller building-block tensors. From the perspective of algebraic geometry, tensor networks provide a natural way of constructing varieties of tensors. The use of graphs to study tensors dates back at least to Clifford in 1881 (see Ref. 1, Fig. 2.11.1). In applied mathematics and physics, tensor networks are versatile tools for efficiently approximating high-dimensional data, such as ground states of many-body quantum systems in condensed matter physics (see, e.g., Refs. 2–4).

In this paper we focus exclusively on *translation-invariant matrix product states with periodic boundary conditions* as defined in §I A. We use them to obtain three sequences of examples of tensor networks where the quantum max-flow (§I B) is strictly less than the quantum min-cut (§I C) for certain partitions. Our results disprove a natural conjecture and verify a numerical prediction from Ref. 5 (§I D).

A. Matrix product states with periodic boundary conditions

In this paper, we work exclusively with tensor networks associated with the oriented cyclic graphs C_m, as in Figure 1 (a), and one building-block tensor T taken from $\mathbb{C}^N \otimes \mathbb{C}^n \otimes (\mathbb{C}^n)^*$, which we write as $A \otimes B \otimes B^*$.

One constructs tensors in $A^{\otimes m} = (\mathbb{C}^N)^{\otimes m}$ from this set-up as follows: For each vertex, we associate a copy of A to the external (physical) edge, a copy of B to the outgoing edge from the circle, and a copy of B^* to

[a)] Electronic mail: fulges@math.ku.dk
[b)] Electronic mail: jml@math.tamu.edu
[c)] Electronic mail: m.walter@uva.nl