

2017–
2021

Conditional Decoupling of Quantum Information

Mario Berta,[1] Fernando G. S. L. Brandão,[2] Christian Majenz,[3,4] and Mark M. Wilde[5]

[1]*Department of Computing, Imperial College London, London SW7 2AZ, UK*
[2]*Institute for Quantum Information and Matter, California Institute of Technology, Pasadena, California 91125, USA*
[3]*Institute for Language, Logic and Computation, University of Amsterdam, and QuSoft, 1098XG Amsterdam, Netherlands*
[4]*Department of Mathematical Sciences, University of Copenhagen, Universitetsparken 5, DK-2100 Copenhagen*
[5]*Hearne Institute for Theoretical Physics, Department of Physics and Astronomy, Center for Computation and Technology, Louisiana State University, Baton Rouge, Louisiana 70803, USA*

(Dated: August 2, 2018)

Insights from quantum information theory show that correlation measures based on quantum entropy are fundamental tools that reveal the entanglement structure of multipartite states. In that spirit, Groisman, Popescu, and Winter [Physical Review A **72**, 032317 (2005)] showed that the quantum mutual information $I(A;B)$ quantifies the minimal rate of noise needed to erase the correlations in a bipartite state of quantum systems AB. Here, we investigate correlations in tripartite systems ABE. In particular, we are interested in the minimal rate of noise needed to apply to the systems AE in order to erase the correlations between A and B given the information in system E, in such a way that there is only negligible disturbance on the marginal BE. We present two such models of conditional decoupling, called deconstruction and conditional erasure cost of tripartite states ABE. Our main result is that both are equal to the conditional quantum mutual information $I(A;B|E)$ – establishing it as an operational measure for tripartite quantum correlations.

arXiv:1808.00135v1 [quant-ph] 1 Aug 2018

Introduction. Landauer's principle states that the amount of work needed for erasing a memory is proportional to the amount of information stored in the memory [1]. Motivated by this principle, the correlations of a bipartite quantum state ρ_{AB} shared between two parties Alice and Bob can be quantified by the amount of noise that is required to erase the correlations in ρ_{AB}. This erasure cost is closely connected to the thermodynamical cost of erasing the correlations [2], which in turn is part of the larger context of the physics of erasure (see, e.g., Refs. [3–6]). In a model of Groisman, Popescu, and Winter [2] [7], Alice is allowed to pick a free ancilla, in the form of an already decoupled state $\theta_{A'}$, and then applies a *unitary randomizing channel*

$$\Lambda_{AA'}(\cdot) := \frac{1}{M}\sum_{i=1}^{M} U^i_{AA'}(\,\cdot\,)\left(U^i_{AA'}\right)^\dagger , \tag{1}$$

where the noise injected into the system comes from averaging over the unitaries. The goal is for the resulting state to become close to a product state (or, in other words, *decoupled*)

$$F\left(\Lambda_{AA'}(\rho_{AB}\otimes\theta_{A'}), \pi_{A'A}\otimes\rho_B\right) \geq 1-\varepsilon , \tag{2}$$

where $\pi_{AA'}$ is a maximally mixed state on a subspace of AA'. Here, the action of the channel $\Lambda_{AA'}$ on systems $AA'B$ is understood as $\Lambda_{AA'}\otimes\mathcal{I}_B$, where $\mathcal{I}_B$ denotes the identity channel, and the fidelity between states ξ and χ is given by $F(\xi,\chi) := [\mathrm{Tr}(\sqrt{\sqrt{\chi}\xi\sqrt{\chi}})]^2$. We note that the use of the ancilla is catalytic in the sense that the system A' has to stay decoupled from B (at least approximately), but potentially makes the erasure process more efficient [8]. The main result of Groisman, Popescu, and Winter [2, Thm. 1] is that the minimal rate of unitaries needed in the limit of many copies $\rho_{AB}^{\otimes n}$ and vanishing error $\varepsilon \to 0$ is given by the *quantum mutual information (QMI)*

$$\frac{1}{n}\log M \to I(A;B)_\rho := H(A)_\rho + H(B)_\rho - H(AB)_\rho ,$$

with the quantum entropy of a state η_X on system X given by $H(X)_\eta := -\mathrm{Tr}\left[\eta_X \log \eta_X\right]$. Thus, we can conclude that the QMI is equal to the amount of noise needed for *correlation destruction* between systems A and B. This result gives information-theoretic justification for the diverse use of the QMI as a correlation measure in quantum physics. For instance, it is a stepping stone in a quantitative understanding of *decoupling*, a central concept both in quantum information theory and in physics in general, with implications ranging from the black-hole information paradox [9–11] to area laws in quantum many-body systems [12].

Conditional measures of correlations. Here, we aim to quantify the correlations in a tripartite quantum state ρ_{ABE}. A measure that is (informally) understood as quantifying the correlations between A and B from the perspective of system E is the *conditional quantum mutual information (CQMI)*

$$I(A;B|E)_\rho := I(AE;B)_\rho - I(E;B)_\rho . \tag{3}$$

The CQMI is always non-negative $I(A;B|E)_\rho \geq 0$, an entropy inequality known as strong sub-additivity [13]. The mentioned informal interpretation of the CQMI can