

Bit Threads and Holographic Monogamy

Shawn X. Cui,[1*] Patrick Hayden,[1] Temple He,[2†] Matthew Headrick,[3,4] Bogdan Stoica,[3,5] and Michael Walter[6]

[1] *Stanford Institute for Theoretical Physics, Stanford University, Stanford, CA 94305, USA*

[2] *Center for the Fundamental Laws of Nature, Harvard University, Cambridge, MA 02138, USA*

[3] *Martin A. Fisher School of Physics, Brandeis University, Waltham, MA 02453, USA*

[4] *Center for Theoretical Physics, Massachusetts Institute of Technology, Cambridge, MA 02139, USA*

[5] *Department of Physics, Brown University, Providence, RI 02912, USA*

[6] *Korteweg-de Vries Institute for Mathematics, Institute of Physics, Institute for Logic, Language & Computation, and QuSoft, University of Amsterdam, 1098 XG Amsterdam, The Netherlands*

xingshan@vt.edu, phayden@stanford.edu, tmhe@ucdavis.edu, headrick@brandeis.edu, bstoica@brandeis.edu, m.walter@uva.nl

arXiv:1808.05234v2 [hep-th] 28 Jun 2019

Abstract

Bit threads provide an alternative description of holographic entanglement, replacing the Ryu-Takayanagi minimal surface with bulk curves connecting pairs of boundary points. We use bit threads to prove the monogamy of mutual information (MMI) property of holographic entanglement entropies. This is accomplished using the concept of a so-called multicommodity flow, adapted from the network setting, and tools from the theory of convex optimization. Based on the bit thread picture, we conjecture a general ansatz for a holographic state, involving only bipartite and perfect-tensor type entanglement, for any decomposition of the boundary into four regions. We also give new proofs of analogous theorems on networks.

BRX-6330, Brown HET-1764, MIT-CTP/5036

*Present affiliation: Department of Mathematics, Virginia Tech, Blacksburg, VA 24061, USA

†Present affiliation: Center for Quantum Mathematics and Physics (QMAP), Department of Physics, University of California, Davis, CA 95616, USA