

Distributional property testing in a quantum world

András Gilyén* Tongyang Li†

February 5, 2019

arXiv:1902.00814v1 [quant-ph] 2 Feb 2019

Abstract

A fundamental problem in statistics and learning theory is to test properties of distributions. We show that quantum computers can solve such problems with significant speed-ups. In particular, we give fast quantum algorithms for testing closeness between unknown distributions, testing independence between two distributions, and estimating the Shannon / von Neumann entropy of distributions. The distributions can be either classical or quantum, however our quantum algorithms require coherent quantum access to a process preparing the samples. Our results build on the recent technique of quantum singular value transformation, combined with more standard tricks such as divide-and-conquer. The presented approach is a natural fit for distributional property testing both in the classical and the quantum case, demonstrating the first speed-ups for testing properties of density operators that can be accessed coherently rather than only via sampling; for classical distributions our algorithms significantly improve the precision dependence of some earlier results.

1 Introduction

Distributional property testing is a fundamental problem in theoretical computer science (see, e.g. Goldreich (2017)). In such property testing questions the goal is to determine properties of probability distributions with the least number of independent samples. This has intimate connections and applications to statistics, learning theory, and algorithm design.

The merit of distributional property testing mainly comes from the fact that the testing of many properties admits *sublinear* algorithms. For instance, given the ability to take samples from a discrete distribution p on $[n] := \{1, \dots, n\}$, it requires $\Theta(n/\epsilon^2)$ samples to "learn" p, i.e., to construct a distribution q on $[n]$ such that $\|p - q\|_1 \leq \epsilon$ with success probability at least $2/3$ ($\|\cdot\|_1$ being ℓ^1-distance). However, testing whether $p = q$ or $\|p - q\|_1 > \epsilon$ requires only $\Theta(\max\{\frac{n^{2/3}}{\epsilon^{4/3}}, \frac{n^{1/2}}{\epsilon^2}\})$ samples from p and q (Chan et al. (2014)), which is sublinear in n and significantly smaller than the complexity of learning the entire distributions. See Section 1.4 for more examples and discussions.

In this paper, we study the impact of quantum computation on distributional property testing problems. We are motivated by the emerging topic of "quantum property testing" (see the survey of Montanaro and de Wolf (2016)) which focuses on investigating the quantum advantage in testing classical statistical properties. Quantum speed-ups have already been established for a

*QuSoft, CWI and University of Amsterdam, the Netherlands. Supported by ERC Consolidator Grant QPROGRESS and partially supported by QuantERA project QuantAlgo 680-91-034. gilyen@cwi.nl

†Department of Computer Science, Institute for Advanced Computer Studies, and Joint Center for Quantum Information and Computer Science, University of Maryland. Supported by IBM PhD Fellowship, QISE-NET Triplet Award (NSF DMR-1747426), and the U.S. Department of Energy, Office of Science, Office of Advanced Scientific Computing Research, Quantum Algorithms Teams program. tongyang@cs.umd.edu