

Towards a theory of non-commutative optimization: geodesic first and second order methods for moment maps and polytopes

Peter Bürgisser* Cole Franks† Ankit Garg‡ Rafael Oliveira§
Michael Walter¶ Avi Wigderson‖

arXiv:1910.12375v3 [math.OC] 26 Jul 2021

Abstract

This paper initiates a systematic development of a theory of *non-commutative optimization*. It aims to unify and generalize a growing body of work from the past few years which developed and analyzed algorithms for natural *geodesically convex* optimization problems on Riemannian manifolds that arise from the symmetries of non-commutative groups. These algorithms minimize the *moment map* (a non-commutative notion of the usual *gradient*) and test membership in *null cones* and *moment polytopes* (a vast class of polytopes, typically of exponential vertex and facet complexity, which arise from this a priori non-convex, non-linear setting). This setting captures a diverse set of problems in different areas of computer science, mathematics, and physics. Several of them were solved efficiently for the first time using non-commutative methods; the corresponding algorithms also lead to solutions of purely structural problems and to many new connections between disparate fields.

In the spirit of standard convex optimization, we develop two general methods in the geodesic setting, a first order and a second order method, which respectively receive first and second order information on the "derivatives" of the function to be optimized. These in particular subsume all past results. The main technical work goes into identifying the key parameters of the underlying group actions which control convergence to the optimum in each of these methods. These non-commutative analogues of "smoothness" are far more complex and require significant algebraic and analytic machinery. Despite this complexity, the way in which these parameters control convergence in both methods is quite simple and elegant. We show how to bound these parameters and hence obtain efficient algorithms for null cone membership in several concrete situations. Our work points to intriguing open problems and suggests further research directions.

*Institut für Mathematik, Technische Universität Berlin, pbuerg@math.tu-berlin.de. Supported by DFG grant BU 1371 2-2 and by the ERC under the European's Horizon 2020 research and innovation programme (grant agreement no. 787840).

†Department of Mathematics, Rutgers University and MIT, franks@mit.edu. Partially supported by Simons Foundation award 32622.

‡Microsoft Research India, garga@microsoft.com.

§Cheriton School of Computer Science, University of Waterloo, rafael@uwaterloo.ca. Research done while Rafael was at Princeton, University of Toronto and at the Simons Institute for the Theory of Computing.

¶Korteweg-de Vries Institute for Mathematics, Institute for Theoretical Physics, Institute for Logic, Language, and Computation, QuSoft, University of Amsterdam, m.walter@uva.nl. Supported by NWO Veni grant no. 680-47-459 and NWO grant OCENW.KLEIN.267.

‖Institute for Advanced Study, Princeton, avi@ias.edu. Supported by NSF grants CCF-1412958 and CCF-1900460.