

Quantum Gravity in the Lab: Teleportation by Size and Traversable Wormholes

Adam R. Brown,[1,2] Hrant Gharibyan,[2,3] Stefan Leichenauer,[1] Henry W. Lin,[1,4] Sepehr Nezami,[5,1,2] Grant Salton,[6,7,5,2] Leonard Susskind,[1,2] Brian Swingle,[8] and Michael Walter[9]

[1]*Google, Mountain View, CA 94043, USA*
[2]*Department of Physics, Stanford University, Stanford, CA 94305, USA*
[3]*Institute for Quantum Information and Matter, Caltech, Pasadena CA 91125, USA*
[4]*Physics Department, Princeton University, Princeton, NJ 08540, USA*
[5]*Institute for Quantum Information and Matter, Caltech, Pasadena, CA 91125, USA*
[6]*Amazon Quantum Solutions Lab, Seattle, WA 98170, USA*
[7]*AWS Center for Quantum Computing, Pasadena, CA 91125, USA*
[8]*Condensed Matter Theory Center, Joint Center for Quantum Information and Computer Science, Maryland Center for Fundamental Physics, and Department of Physics, University of Maryland, College Park, MD 20742, USA*
[9]*Korteweg-de Vries Institute for Mathematics, Institute for Theoretical Physics, Institute for Logic, Language and Computation & QuSoft, University of Amsterdam, The Netherlands*

arXiv:1911.06314v2 [quant-ph] 1 Feb 2021

With the long-term goal of studying models of quantum gravity in the lab, we propose holographic teleportation protocols that can be readily executed in table-top experiments. These protocols exhibit similar behavior to that seen in the recent traversable wormhole constructions of [1, 2]: information that is scrambled into one half of an entangled system will, following a weak coupling between the two halves, unscramble into the other half. We introduce the concept of *teleportation by size* to capture how the physics of operator-size growth naturally leads to information transmission. The transmission of a signal *through* a semi-classical holographic wormhole corresponds to a rather special property of the operator-size distribution we call *size winding*. For more general systems (which may not have a clean emergent geometry), we argue that imperfect size winding is a generalization of the traversable wormhole phenomenon. In addition, a form of signaling continues to function at high temperature and at large times for generic chaotic systems, even though it does *not* correspond to a signal going through a geometrical wormhole, but rather to an interference effect involving macroscopically different emergent geometries. Finally, we outline implementations feasible with current technology in two experimental platforms: Rydberg atom arrays and trapped ions.