arXiv:2002.02743v1 [quant-ph] 7 Feb 2020

The Haemers bound of noncommutative graphs

Sander Gribling* Yinan Li*

Abstract

We continue the study of the quantum channel version of Shannon's zero-error capacity problem. We generalize the celebrated Haemers bound to noncommutative graphs (obtained from quantum channels). We prove basic properties of this bound, such as additivity under the direct sum and submultiplicativity under the tensor product. The Haemers bound upper bounds the Shannon capacity of noncommutative graphs, and we show that it can outperform other known upper bounds, including noncommutative analogues of the Lovász theta function (Duan–Severini–Winter, *IEEE Trans. Inform. Theory*, 2013 and Boreland–Todorov–Winter, *arXiv*, 2019).

Keywords: Haemers bound – Noncommutative graphs – Quantum channels – Shannon capacity – Zero-error information theory

1 Introduction

The celebrated *Shannon capacity* of a graph G is defined as

$$\Theta(G) = \sup_k \sqrt[k]{\alpha(G^{\boxtimes k})} = \lim_{k\to\infty} \sqrt[k]{\alpha(G^{\boxtimes k})}, \tag{1}$$

where $\alpha(G)$ denotes the independence number of G and $\boxtimes$ denotes the strong graph product [Sha56]. The logarithm of $\Theta(G)$ characterizes the amount of information that can be transmitted through a classical communication channel, with zero error, where we allow an arbitrary number of uses of the channel and we measure the average amount of information transmitted per use of the channel. (The graph G is the so-called *confusability graph* associated to the channel, see Section 2.1.) The Shannon capacity is not known to be *computable*: Even though computing the independence number is NP-complete [Kar72], there exist graphs whose Shannon capacities are not achieved by taking the strong graph product with itself *finitely* many times [GW90].

To upper bound the Shannon capacity, Lovász introduced the celebrated theta function [Lov79], which can be cast as a semidefinite program and can be used to compute, e.g., $\Theta(C_5)$. Lovász posed the question whether the Shannon capacity equals the theta function in general, which has been refuted by Haemers: He introduced another upper bound on the Shannon capacity, now known as the Haemers bound, which can be strictly smaller than the theta function on some graphs [Hae78, Hae79].

Instead of a *classical* communication channel, we could also consider a *quantum* communication channel. Doing so leads to quantum information analogues of the aforementioned questions, the study of which was systematically initiated by Duan, Severini and Winter [DSW13]. In Section 2.1 we show how the quantum setting generalizes the classical setting, which also motivates the following definitions. To (the Choi-Kraus representation of) a quantum channel $\Phi(A) = \sum_{k=1}^m E_k A E_k^\dagger$ ($\forall\ A \in$

*Centrum Wiskunde & Informatica (CWI) and QuSoft, Science Park 123, 1098XG, Amsterdam, Netherlands. {gribling,Yinan.Li}@cwi.nl.