arXiv:2002.08070v2 [quant-ph] 14 Jul 2020

Complete classification of trapping coins for quantum walks on the 2D square lattice

B. Kollár[1], A. Gilyén[1,4,5], I. Tkáčová[2], T. Kiss[1], I. Jex[3] and M. Štefaňák[3]
[1] *Wigner RCP, Konkoly-Thege M. u. 29-33, H-1121 Budapest, Hungary*
[2] *Department of Physics, VŠB Technical University of Ostrava, 17. Listopadu 15, 70800 Ostrava-Poruba, Czech Republic*
[3] *Department of Physics, Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague, Břehová 7, 115 19 Praha 1 - Staré Město, Czech Republic*
[4] *QuSoft / CWI, Science Park 123, 1098 XG Amsterdam, the Netherlands and*
[5] *Institute for Quantum Information and Matter, California Institute of Technology, 1200 E California Blvd, Pasadena, CA 91125, US*
(Dated: July 15, 2020)

One of the unique features of discrete-time quantum walks is called trapping, meaning the inability of the quantum walker to completely escape from its initial position, albeit the system is translationally invariant. The effect is dependent on the dimension and the explicit form of the local coin. A four state discrete-time quantum walk on a square lattice is defined by its unitary coin operator, acting on the four dimensional coin Hilbert space. The well known example of the Grover coin leads to a partial trapping, i.e., there exists some escaping initial state for which the probability of staying at the initial position vanishes. On the other hand, some other coins are known to exhibit strong trapping, where such escaping state does not exist. We present a systematic study of coins leading to trapping, explicitly construct all such coins for discrete-time quantum walks on the 2D square lattice, and classify them according to the structure of the operator and the manifestation of the trapping effect. We distinguish three types of trapping coins exhibiting distinct dynamical properties, as exemplified by the existence or non-existence of the escaping state and the area covered by the spreading wave-packet.

I. INTRODUCTION

Discrete time quantum walks [1, 2] are non-trivial generalizations of discrete-time classical walks. These elementary constructs follow the rules of quantum mechanics and became versatile tools in various field of physics (for reviews see [3–8]). The motion of a single excitation in a solid state, the spreading of quantum information in a quantum network and even quantum computation can be modeled by quantum walks [9].

Recently, quantum walks have attracted interest as simple quantum simulators, modeling the behavior of quantum particles under various conditions: the effect of decoherence [10, 11], electric fields [12, 13], and percolation [14–20] were studied in detail. During the last decade, a number of state-of-the-art experiments [7, 13, 21–34] were performed validating the theoretical results and also benchmarking the achievable degree of quantum control and visibility.

Quantum walks serve as an elementary model for transport phenomena in physical systems. Spreading properties of quantum walks significantly differ from classical random walks. They can spread faster, thus speeding up random walk based search [35–38], leading to a number of possible applications in quantum information [39]. Nevertheless, there might be vertices which are almost never reached by the walker due to destructive interference, leading to infinite hitting times even for finite graphs [40, 41]. However, for the initial vertex the expected return time is always finite for a finite graph, as follows from a general results for discrete-time unitary evolution [42]. The expected return time to the exact initial state (state recurrence) is an integer [42]. This holds even for iterated open quantum evolution, provided it is described by a unital quantum channel [43]. The investigation was later extended to a broader class of iterated open quantum dynamics [44] and the result can be understood as a generalization of the Kac lemma [45]. We note that in the case of subspace recurrence the expected return time is a rational number [46].

Quantum walks are known for their typical ballistic spreading [47]. However, for a quantum walk on a two-dimensional lattice there exist some coins which lead to limited spreading for some initial states. In particular, for a Grover coin one can observe a probability peak situated at the origin of the walk, discovered by Mackay, Bartlett, Stephenson and Sanders [48]. We will refer to this property as trapping. Let us note that the term localization is sometimes used for the same effect in the literature, however localization [49] is often used in a different context, e.g., in Anderson localization – a phenomenon arising from spatial randomness [24, 28, 50–56], exponential localization of topologically protected states [26, 57, 58], or oscillatory localization [59]. The effect of trapping by a Grover coin for discrete-time quantum walks on 2D integer lattice was rigorously proven by Inui, Konishi and Konno [60]. Implications of trapping for stationary measures of quantum walks were discussed in [61, 62]. We note that trapping is not limited to the square lattice, but can be found in any d-dimensional lattice. For quantum walks on a line non-trivial trapping coins need to have at least three-dimensions [63]. Trapping coins of dimensions greater than 3 were also identified [64] and further studied in [65, 66]. Several extensions of the three-state Grover coin featuring trapping were introduced [67] and investigated in detail [61, 68–