

2018–
2021



Witnessing Entanglement in Experiments with Correlated Noise

Bas Dirkse,[1,2,3] Matteo Pompili,[1,2] Ronald Hanson,[1,2] Michael Walter,[3,4] and Stephanie Wehner[1,2]

[1] *QuTech, Delft University of Technology, The Netherlands*
[2] *Kavli Institute of Nanoscience, Delft University of Technology, The Netherlands*
[3] *QuSoft, University of Amsterdam, The Netherlands*
[4] *Korteweg-de Vries Institute for Mathematics, Institute for Theoretical Physics, Institute for Logic, Language, and Computation, University of Amsterdam, The Netherlands*

(Dated: April 29, 2020)

The purpose of an entanglement witness experiment is to certify the creation of an entangled state from a finite number of trials. The statistical confidence of such an experiment is typically expressed as the number of observed standard deviations of witness violations. This method implicitly assumes that the noise is well-behaved so that the central limit theorem applies. In this work, we propose two methods to analyze witness experiments where the states can be subject to arbitrarily correlated noise. Our first method is a *rejection experiment*, in which we certify the creation of entanglement by rejecting the hypothesis that the experiment can only produce separable states. We quantify the statistical confidence by a p-value, which can be interpreted as the likelihood that the observed data is consistent with the hypothesis that only separable states can be produced. Hence a small p-value implies large confidence in the witnessed entanglement. The method applies to general witness experiments and can also be used to witness genuine multipartite entanglement. Our second method is an *estimation experiment*, in which we estimate and construct confidence intervals for the average witness value. This confidence interval is statistically rigorous in the presence of correlated noise. The method applies to general estimation problems, including fidelity estimation. To account for systematic measurement and random setting generation errors, our model takes into account device imperfections and we show how this affects both methods of statistical analysis. Finally, we illustrate the use of our methods with detailed examples based on a simulation of NV centers.

arXiv:2002.12400v2 [quant-ph] 28 Apr 2020

I. INTRODUCTION

Entanglement is a fundamental property of quantum mechanical systems [1] and an important resource for many quantum information processing tasks. In quantum computing, coherently creating entanglement between several qubits is necessary for computational speedups [1–3]. In quantum networks, remote entanglement is an essential resource for quantum cryptography [4–6] and distributed computing applications [7–9]. Entanglement also plays a crucial role in quantum sensing and metrology [10–12], enabling more precise measurement of physical quantities. With the rapid experimental advances in the manipulation and control of quantum systems, much progress had been made towards the generation of entangled states in various physical platforms [13–19]. Yet, the creation of high-quality many-body entanglement is still a significant challenge. It is therefore important to have good tools to certify the creation of entanglement. The main tools used for this purpose are entanglement witnesses.

An *entanglement witness* is an observable W on a quantum system that can certify entanglement of a state ρ^* under investigation [20]. By definition, the witness W satisfies

$$\mathrm{Tr}[\rho W] \geq 0 \quad \forall \rho \in \mathcal{S}, \tag{1}$$

for all separable states $\rho \in \mathcal{S}$. As a consequence, it can be used to certify entanglement: If a state ρ^* has negative witness expectation value, $\mathrm{Tr}[W\rho^*] < 0$, then it is necessarily entangled. If the expectation value is non-negative, the test is inconclusive (the state can either be separable or not). The witness method applies more generally than just for separating entangled from separable states. If $\mathcal{S}$ is an arbitrary convex set of states and $\rho^* \notin \mathcal{S}$, then there always exists a witness W such that Eq. (1) holds, while $\mathrm{Tr}[W\rho^*] < 0$. For example, a witness can be used to certify that states are genuinely multipartite entangled. Geometrically, the witness W can be interpreted as a hyperplane that separates the convex set $\mathcal{S}$ from the state $\rho^* \notin \mathcal{S}$. This is illustrated in Fig. 1. In general, finding an appropriate witness W for a state ρ^* is a difficult problem that has been studied extensively in literature [21–23]. For the remainder of this article, we will assume that W is chosen and fixed.




FIG. 1. Geometric interpretation of a witness W as a hyperplane that separates the state ρ^* from the convex set $\mathcal{S}$. If $\mathcal{S}$ is the set of separable states, then W certifies that ρ^* is entangled.

Often, a witness W is a non-local observable of the system for which entanglement is to be certified. Such measurements are typically hard to perform, particularly in a network setting. Therefore, in experiments, W is usually decomposed into a sum of locally measurable observables which are then measured individually on the constituent