


SciPost Physics Submission

Bosonic entanglement renormalization circuits from wavelet theory

Freek Witteveen[1,*], Michael Walter[1,2]

1 Korteweg-de Vries Institute for Mathematics and QuSoft, University of Amsterdam, Netherlands
2 Institute for Theoretical Physics and Institute for Language, Logic, and Computation, University of Amsterdam, Netherlands
* f.g.witteveen@uva.nl

April 20, 2021

arXiv:2004.11952v3 [quant-ph] 24 Jun 2021

Abstract

Entanglement renormalization is a unitary real-space renormalization scheme. The corresponding quantum circuits or tensor networks are known as MERA, and they are particularly well-suited to describing quantum systems at criticality. In this work we show how to construct Gaussian bosonic quantum circuits that implement entanglement renormalization for ground states of arbitrary free bosonic chains. The construction is based on wavelet theory, and the dispersion relation of the Hamiltonian is translated into a filter design problem. We give a general algorithm that approximately solves this design problem and provide an approximation theory that relates the properties of the filters to the accuracy of the corresponding quantum circuits. Finally, we explain how the continuum limit (a free bosonic quantum field) emerges naturally from the wavelet construction.

Contents

1 Introduction **2**
1.1 Main results 3
1.2 Organization of the paper 4

2 Perfect reconstruction and biorthogonal filters **5**

3 Entanglement renormalization and filter design **6**
3.1 Derivation of filter condition 7

4 Entanglement renormalization circuits **9**

5 The continuum limit **10**
5.1 Scaling and wavelet functions 12
5.2 Entanglement renormalization for the massless boson 13
5.3 The massive bosonic field 15
5.4 Other perspectives 15