



A High-Fidelity Method for a Single-Step N-bit Toffoli Gate in Trapped Ions

Juan Diego Arias Espinoza,[1] Koen Groenland,[2] Matteo Mazzanti,[1] Kareljan Schoutens,[2,3] and Rene Gerritsma[1]

[1] *Van der Waals-Zeeman Institute, Institute of Physics, University of Amsterdam, 1098 XH Amsterdam, the Netherlands*
[2] *QuSoft Science Park 123, 1098 XG Amsterdam, the Netherlands*
[3] *Institute for Theoretical Physics, Institute of Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, the Netherlands*

(Dated: October 19, 2020)

Conditional multi-qubit gates are a key component for elaborate quantum algorithms. In a recent work, Rasmussen *et al.* (Phys. Rev. A 101, 022308) proposed an efficient single-step method for a prototypical multi-qubit gate, a Toffoli gate, based on a combination of Ising interactions between control qubits and an appropriate driving field on a target qubit. Trapped ions are a natural platform to implement this method, since Ising interactions mediated by phonons have been demonstrated in increasingly large ion crystals. However, the simultaneous application of these interactions and the driving field required for the gate results in undesired entanglement between the qubits and the motion of the ions, reducing the gate fidelity. In this work, we propose a solution based on adiabatic switching of these phonon mediated Ising interactions. We study the effects of imperfect ground state cooling, and use spin-echo techniques to undo unwanted phase accumulation in the achievable fidelities. For gates coupling to all axial modes of a linear crystal, we calculate high fidelities ($>$ 99%) N-qubit rotations with N = 3-7 ions cooled to their ground state of motion and a gate time below 1 ms. The high fidelities obtained also for large crystals could make the gate competitive with gate-decomposed, multi-step variants of the N-qubit Toffoli gate, at the expense of requiring ground state cooling of the ion crystal.

arXiv:2010.08490v1 [quant-ph] 16 Oct 2020

I. INTRODUCTION

Quantum computers promise dramatic speedups in a variety of disciplines [1–5], but remain challenging to scale up in practice. A major obstacle to executing elaborate quantum algorithms, is the need for gates that act conditionally on a large number of qubits. The prototypical example of such a gate is the N-qubit Toffoli gate, which flips a single 'target' qubit if and only if all $N-1$ 'control' qubits are in the state $|1\rangle$. Even though quantum devices with over 50 qubits have been reported [6, 7], the largest Toffoli gate ever performed is, to our best knowledge, the case $N = 4$ [8]. This gap is surprising, because Toffoli gates (or equivalents) are essential ingredients of many basic computation steps, such as elementary arithmetic [9–11], error correction [12], and the Grover diffusion operator [13].

Two different strategies exist to implement Toffoli gates. The first consists of decomposing a single N-qubit Toffoli gate into a circuit consisting of one- and two-qubit gates [14–16] or multiqubit gates, such as the Mølmer-Sørensen gate in trapped ions [17–19]. The second approach is to perform the gate in a single step using interactions that are native to the specific platform [20–23]. In particular, a recent proposal [23] has demonstrated that by exploiting systems with an all-to-all Ising interaction in combination with a drive field on a single target qubit an i-Toffoli gate can be implemented. This gate differs only from the regular Toffoli by a phase $+i$ on the target qubit.

Trapped ions are a natural candidate to implement this proposal, as intrinsic Ising interactions have been demonstrated in increasingly large ion crystals [6, 17, 24–26]. Moreover, quantum operations have been demonstrated [27, 28] with fidelities higher than 99.9%. Ising interactions generally arise from qubit-phonon couplings $\hat{H}_{\text{q-ph}}$ generated from state-dependent laser-induced forces on the ions. Combining this mechanism with the driving field $\hat{H}_{\text{drive}}$ required for an i-Toffoli gate poses a problem, as both process do not commute i.e. $[\hat{H}_{\text{q-ph}}, \hat{H}_{\text{drive}}] \neq 0$. As a result, the qubit states and the motion of the ions remain entangled at the end of the gate sequence, which leads to fidelity loss. This effect could be mitigated by restricting the strength of the spin-phonon coupling such that the phonons are only virtually excited [26]. However, limiting the strength of the Ising interactions leads to undesirably long gate times.

In this work, we show that this residual qubit-phonon entanglement can be suppressed by adiabatic ramping of $\hat{H}_{\text{q-ph}}$. In this way, the i-Toffoli gate operates on the dressed eigenstates of $\hat{H}_{\text{q-ph}}$, that are adiabatically connected to the Fock eigenstates of the non-interacting system. The benefit of this approach is that the effective Ising interaction strength does not have to be limited to the regime of virtual phonon excitation. We show that high-fidelity $\bar{F} > 99\%$, single step, i-Toffoli gates should be possible with up to 7 ions at gate times ~ 600 µs.

We start in Sec. II with the derivation of the model for a N-qubit i-Toffoli gate for a system of trapped ions and introduce our proposal for adiabatic preparation of dressed states. In Sec. III we analyze the results of numerical simulations for a linear 3 crystal and consider the role of inhomogeneous Ising interactions mediated by multiple phonon modes. We discuss the implementation of a method based on multi-frequency laser fields [29] to eliminate undesired phases originating from these inho-