


State selective cooling of SU(N) Fermi-gases

Aaron Merlin Müller,[1,2] Miklós Lajkó,[2] Florian Schreck,[3,4] Frédéric Mila,[2] and Jiří Minář[5,4]

[1] *Institute for Theoretical Physics, ETH Zürich, 8093 Zürich, Switzerland*
[2] *Institute of Physics, École Polytechnique Fédérale de Lausanne (EPFL), CH-1015 Lausanne, Switzerland*
[3] *Van der Waals-Zeeman Institute, Institute of Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, the Netherlands*
[4] *QuSoft, Science Park 123, 1098 XG Amsterdam, the Netherlands*
[5] *Institute for Theoretical Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, the Netherlands*

(Dated: June 15, 2021)

We investigate a species selective cooling process of a trapped SU(N) Fermi gas using entropy redistribution during adiabatic loading of an optical lattice. Using high-temperature expansion of the Hubbard model, we show that when a subset $N_A < N$ of the single-atom levels experiences a stronger trapping potential in a certain region of space, the dimple, it leads to improvement in cooling as compared to a SU(N_A) Fermi gas only. We show that optimal performance is achieved when all atomic levels experience the same potential outside the dimple and we quantify the cooling for various N_A by evaluating the dependence of the final entropy densities and temperatures as functions of the initial entropy. Furthermore, considering ^{87}Sr and ^{173}Yb for specificity, we provide a quantitative discussion of how the state selective trapping can be achieved with readily available experimental techniques.

arXiv:2102.08656v2 [cond-mat.quant-gas] 12 Jun 2021

I. INTRODUCTION

In recent years, there has been a considerable effort in experimental control of ultracold Fermi gases with the aim of realizing models of strongly interacting electrons, in particular the Hubbard model, upon loading the atoms into a deep optical lattice [1]. Of particular interest are ultracold quantum degenerate Fermi gases with nuclear spin I that is decoupled from the electronic spin, such as ^{173}Yb [2–4] or ^{87}Sr [5–7], which feature $N = 2I + 1$ hyperfine states in the ground state manifold.

The SU(N) Fermi gases have attracted considerable attention as they allow for SU(N) generalizations of the Hubbard model [8] and can host a plethora of exotic phases including various spin orders and liquids [9–14], Mott insulator-metal transitions and crossovers [15, 16], valence bond solids and semimetals [17, 18], unconventional superconductors [19] or collective motional modes [20]. Remarkably, some of these scenarios have been probed also experimentally for $N > 2$ [21–25]. The limit of large interaction gives rise to SU(N) magnetism [26, 27], where the system can be effectively described in terms of a Heisenberg model. This stimulated theoretical investigations using representation theory [28–31], variational approaches [32] or large scale simulations at finite temperature [33]. Furthermore, depending on N and the lattice geometry, the Heisenberg Hamiltonians can be linked to Wess-Zumino-Witten models when at a critical point [34, 35], feature chiral spin liquids [36] and magnetic orders such as generalized valence bond solids [37], plaquette [38, 39], Néel and stripelike long-range [40, 41] or antiferromagnetic order [42].

To observe these magnetic orders the atoms need to be cooled to temperatures below the superexchange energy $4t^2/U$, where t and U are the tunneling rate and interaction strength of the parent Hubbard model. Here, a promising approach is based on an (adiabatic) entropy redistribution akin to the Pomeranchuk effect in solid helium [43]. For cold atoms in optical lattices this effect has been studied theoretically by means of dynamical mean field theory in [44], where the entropy was removed from a certain region - a dimple - by appropriately shaping the trapping potential. In the context of SU(N) fermions, Refs. [45, 46] have studied the enhancement of the cooling due to higher N (see also [47] for adiabatic cooling of interacting and [48, 49] of non-interacting fermions). Pomeranchuk and dimple cooling were experimentally demonstrated in [50] and [51] respectively, leading to an antiferromagnetic order [23, 52] with [51] reporting the final temperature of $T/t = 0.25$ (see also [53] for experimental realization of short-ranged antiferromagnetic order, [22] for probing the Mott-insulator transition, and [25] for the thermodynamics of the interacting SU(N) Fermi gas).

Motivated by these developments, in this work we study the effect of adiabatically loading an initally harmonically trapped SU(N) Fermi gas into a deep optical lattice in a species selective way: specifically, we consider a bi-partition of the atomic levels in two families, A and B such that $N = N_A + N_B$ and an optical potential which forms a dimple for only the A-family (hereafter we refer to the different atomic levels as colors). Using the high-temperature expansion of the Hubbard model we compute the entropy density and show that this results in further enhancement of the cooling of the Mott-insulating state of the A-family atoms in the dimple compared to a SU(N_A) Fermi gas only.

The paper is structured as follows. In Sec. II we describe the model and methodology, present the results in Sec. III, discuss a possible experimental implementation in Sec. IV and conclude in Sec. V.