



Engineering spin-spin interactions with optical tweezers in trapped ions

J. D. Arias Espinoza,[1] M. Mazzanti,[1] K. Fouka,[1] R. X. Schüssler,[1]
Z. Wu,[1] P. Corboz,[2,3] R. Gerritsma,[1] and A. Safavi-Naini[2,3]

[1] *Van der Waals-Zeeman Institute, Institute of Physics, University of Amsterdam, 1098 XH Amsterdam, the Netherlands*
[2] *QuSoft, Science Park 123, 1098 XG Amsterdam, the Netherlands*
[3] *Institute for Theoretical Physics, Institute of Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, the Netherlands*

arXiv:2103.10425v3 [quant-ph] 23 Mar 2021

We propose a new method for generating programmable interactions in one- and two-dimensional trapped-ion quantum simulators. Here we consider the use of optical tweezers to engineer the sound-wave spectrum of trapped ion crystals. We show that this approach allows us to tune the interactions and connectivity of the ion qubits beyond the power-law interactions accessible in current setups. We demonstrate the experimental feasibility of our proposal using realistic tweezer settings and experimentally relevant trap parameters to generate the optimal tweezer patterns to create target spin-spin interaction patterns in both one- and two-dimensional crystals. Our approach will advance quantum simulation in trapped-ion platforms as it allows them to realize a broader family of quantum spin Hamiltonians.

I. INTRODUCTION

Trapped ions are one of the leading platforms for quantum computation and quantum simulation [1–3]. Numerous experiments have demonstrated the ability of the analog trapped-ion quantum simulator to emulate the dynamics of quantum magnetism models [3–7] and to study the dynamics of quantum information and quantum entanglement [8–10].

One of the main advantages of the trapped-ion quantum simulators is the tunability of the interaction range, as well as the ability to realize one- and two-dimensional (1D and 2D) systems. Hence, this platform provides an ideal setup in which one can explore the interplay of interaction range and dimensionality in the dynamics of quantum information, entanglement, and speed of thermalization [8, 9, 11, 12], while simultaneously allowing one to simulate models relevant to condensed matter physics that are beyond the state-of-the-art numerical methods.

However, the current simulator setups do not offer enough versatility to explore the above questions. This is because, theoretically, the types of engineered interactions which can be realized in these systems is limited to those with power-law decay, $1/r^{\xi}$, where r is the separation between two ions and $0 \leq \xi \leq 3$ [13]. In order to understand the source of this limitation, we note that the ion-ion interactions in the simulator are phonon-mediated and depend on the spectrum and structure of the collective vibrational modes of the ion crystal [14]. Furthermore, considering experimental constraints such as laser power and decoherence rates, the range of interactions tends to be even more limited, and most experiments are operated with $0 \leq \xi \leq 1.5$ [11, 13, 15].

In the following, we illustrate a new approach for realizing a highly tunable trapped-ion simulator in terms of connectivity, range, and sign of the interactions in both linear (or 1D) and triangular (2D) ion crystals in Paul traps [16–20]. We use optical tweezers to manipulate the frequencies and structure of the collective vibrational modes of the crystal. The triangular crystal structure in 2D makes it a natural platform for implementing quantum simulation of frustrated spin systems [21, 22] and there has been rapid experimental progress in this area in recent years [23–25].

II. EFFECTIVE ISING INTERACTIONS

The spin-spin interactions in an ion crystal are generated when the electronic state of each ion is coupled to the phonon modes of the ion crystal by applying a state-dependent force generated by a pair of counter-propagating laser beams. The resulting Hamiltonian, in the Lamb-Dicke (LD) limit and in the interaction picture with respect to $\hat{H}_0 = \sum_{m=1}^{N} \hbar\omega_m \hat{n}_m$, is given by

$$\hat{H}_{\mathrm{I}} = -i \sum_{j=1}^{N} g \cos(\mu t) \hat{\sigma}_j^x \times \sum_{m=1}^{N} b_{jm} \sqrt{\frac{\hbar}{2M\omega_m}} \left(\hat{a}_m e^{-i\omega_m t} - \hat{a}_m^{\dagger} e^{i\omega_m t} \right). \tag{1}$$

where where μ is the beat-note frequency between the two counter-propagating bichromatic lasers used to generate the state-dependent force and g is the interaction strength of the laser which is assumed to be homogeneous throughout the crystal. Here $\hat{a}_m$ ($\hat{a}_m^{\dagger}$) is the annihilation (creation) operator for a phonon in mode m, $\hat{n}_m = \hat{a}_m^{\dagger}\hat{a}_m$, and b_{jm} is the amplitude of the corresponding eigenvector at the position of ion j. The LD parameter is given by $\eta_j^{(m)} \equiv k_q b_{jm} \sqrt{\frac{\hbar}{2M\omega_m}}$ where k_q is the wavevector for the Raman beam pair and M the ion mass.

The form of the Hamiltonian in Eq. (1) allows us to find an explicit expression for the propagator governing