

2022–
2024

Entanglement Classification via Single Entanglement Measure

Adam Burchardt,[1,*] Gonçalo M. Quinta,[2] and Rui André[3]

[1] QuSoft, CWI and University of Amsterdam, Science Park 123, 1098 XG Amsterdam, the Netherlands
[2] Instituto de Telecomunicações, Physics of Informations and Quantum Technologies Group, Lisboa, Portugal
[3] Centro de Astrofísica e Gravitação - CENTRA,
Departamento de Física, Instituto Superior Técnico - IST,
Universidade de Lisboa - UL, Av. Rovisco Pais 1, 1049-001 Lisboa, Portugal

We show that a single polynomial entanglement measure is enough to verify equivalence between generic n-qubit states under Stochastic Local Operations with Classical Communication (SLOCC). SLOCC operations may be represented geometrically by Möbius transformations on the roots of the entanglement measure on the Bloch sphere. Moreover, we show how the roots of the 3-tangle measure classify 4-qubit generic states, and propose a method to obtain the normal form of a 4-qubit state which bypasses the possibly infinite iterative procedure.

arXiv:2106.00850v3 [quant-ph] 4 Apr 2024

I. INTRODUCTION

Quantum entanglement is one of the key manifestations of quantum mechanics and the main resource for technologies founded on quantum information science. In particular, quantum states with non-equivalent entanglement represent distinct resources which may be useful for different protocols. The idea of clustering states into classes exhibiting different qualities under quantum information processing tasks resulted in their classification under stochastic local operations assisted by classical communication (SLOCC). Such a classification was successfully presented for two, three and four qubits [1–4]. However, the full classification of larger systems is completely unkown. Even the much simpler problem of detecting if two n-qubit states ($n > 4$) are SLOCC-equivalent is, in general, quite demanding [5–8].

Among several approaches to the entanglement quantification and classification problem, a particularly useful one is via *SL-invariant polynomial* (SLIP) measures. Well-known examples are concurrence and 3-tangle, which measure the 2-body and 3-body quantum correlations of the system [9, 10]. SLIP measures provide not only a convenient method for entanglement classification but also its practical detection. Indeed, it was shown that almost all SLOCC equivalence classes can be distinguished by ratios of such measures [3]. Any given two n-qubit states are then SLOCC-equivalent if a *complete set of SLIP measures* has the same values for both of them [5]. For more than four qubits, however, the size of such a set grows exponentially, making it intractable to use this approach to discriminate SLOCC-equivalent states with more than four qubits [11].

In this paper, we show that, contrary to intuition, a single SLIP measure is enough to verify the SLOCC-equivalence between any two generic pure n-qubit states. By generic state, we mean the set of all pure states except of the measure zero subset with respect to the Haar measure. We achive this by using a mathematical trick, in particular, we use a SLIP measure defined for n qubit systems to verify equivalence of systems of $n+1$ quibits. In essence, we look how the roots of the SLIP measure for those states behave under SLOCC. We show that if the states are SLOCC-equivalent, then the roots of the SLIP measure for each state must be related by a Möbius transformation, which is straightforward to verify. In particular, we use this procedure to show that the 3-tangle measure is enough to discriminate between generic 4-qubit states. Finally, we show how one may use the roots of a SLIP entanglement measure to obtain the normal form of a 4-qubit state which bypasses the possibly infinite iterative standard procedure.

II. POLYNOMIAL INVARIANT MEASURES

An *entanglement measure* is a function $E(|\psi\rangle)$ defined for pure states of n qubits which vanishes on the set of separable states. One of the desired features of entanglement measures is invariance under SLOCC operations. Mathematically, a SLOCC operation might be uniquely determined by the action of local invertible operators $L \in \mathrm{SL}(2,\mathbb{C})^{\otimes n}$ [1]. An entanglement measure E defined for all pure states of n qubits is called a *SL-invariant polynomial of homogeneous degree h* if it is homogeneous polynomial of degree h in the state coefficients, an it is invariant under any local operation $\mathcal{O} = \mathcal{O}_1 \otimes \cdots \otimes \mathcal{O}_n$, where $\mathcal{O}_i \in \mathrm{SL}(2,\mathbb{C})$. It is easy to see that those two conditions are equivalent to the fact that E satisfies

$$E\big(\kappa\,\mathcal{O}\,|\psi\rangle\big) = \kappa^h E\big(\,|\psi\rangle\big)$$

for each real constant $\kappa > 0$ and invertible linear operator $\mathcal{O}$ [1, 3, 12]. Such polynomial will be denoted by SLIP^h_n, where the upper index indicates the degree of the polynomial and the lower index is related to the number of qubits. Well-known examples are concurrence and 3-tangle [9, 10]. Both of those measures, concurrence and 3-tangle, can be written as the absolute value of the anti-linear expectation value of simple operators. Indeed, for a two-qubit pure state $|\psi\rangle = c_{00}\,|00\rangle + c_{01}\,|01\rangle + c_{10}\,|10\rangle + c_{11}\,|11\rangle$ its concurrence reads

* adam.burchardt.uam@gmail.com