



The Effect of Micromotion and Local Stress in Quantum simulation with Trapped Ions in Optical Tweezers

Liam Bond,[1,2] Lisa Lenstra,[3] Rene Gerritsma,[1,3] and Arghavan Safavi-Naini[1,2]

[1] *QuSoft, Science Park 123, 1098 XG Amsterdam, the Netherlands*
[2] *Institute for Theoretical Physics, Institute of Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, the Netherlands*
[3] *Van der Waals-Zeeman Institute, Institute of Physics, University of Amsterdam, 1098 XH Amsterdam, Netherlands*

(Dated: March 1, 2022)

The ability to program and control interactions provides the key to implementing large-scale quantum simulation and computation in trapped ion systems. Adding optical tweezers, which can tune the phonon spectrum and thus modify the phonon-mediated spin-spin interaction, was recently proposed as a way of programming quantum simulators for a broader range of spin models [Arias Espinoza et al., Phys. Rev. A **103**, 052437]. In this work we study the robustness of our findings in the presence of experimental imperfections: micromotion, local stress, and intensity noise. We show that the effects of micromotion can be easily circumvented when designing and optimizing tweezer patterns to generate a target interaction. Furthermore, while local stress, whereby the tweezers apply small forces on individual ions, may appear to enable further tuning of the spin-spin interactions, any additional flexibility is negligible. We conclude that optical tweezers are a useful method for controlling interactions in trapped ion quantum simulators in the presence of micromotion and imperfections in the tweezer alignment, but require intensity stabilization on the sub-percent level.

arXiv:2202.13681v1 [quant-ph] 28 Feb 2022

I. INTRODUCTION

Trapped ions are at the forefront of both digital and analog quantum simulation [1–3]. On the digital side, trapped-ions are the building blocks of the highest fidelity two-qubit universal gates [4–6], and the recent demonstration of on-the-fly quantum error correction adds to the robustness of this architecture [7]. On the analog side, they have been used to emulate the dynamics and prepare the ground states of quantum magnets, as well as study the dynamics of quantum correlations, quantum information and entanglement in the presence of engineered, variable-range interactions [8–12].

Trapped-ion quantum simulators allow one to engineer power-law spin-spin interactions which decay as $1/r^\alpha$ where $0 < \alpha < 3$ and r is the distance between two ions. This is the direct result of the mechanism behind the interactions. The inter-ion interactions are phonon-mediated and as such depend on the spectrum and structure of the collective vibrational modes of the ion crystal [13, 14]. So far experimental efforts utilizing trapped-ions as analog simulators have been restricted to the aforementioned power-law interactions. Recently it has been shown that the addition of optical tweezers to the typical trapped-ion platform produces a highly tunable quantum simulator in terms of connectivity, range, and sign of the interactions in both linear (or 1D) and triangular (2D) ion crystals in Paul traps [15–18]. If a target interaction matrix passes our feasibility criterion, we search for the optimal optical tweezer pattern to manipulate the frequencies and structure of the collective vibrational modes of the crystal.

In this work we study the robustness of our scheme in presence of typical experimental imperfections: micromotion, tweezer misalignment, and tweezer intensity noise. In Section. II we review the radio-frequency (r.f.) Paul trap and the formalism describing the motion (including micromotion) of ion crystals. In Section. III we extend previous studies to characterize the effect of small-amplitude micromotion [19–21] and correct for it in our tweezer patterns, before including first-order Doppler modulation. Section. IV investigates if local stress due to misalignment of the tweezers can improve the optimization and considers the effect of laser intensity fluctuations.

II. TRAPPED-ION QUANTUM SIMULATOR

We consider a one or two dimensional crystal of N ions in a Paul trap. The potential energy of the system is given by $V_0 = V_{\text{coulomb}} + V_{\text{trap}}$. The first term is the contribution due to the Coloumb repulsion between the ions $V_{\text{coulomb}}(\boldsymbol{r}_i) = \frac{1}{2}\sum_{i\neq j} |\boldsymbol{r}_i - \boldsymbol{r}_j|^{-1}$, whilst the second term is the confinement supplied by the external trapping potential

$$V_{\text{trap}}(r_{i,\alpha}, t) = \frac{\Omega_{\text{rf}}^2}{8}\sum_{i,\alpha}[a_\alpha - 2q_\alpha \cos(\Omega_{\text{rf}} t)]r_{i,\alpha}^2, \qquad (1)$$

generated by DC fields and AC components oscillating at Ω_{rf}. Here a and q are the (dimensionless) Mathieu parameters and $r_{i,\alpha}$ is the position of the i-th ion in the $\alpha = x, y, z$ direction. The ion positions and the oscillation frequency are dimensionless and in terms of the characteristic length scale $d = \left(e^2/(4\pi\epsilon_0 m\bar{\omega}^2)\right)^{1/3}$ and a characteristic frequency $\bar{\omega}$ respectively. Here e is the electron charge, ϵ_0 is the vacuum permittivity and m is the ion mass. This allows us to define time t in units of