

Influence in Completely Bounded Block-multilinear Forms and Classical Simulation of Quantum Algorithms

Nikhil Bansal*
University of Michigan

Makrand Sinha†
Simons Institute and UC Berkeley

Ronald de Wolf‡
QuSoft, CWI and U. of Amsterdam

arXiv:2203.00212v1 [quant-ph] 1 Mar 2022

Abstract

The Aaronson-Ambainis conjecture (Theory of Computing '14) says that every low-degree bounded polynomial on the Boolean hypercube has an influential variable. This conjecture, if true, would imply that the acceptance probability of every d-query quantum algorithm can be well-approximated almost everywhere (i.e., on almost all inputs) by a poly(d)-query classical algorithm. We prove a special case of the conjecture: in every completely bounded degree-d block-multilinear form with constant variance, there always exists a variable with influence at least $1/\text{poly}(d)$. In a certain sense, such polynomials characterize the acceptance probability of quantum query algorithms, as shown by Arunachalam, Briët and Palazuelos (SICOMP '19). As a corollary we obtain efficient classical almost-everywhere simulation for a particular class of quantum algorithms that includes for instance k-fold Forrelation. Our main technical result relies on connections to free probability theory.

*bansaln@umich.edu. Supported in part by the NWO VICI grant 639.023.812.

†makrand@berkeley.edu. Supported by a Simons-Berkeley postdoctoral fellowship.

‡rdewolf@cwi.nl. Partially supported by the Dutch Research Council (NWO/OCW), as part of the Quantum Software Consortium programme (project number 024.003.037), and through QuantERA ERA-NET Cofund project QuantAlgo (680-91-034).